

Ann. Funct. Anal. 4 (2013), no. 2, 97–109

ANNALS OF FUNCTIONAL ANALYSIS

ISSN: 2008-8752 (electronic)

URL:www.emis.de/journals/AFA/

## ISHIKAWA TYPE ALGORITHM OF TWO MULTI-VALUED QUASI-NONEXPANSIVE MAPS ON NONLINEAR DOMAINS

HAFIZ FUKHAR-UD-DIN<sup>1,2</sup>, ABDUL RAHIM KHAN<sup>1,\*</sup> AND M. UBAID- UR-REHMAN<sup>2</sup>

Communicated by M. A. Japon Pineda

ABSTRACT. We study an Ishikawa type algorithm for two multi-valued quasinonexpansive maps on a special class of nonlinear spaces namely hyperbolic metric spaces; in particular, strong and  $\triangle$ —convergence theorems for the proposed algorithms are established in a uniformly convex hyperbolic space which improve and extend the corresponding known results in uniformly convex Banach spaces. Our new results are also valid in geodesic spaces.

 $E ext{-}mail\ address: hfdin@kfupm.edu.sa; hfdin@yahoo.com}$ 

E-mail address: arahim@kfupm.edu.sa

<sup>2</sup>Department of Mathematics, The Islamia University of Bahawalpur, Bahawalpur 63100, Pakistan

 $E ext{-}mail\ address: mubaid@188yahoo.com}$ 

<sup>&</sup>lt;sup>1</sup>Department of Mathematics and Statistics, King Fahd University of Petroleum and Minerals, Dhahran 31261, Saudi Arabia.

Date: Received: 12 September 2012; Revised: 11 December 2012; Accepted: 10 January 2013.

<sup>\*</sup> Corresponding author.

<sup>2010</sup> Mathematics Subject Classification. Primary 47H04; Secondary 47H09.

Key words and phrases. Hyperbolic space, multi-valued map, common fixed point, asymptotic centre, convergence.