

ON THE CONVEXITY OF CERTAIN INTEGRAL OPERATORS

VASILE MARIUS MACARIE^{1*} AND DANIEL BREAZ²

Communicated by K. Gurlebeck

ABSTRACT. In this paper we consider the classes of starlike functions of order α , convex functions of order α and we study the convexity and α -order convexity for some general integral operators. Several corollaries of the main results are also considered.

1. INTRODUCTION AND PRELIMINARIES

Let $U = \{z \in \mathbb{C} : |z| < 1\}$ be the unit disk of the complex plane and denote by $H(U)$ the class of the holomorphic functions in U . Consider $\mathcal{A} = \{f \in H(U) : f(z) = z + a_2 z^2 + a_3 z^3 + \dots, z \in U\}$ be the class of analytic functions in U and $S = \{f \in \mathcal{A} : f \text{ is univalent in } U\}$.

Consider S^* the class of starlike functions in unit disk, defined by

$$S^* = \left\{ f \in \mathcal{A} : \operatorname{Re} \left(\frac{zf'(z)}{f(z)} \right) > 0, z \in U \right\}.$$

Definition 1.1. A function $f \in S$ is a *starlike function of order α* , $0 \leq \alpha < 1$ and denote this class by $S^*(\alpha)$ if f verifies the inequality

$$\operatorname{Re} \left(\frac{zf'(z)}{f(z)} \right) > \alpha, \quad (z \in U).$$

Denote with K the class of convex functions in U , defined by

$$K = \left\{ f \in \mathcal{A} : \operatorname{Re} \left(\frac{zf''(z)}{f'(z)} + 1 \right) > 0, z \in U \right\}.$$

Date: Received: 30 January 2012; Accepted: 14 June 2012.

* Corresponding author.

2010 *Mathematics Subject Classification.* Primary 30C45; Secondary 30C55, 47G10.

Key words and phrases. Analytic function, integral operator, open unit disk, convexity, starlike function.

Definition 1.2. A function $f \in S$ is *convex function of order α* , $0 \leq \alpha < 1$ and denote this class by $K(\alpha)$ if f verifies the inequality

$$\operatorname{Re} \left(\frac{zf''(z)}{f'(z)} + 1 \right) > \alpha, \quad (z \in U).$$

Definition 1.3. A function $f \in \mathcal{A}$ is said to be in the class $R(\alpha)$ if $\operatorname{Re}(f'(z)) > \alpha$, $(z \in U)$.

It is well known that $K(\alpha) \subset S^*(\alpha) \subset S$.

Frasin and Jahangiri defined in [1] the family $B(\mu, \alpha)$, $\mu \geq 0$, $0 \leq \alpha < 1$ so that it consists of functions $f \in \mathcal{A}$ satisfying the condition

$$\left| f'(z) \left(\frac{z}{f(z)} \right)^\mu - 1 \right| < 1 - \alpha, \quad (z \in U). \quad (1.1)$$

For $\mu = 0$ we have $B(0, \alpha) \equiv R(\alpha)$ and for $\mu = 1$ we have $B(1, \alpha) \equiv S^*(\alpha)$.

In this paper we will obtain the order of convexity of the following integral operators:

$$H_n(z) = \int_0^z \prod_{i=1}^n (te^{f_i(t)})^{\gamma_i} dt, \quad (1.2)$$

$$H(z) = \int_0^z \prod_{i=1}^n \left(\frac{f_i(t)}{t} \right)^{\frac{1}{\beta}} t^{n(\beta-1)} dt, \quad (1.3)$$

$$G(z) = \int_0^z \prod_{i=1}^n (f_i(t))^{\beta-1} dt, \quad (1.4)$$

where the functions $f_i(t)$ are in $B(\mu_i, \alpha_i)$ for all $i = 1, 2, \dots, n$.

Lemma 1.4. [2] (*General Schwarz lemma*). Let the function f be regular in the disk $U_R = \{z \in \mathbb{C} : |z| < R\}$, with $|f(z)| < M$ for fixed M . If f has one zero with multiplicity order bigger than m for $z = 0$, then

$$|f(z)| \leq \frac{M}{R^m} \cdot |z|^m \quad (z \in U_R).$$

The equality can hold only if

$$f(z) = e^{i\theta} \cdot \frac{M}{R^m} \cdot z^m,$$

where θ is constant.

2. MAIN RESULTS

Theorem 2.1. Let $f_i(z) \in \mathcal{A}$ be in the class $B(\mu_i, \alpha_i)$, $\mu_i \geq 0$, $0 \leq \alpha_i < 1$ for all $i = 1, 2, \dots, n$. If $|f_i(z)| \leq M_i$ ($M_i \geq 1$, $z \in U$) for all $i = 1, 2, \dots, n$, then the integral operator

$$H_n(z) = \int_0^z \prod_{i=1}^n (te^{f_i(t)})^{\gamma_i} dt$$

is in $K(\delta)$, where

$$\delta = 1 - \sum_{i=1}^n |\gamma_i| \cdot [1 + (2 - \alpha_i) M_i^{\mu_i}]$$

and $\sum_{i=1}^n |\gamma_i| \cdot [1 + (2 - \alpha_i) M_i^{\mu_i}] < 1$, $\gamma_i \in \mathbb{C}$ for all $i = 1, 2, \dots, n$.

Proof. Let $f_i \in \mathcal{A}$ be in the class $B(\mu_i, \alpha_i)$, $\mu_i \geq 0$, $0 \leq \alpha_i < 1$. It follows from (1.2) that

$$H_n(z) = \int_0^z t^{\sum_{i=1}^n \gamma_i} e^{\sum_{i=1}^n \gamma_i f_i(t)} dt \quad \text{and} \quad H'_n(z) = z^{\sum_{i=1}^n \gamma_i} e^{\sum_{i=1}^n \gamma_i f_i(z)}.$$

Also

$$H''_n(z) = z^{\sum_{i=1}^n \gamma_i - 1} \cdot e^{\sum_{i=1}^n \gamma_i f_i(z)} \left[\sum_{i=1}^n \gamma_i + z \sum_{i=1}^n \gamma_i f'_i(z) \right]$$

Then

$$\frac{H''_n(z)}{H'_n(z)} = \frac{\sum_{i=1}^n \gamma_i + z \sum_{i=1}^n \gamma_i f'_i(z)}{z}$$

and, hence

$$\begin{aligned} \left| \frac{z H''_n(z)}{H'_n(z)} \right| &= \left| \sum_{i=1}^n \gamma_i + z \sum_{i=1}^n \gamma_i f'_i(z) \right| \leq \sum_{i=1}^n |\gamma_i| + |z| \sum_{i=1}^n |\gamma_i| \cdot |f'_i(z)| \\ &\leq \sum_{i=1}^n |\gamma_i| + |z| \cdot \sum_{i=1}^n |\gamma_i| \cdot \left| f'_i(z) \left(\frac{z}{f_i(z)} \right)^{\mu_i} \right| \cdot \left| \left(\frac{f_i(z)}{z} \right)^{\mu_i} \right| \end{aligned} \quad (2.1)$$

Applying the General Schwarz lemma, we have $\left| \frac{f_i(z)}{z} \right| \leq M_i$, for all $i = 1, 2, \dots, n$.

Therefore, from (2.1), we obtain

$$\left| \frac{z H''_n(z)}{H'_n(z)} \right| \leq \sum_{i=1}^n |\gamma_i| + \sum_{i=1}^n |\gamma_i| \cdot \left| f'_i(z) \left(\frac{z}{f_i(z)} \right)^{\mu_i} \right| \cdot M_i^{\mu_i}, \quad (z \in U). \quad (2.2)$$

From (1.1) and (2.2), we see that

$$\left| \frac{z H''_n(z)}{H'_n(z)} \right| \leq \sum_{i=1}^n |\gamma_i| \cdot [1 + (2 - \alpha_i) M_i^{\mu_i}] = 1 - \delta.$$

This completes the proof. \square

Letting $M_i = M$ for all $i = 1, 2, \dots, n$ in Theorem 2.1, we have

Corollary 2.2. *Let $f_i(z) \in \mathcal{A}$ be in the class $B(\mu_i, \alpha_i)$, $\mu_i \geq 0$, $0 \leq \alpha_i < 1$ for all $i = 1, 2, \dots, n$. If $|f_i(z)| \leq M$ ($M \geq 1$, $z \in U$) for all $i = 1, 2, \dots, n$, then the integral operator*

$$H_n(z) = \int_0^z \prod_{i=1}^n (t e^{f_i(t)})^{\gamma_i} dt$$

is in $K(\delta)$, where

$$\delta = 1 - \sum_{i=1}^n |\gamma_i| \cdot [(2 - \alpha_i)M^{\mu_i} + 1]$$

and $\sum_{i=1}^n |\gamma_i| \cdot [(2 - \alpha_i)M^{\mu_i} + 1] < 1$, $\gamma_i \in \mathbb{C}$ for all $i = 1, 2, \dots, n$.

Letting $\mu_i = 0$ for all $i = 1, 2, \dots, n$ in Corollary 2.2, we have

Corollary 2.3. Let $f_i(z) \in \mathcal{A}$ be in the class $R(\alpha_i)$, $0 \leq \alpha_i < 1$ for all $i = 1, 2, \dots, n$. Then the integral operator defined in (1.2) is in $K(\delta)$, where

$$\delta = 1 - \sum_{i=1}^n |\gamma_i|(3 - \alpha_i)$$

and $\sum_{i=1}^n |\gamma_i|(3 - \alpha_i) < 1$, $\gamma_i \in \mathbb{C}$ for all $i = 1, 2, \dots, n$.

Letting $\mu_i = 1$ for all $i = 1, 2, \dots, n$ in Corollary 2.2, we have

Corollary 2.4. Let $f_i \in \mathcal{A}$ be in the class $S^*(\alpha_i)$, $0 \leq \alpha_i < 1$ for all $i = 1, 2, \dots, n$. If $|f_i(z)| \leq M$ ($M \geq 1$, $z \in U$) for all $i = 1, 2, \dots, n$, then the integral operator defined in (1.2) is in $K(\delta)$, where

$$\delta = 1 - \sum_{i=1}^n |\gamma_i| \cdot [1 + (2 - \alpha_i)M]$$

and $\sum_{i=1}^n |\gamma_i| \cdot [1 + (2 - \alpha_i)M] < 1$, $\gamma_i \in \mathbb{C}$ for all $i = 1, 2, \dots, n$.

Letting $\alpha_i = \delta = 0$ for all $i = 1, 2, \dots, n$ in Corollary 2.4, we have

Corollary 2.5. Let $f_i \in \mathcal{A}$ be starlike functions in U for all $i = 1, 2, \dots, n$. If $|f_i(z)| \leq M$ ($M \geq 1$, $z \in U$) for all $i = 1, 2, \dots, n$ then the integral operator defined in (1.2) is convex in U , where $\sum_{i=1}^n |\gamma_i| = \frac{1}{2M+1}$, $\gamma_i \in \mathbb{C}$ for all $i = 1, 2, \dots, n$.

Theorem 2.6. Let $f_i(z)$ be in the class $B(\mu_i, \alpha_i)$, $\mu_i \geq 1$, $0 \leq \alpha_i < 1$ for all $i = 1, 2, \dots, n$. If $|f_i(z)| \leq M_i$ ($M_i \geq 1$, $z \in U$) for all $i = 1, 2, \dots, n$, then the integral operator

$$H(z) = \int_0^z \prod_{i=1}^n \left(\frac{f_i(t)}{t} \right)^{\frac{1}{\beta}} t^{n(\beta-1)} dt,$$

is in $K(\delta)$, where

$$\delta = 1 - \left\{ \frac{1}{|\beta|} \sum_{i=1}^n [(2 - \alpha_i)M_i^{\mu_i-1} + 1] + n|\beta - 1| \right\}$$

and $\frac{1}{|\beta|} \sum_{i=1}^n [(2 - \alpha_i)M_i^{\mu_i-1} + 1] + n|\beta - 1| < 1$, $\beta \in \mathbb{C} \setminus \{0\}$.

Proof. Let $f_i(z)$ be in the class $B(\mu_i, \alpha_i)$, $\mu_i \geq 1$, $0 \leq \alpha_i < 1$, for all $i = 1, 2, \dots, n$. It follows from (1.3) that

$$\frac{H''(z)}{H'(z)} = \frac{1}{\beta} \cdot \frac{f'_1(z) \cdot z - f_1(z)}{zf_1(z)} + \dots + \frac{1}{\beta} \cdot \frac{f'_n(z) \cdot z - f_n(z)}{zf_n(z)} + \frac{n(\beta - 1)}{z}$$

and

$$\frac{zH''(z)}{H'(z)} = \frac{1}{\beta} \cdot \sum_{i=1}^n \left(\frac{f'_i(z) \cdot z}{f_i(z)} - 1 \right) + n(\beta - 1).$$

So that

$$\begin{aligned} \left| \frac{zH''(z)}{H'(z)} \right| &\leq \frac{1}{|\beta|} \cdot \sum_{i=1}^n \left(\left| \frac{zf'_i(z)}{f_i(z)} \right| + 1 \right) + n|\beta - 1| \\ &\leq \frac{1}{|\beta|} \cdot \sum_{i=1}^n \left(\left| f'_i(z) \left(\frac{z}{f_i(z)} \right)^{\mu_i} \right| \cdot \left| \left(\frac{f_i(z)}{z} \right)^{\mu_i-1} \right| + 1 \right) + n|\beta - 1|. \end{aligned} \quad (2.3)$$

Applying the General Schwarz lemma, we have $\left| \frac{f_i(z)}{z} \right| \leq M_i$, ($z \in U$) for all $i = 1, 2, \dots, n$. Therefore, from (2.3), we obtain

$$\left| \frac{zH''(z)}{H'(z)} \right| \leq \frac{1}{|\beta|} \cdot \sum_{i=1}^n \left(\left| f'_i(z) \left(\frac{z}{f_i(z)} \right)^{\mu_i} \right| \cdot M_i^{\mu_i-1} + 1 \right) + n|\beta - 1|, \quad (z \in U). \quad (2.4)$$

From (1.1) and (2.4), we see that

$$\left| \frac{zH''(z)}{H'(z)} \right| \leq \frac{1}{|\beta|} \sum_{i=1}^n [(2 - \alpha_i)M_i^{\mu_i-1} + 1] + n|\beta - 1| = 1 - \delta.$$

This completes the proof. \square

Letting $M_i = M$ for all $i = 1, 2, \dots, n$ in Theorem 2.6, we have

Corollary 2.7. *Let $f_i(z)$ be in the class $B(\mu_i, \alpha_i)$, $\mu_i \geq 1$, $0 \leq \alpha_i < 1$ for all $i = 1, 2, \dots, n$. If $|f_i(z)| \leq M$ ($M \geq 1$, $z \in U$) for all $i = 1, 2, \dots, n$, then the integral operator*

$$H(z) = \int_0^z \prod_{i=1}^n \left(\frac{f_i(t)}{t} \right)^{\frac{1}{\beta}} t^{n(\beta-1)} dt$$

is in $K(\delta)$, where

$$\delta = 1 - \left\{ \frac{1}{|\beta|} \sum_{i=1}^n [(2 - \alpha_i)M^{\mu_i-1} + 1] + n|\beta - 1| \right\}$$

and $\frac{1}{|\beta|} \sum_{i=1}^n [(2 - \alpha_i)M^{\mu_i-1} + 1] + n|\beta - 1| < 1$, $\beta \in \mathbb{C} \setminus \{0\}$.

Letting $\delta = 0$ in Corollary 2.7 we have

Corollary 2.8. Let $f_i(z)$ be in the class $B(\mu_i, \alpha_i)$, $\mu_i \geq 1$, $0 \leq \alpha_i < 1$ for all $i = 1, 2, \dots, n$. If $|f_i(z)| \leq M$ ($M \geq 1$, $z \in U$) for all $i = 1, 2, \dots, n$, then the integral operator defined in (1.3) is convex function in U , where

$$\frac{1}{|\beta|} \sum_{i=1}^n [(2 - \alpha_i)M^{\mu_i-1} + 1] + n|\beta - 1| = 1, \quad \beta \in \mathbb{C} \setminus \{0\}.$$

Letting $\mu_i = 1$ for all $i = 1, 2, \dots, n$ in Corollary 2.7, we have

Corollary 2.9. Let $f_i(z)$ be in the class $S^*(\alpha_i)$, $0 \leq \alpha_i < 1$ for all $i = 1, 2, \dots, n$. If $|f_i(z)| \leq M$ ($M \geq 1$, $z \in U$) for all $i = 1, 2, \dots, n$, then the integral operator defined in (1.3) is in $K(\delta)$, where

$$\delta = 1 - \left[\frac{1}{|\beta|} \sum_{i=1}^n (3 - \alpha_i) + n|\beta - 1| \right]$$

and $\frac{1}{|\beta|} \sum_{i=1}^n (3 - \alpha_i) + n|\beta - 1| < 1$ for all $\beta \in \mathbb{C} \setminus \{0\}$.

Letting $n = 1$ and $\alpha_i = \delta = 0$ for all $i = 1, 2, \dots, n$ in Corollary 2.9, we have

Corollary 2.10. Let $f(z)$ be a starlike function in U . If $|f(z)| \leq M$ ($M \geq 1$, $z \in U$) then the integral operator $\int_0^z \left(\frac{f(t)}{t} \right)^{\frac{1}{\beta}} t^{\beta-1} dt$ is convex in U , where $\frac{3}{|\beta|} + |\beta - 1| = 1$, $\beta \in \mathbb{C} \setminus \{0\}$.

Theorem 2.11. Let $f_i(z)$ be in the class $B(\mu_i, \alpha_i)$, $\mu_i \geq 1$, $0 \leq \alpha_i < 1$ for all $i = 1, 2, \dots, n$. If $|f_i(z)| \leq M_i$ ($M_i \geq 1$, $z \in U$), for all $i = 1, 2, \dots, n$ then the integral operator

$$G(z) = \int_0^z \prod_{i=1}^n (f_i(t))^{\beta-1} dt$$

is in $K(\delta)$, where

$$\delta = 1 - |\beta - 1| \cdot \sum_{i=1}^n (2 - \alpha_i) M_i^{\mu_i-1}$$

and $|\beta - 1| \cdot \sum_{i=1}^n (2 - \alpha_i) M_i^{\mu_i-1} < 1$, $\beta \in \mathbb{C}$.

Proof. Let $f_i(z)$ be in the class $B(\mu_i, \alpha_i)$, $\mu_i \geq 1$, $0 \leq \alpha_i < 1$ for all $i = 1, 2, \dots, n$. It follows from (1.4) that

$$\frac{G''(z)}{G'(z)} = (\beta - 1) \sum_{i=1}^n \frac{f'_i(z)}{f_i(z)}$$

and, hence

$$\begin{aligned} \left| \frac{zG''(z)}{G'(z)} \right| &\leq |\beta - 1| \left(\sum_{i=1}^n \left| \frac{zf'_i(z)}{f_i(z)} \right| \right) \\ &\leq |\beta - 1| \left(\sum_{i=1}^n \left| f'_i(z) \left(\frac{z}{f_i(z)} \right)^{\mu_i} \right| \cdot \left| \left(\frac{f_i(z)}{z} \right)^{\mu_i-1} \right| \right) \end{aligned} \quad (2.5)$$

Applying the General Schwarz lemma, we have $\left| \frac{f_i(z)}{z} \right| \leq M_i$, ($z \in U$) for all $i = 1, 2, \dots, n$. Therefore, from (2.5), we obtain

$$\left| \frac{zG''(z)}{G'(z)} \right| \leq |\beta - 1| \left(\sum_{i=1}^n \left| f'_i(z) \left(\frac{z}{f_i(z)} \right)^{\mu_i} \right| \cdot M_i^{\mu_i-1} \right), \quad z \in U. \quad (2.6)$$

From (1.1) and (2.6), we see that

$$\left| \frac{zG''(z)}{G'(z)} \right| \leq |\beta - 1| \cdot \sum_{i=1}^n (2 - \alpha_i) M_i^{\mu_i-1} = 1 - \delta.$$

This completes the proof. \square

Letting $M_i = M$ for all $i = 1, 2, \dots, n$ in Theorem 2.11, we have

Corollary 2.12. *Let $f_i(z)$ be in the class $B(\mu_i, \alpha_i)$, $\mu_i \geq 1$, $0 \leq \alpha_i < 1$ for all $i = 1, 2, \dots, n$. If $|f_i(z)| \leq M$ ($M \geq 1$, $z \in U$) then the integral operator*

$$G(z) = \int_0^z \prod_{i=1}^n (f_i(t))^{\beta-1} dt$$

is in $K(\delta)$, where

$$\delta = 1 - |\beta - 1| \sum_{i=1}^n (2 - \alpha_i) M^{\mu_i-1}$$

and $|\beta - 1| \sum_{i=1}^n (2 - \alpha_i) M^{\mu_i-1} < 1$, $\beta \in \mathbb{C}$.

Letting $\delta = 0$ in Corollary 2.12 we have

Corollary 2.13. *Let $f_i(z)$ be in the class $B(\mu_i, \alpha_i)$, $\mu_i \geq 1$, $0 \leq \alpha_i < 1$ for all $i = 1, 2, \dots, n$. If $|f_i(z)| \leq M$ ($M \geq 1$, $z \in U$) then the integral operator defined in (1.4) is convex function in U , where*

$$|\beta - 1| = \frac{1}{\sum_{i=1}^n (2 - \alpha_i) M^{\mu_i-1}}, \quad \beta \in \mathbb{C}.$$

Letting $\mu_i = 1$ for all $i = 1, 2, \dots, n$ in Corollary 2.12, we have

Corollary 2.14. *Let $f_i(z)$ be in the class $S^*(\alpha_i)$, $0 \leq \alpha_i < 1$ for all $i = 1, 2, \dots, n$. If $|f_i(z)| \leq M$ ($M \geq 1$, $z \in U$) then the integral operator defined in (1.4) is in $K(\delta)$, where*

$$\delta = 1 - |\beta - 1| \sum_{i=1}^n (2 - \alpha_i)$$

and $|\beta - 1| \sum_{i=1}^n (2 - \alpha_i) < 1$, $\beta \in \mathbb{C}$.

Letting $n = 1$ and $\alpha_i = \delta = 0$ for all $i = 1, 2, \dots, n$ in Corollary 2.14, we have

Corollary 2.15. *Let $f(z)$ be a starlike function in U . If $|f(z)| \leq M$ ($M \geq 1$, $z \in U$) then the integral operator $\int_0^z f(t)^{\beta-1} dt$ is convex in U , where $|\beta - 1| = \frac{1}{2}$, $\beta \in \mathbb{C}$.*

REFERENCES

1. B.A. Frasin and J. Jahangiri, *A new and comprehensive class of analytic functions*, Anal. Univ. Oradea, Fasc. Math. **15** (2008), 59–62.
2. Z. Nehari, *Conformal Mapping*, Dover, New York, NY, USA, 1975.

¹ DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PITESTI, TARGU DIN VALE STR., NO. 1, 110040, PITESTI, ARGES, ROMANIA.

E-mail address: macariem@yahoo.com

² DEPARTMENT OF MATHEMATICS, "1 DECEMBRIE 1918" UNIVERSITY OF ALBA IULIA, N. IORGA STR., NO. 11-13, 510000, ALBA IULIA, ROMANIA.

E-mail address: dbreaz@uab.ro