

WREATH PRODUCTS IN MODULAR GROUP ALGEBRAS OF SOME FINITE 2-GROUPS

ALEXANDER KONOVALOV

ABSTRACT. Let K be field of characteristic 2 and let G be a finite non-abelian 2-group with the cyclic derived subgroup G' , and there exists a central element z of order 2 in $Z(G)\setminus G'$. We prove that the unit group of the group algebra KG possesses a section isomorphic to the wreath product of a group of order 2 with the derived subgroup of the group G , giving for such groups a positive answer to the question of A. Shalev.

1. INTRODUCTION

Let p be a prime number, G be a finite p -group and K be a field of characteristic p . Denote by $I(KG)$ the augmentation ideal of the modular group algebra KG . The group of normalized units $V(KG)$ consists of all elements of the form $1 + x$, where x belongs to $I(KG)$.

An interest to the structure of the normalized unit group raised a number of questions about kinds of wreath products that may be involved into $V(KG)$ as a subgroup of as a section, i.e. as a factor-group of a certain subgroup of $V(KG)$.

The first result was obtained in [7] by D. Coleman and D. Passman, who proved that for a non-abelian finite p -group G a wreath product of two groups of order p is involved into $V(KG)$. Later it was generalized by A. Bovdi in [2]. Among other related results it is worth to mention [11, 12, 15]. C. Bagiński in [1] described all p -groups, for which $V(KG)$ does not contain a subgroup isomorphic to the wreath product of two groups of order p for the case of odd p . Using results and methods of [5], the case of $p = 2$ was investigated in [3] by V. Bovdi and M. Dokuchaev.

In [14] A. Shalev, motivated by problem of determining of the nilpotency class of $V(KG)$, formulated the question whether $V(KG)$ possesses a section isomorphic to the wreath product of a cyclic group of order p and the derived

2000 *Mathematics Subject Classification.* Primary 16S34, 20C05.

Key words and phrases. wreath product, modular group algebra.

subgroup of G . In [13] he proved that this is true for the case of an odd p and a cyclic derived subgroup of G .

Besides the importance of wreath products with large nilpotency class in the investigation of the unit group, it appears that there are certain connections between wreath products and Lie nilpotency indices of group algebras (see [4, 6]).

When $p = 2$ and G is a 2-group of almost maximal class, in [9] and [10] the author constructed a section of $V(KG)$ isomorphic to the wreath product of a group of order 2 and the derived subgroup of G .

The aim of the present short note is to publish one observation, that allows to confirm the conjecture of A. Shalev for another class of 2-groups.

Theorem. *Let K be field of characteristic 2 and let G be a finite non-abelian 2-group with the cyclic derived subgroup G' , and there exists a central element z of order 2 in $Z(G) \setminus G'$. Then the wreath product of the cyclic group of order 2 and the derived subgroup of G is involved into the $V(KG)$.*

2. PROOF OF THE THEOREM

Proof. Let K be field of characteristic 2 and let G be a finite p -group of order p^n with the cyclic derived subgroup $G' = \langle (b, a) \rangle$ of order 2^s , and there exists an element z of order 2 in $Z(G) \setminus G'$. Let the order of a is p^k and $(b, a^{p^s}) = 1$. Consider an element $h = 1 + b(1 + z)$. Then it is easy to check that

$$\begin{aligned} h^a &= 1 + b^a(1 + z) = 1 + b(b, a)(1 + z), \\ h^{a^2} &= 1 + b^{a^2}(1 + z) = 1 + b(b, a^2)(1 + z), \\ &\dots \\ h^{a^{p^s-1}} &= 1 + b^{a^{p^s-1}}(1 + z) = 1 + b(b, a^{p^s-1})(1 + z), \\ h^{a^{p^s}} &= 1 + b^{a^{p^s}}(1 + z) = h. \end{aligned}$$

Note that h is of order 2 as well as all of its conjugates, and they generate an elementary abelian group $\langle h, h^a, h^{a^2}, \dots, h^{a^{p^s-1}} \rangle$. It is easy to see that this group is actually the direct product

$$X = \langle h \rangle \times \langle h^a \rangle \times \langle h^{a^2} \rangle \times \dots \times \langle h^{a^{p^s-1}} \rangle,$$

since if we multiply its elements, we will obtain an element of the form

$$1 + b((b, a^{i_1}) + (b, a^{i_2}) + \dots + (b, a^{i_t}))(1 + z)$$

which is not equal to 1. Now the wreath product could be obtained if we factorize the semidirect product of X and $\langle a \rangle$ over $\langle a^{p^s} \rangle$, so the theorem is proved. \square

Remark. Note that the family of groups from the conditions of the theorem extends the result of [10] significantly. For example, using the Small Groups

Library of the GAP system [8], we can find that the number of such groups of order 2^n for $n = 4, 5, 6, 7, 8, 9$ is accordingly 4, 20, 72, 231, 662 and 1750.

REFERENCES

- [1] C. Bagiński. Groups of units of modular group algebras. *Proc. Amer. Math. Soc.*, 101(4):619–624, 1987.
- [2] A. A. Bovdi. Construction of a multiplicative group of a group algebra with finiteness conditions. *Mat. Issled.*, (56):14–27, 158, 1980. Algebraic structures.
- [3] V. Bovdi and M. Dokuchaev. Group algebras whose involutory units commute. *Algebra Colloq.*, 9(1):49–64, 2002.
- [4] V. Bovdi, T. Juhász, and E. Spinelli. Modular group algebras with almost maximal Lie nilpotency indices. *Algebr. Represent. Theory*, 9(3):259–266, 2006.
- [5] V. Bovdi and L. G. Kovács. Unitary units in modular group algebras. *Manuscripta Math.*, 84(1):57–72, 1994.
- [6] V. Bovdi and E. Spinelli. Modular group algebras with maximal Lie nilpotency indices. *Publ. Math. Debrecen*, 65(1-2):243–252, 2004.
- [7] D. B. Coleman and D. S. Passman. Units in modular group rings. *Proc. Amer. Math. Soc.*, 25:510–512, 1970.
- [8] The GAP Group. *GAP – Groups, Algorithms, and Programming, Version 4.4.9*, 2006. (<http://www.gap-system.org>).
- [9] A. B. Konovalov. On the nilpotency class of the multiplicative group of a modular group algebra of a dihedral 2-group. *Ukrain. Mat. Zh.*, 47(1):39–45, 1995.
- [10] A. B. Konovalov. Wreath products in the unit group of modular group algebras of 2-groups of maximal class. *Acta Math. Acad. Paedagog. Nyházi. (N.S.)*, 17(3):141–149 (electronic), 2001.
- [11] A. Mann. Wreath products in modular group rings. *Bull. London Math. Soc.*, 23(5):443–444, 1991.
- [12] A. Mann and A. Shalev. The nilpotency class of the unit group of a modular group algebra. II. *Israel J. Math.*, 70(3):267–277, 1990.
- [13] A. Shalev. The nilpotency class of the unit group of a modular group algebra. I. *Israel J. Math.*, 70(3):257–266, 1990.
- [14] A. Shalev. On the conjectures concerning units in p -group algebras. In *Proceedings of the Second International Group Theory Conference (Bressanone, 1989)*, number 23, pages 279–288, 1990.
- [15] A. Shalev. Large wreath products in modular group rings. *Bull. London Math. Soc.*, 23(1):46–52, 1991.

Received March 20, 2007.

SCHOOL OF COMPUTER SCIENCE,
UNIVERSITY OF ST ANDREWS,
NORTH HAUGH, ST ANDREWS,
FIFE, KY16 9SX, SCOTLAND
E-mail address: konovalov@member.ams.org