

# THE CESÀRO $\chi^2$ SEQUENCE SPACES DEFINED BY A MODULUS

N. SUBRAMANIAN

ABSTRACT. In this paper we define the Cesàro  $\chi^2$  sequence space  $\text{Ces}_p^q(\chi_f^2)$  defined by a modulus and exhibit some general properties of the space.

## 1. INTRODUCTION

Throughout  $w, \chi$  and  $\Lambda$  denote the classes of all, gai and analytic scalar valued single sequences, respectively.

We write  $w^2$  for the set of all complex sequences  $(x_{mn})$ , where  $m, n \in \mathbb{N}$ , the set of positive integers. Then,  $w^2$  is a linear space under the coordinate wise addition and scalar multiplication.

An initial work on double sequence spaces is found in Bromwich [4]. Later on, they were investigated by Hardy [5], Moricz [9], Moricz and Rhoades [10], Basarir and Solankan [2], Tripathy [17], Turkmenoglu [19] and many others.

---

Received July 30, 2011.

2010 *Mathematics Subject Classification*. Primary 40A05, 40C05, 40D05.

*Key words and phrases*. Gai sequence; analytic sequence; modulus function; double sequences; Cesàro.



Go back

Full Screen

Close

Quit

Let us define the following sets of double sequences:

$$\mathcal{M}_u(t) := \left\{ (x_{mn}) \in w^2 : \sup_{m,n \in \mathbb{N}} |x_{mn}|^{t_{mn}} < \infty \right\},$$

$$\mathcal{C}_p(t) := \left\{ (x_{mn}) \in w^2 : p - \lim_{m,n \rightarrow \infty} |x_{mn} - l|^{t_{mn}} = 1 \text{ for some } l \in \mathbb{C} \right\},$$

$$\mathcal{C}_{0p}(t) := \left\{ (x_{mn}) \in w^2 : p - \lim_{m,n \rightarrow \infty} |x_{mn}|^{t_{mn}} = 1 \right\},$$

$$\mathcal{L}_u(t) := \left\{ (x_{mn}) \in w^2 : \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} |x_{mn}|^{t_{mn}} < \infty \right\},$$

$$\mathcal{C}_{bp}(t) := \mathcal{C}_p(t) \cap \mathcal{M}_u(t) \quad \text{and} \quad \mathcal{C}_{0bp}(t) = \mathcal{C}_{0p}(t) \cap \mathcal{M}_u(t)$$

where  $t = (t_{mn})$  is the sequence of strictly positive reals  $t_{mn}$  for all  $m, n \in \mathbb{N}$  and  $p - \lim_{m,n \rightarrow \infty}$  denotes the limit in the Pringsheim's sense. In case  $t_{mn} = 1$  for all  $m, n \in \mathbb{N}$ ,  $\mathcal{M}_u(t)$ ,  $\mathcal{C}_p(t)$ ,  $\mathcal{C}_{0p}(t)$ ,  $\mathcal{L}_u(t)$ ,  $\mathcal{C}_{bp}(t)$  and  $\mathcal{C}_{0bp}(t)$  are reduced to the sets  $\mathcal{M}_u$ ,  $\mathcal{C}_p$ ,  $\mathcal{C}_{0p}$ ,  $\mathcal{L}_u$ ,  $\mathcal{C}_{bp}$  and  $\mathcal{C}_{0bp}$ , respectively. Now, we may summarize the knowledge given in some documents related to the double sequence spaces. Gökhan and Colak [21, 22] proved that  $\mathcal{M}_u(t)$  and  $\mathcal{C}_p(t)$ ,  $\mathcal{C}_{bp}(t)$  are complete paranormed spaces of double sequences and gave the  $\alpha$ -,  $\beta$ -,  $\gamma$ - duals of the spaces  $\mathcal{M}_u(t)$  and  $\mathcal{C}_{bp}(t)$ . Quite recently, Zelter [23] in her PhD thesis, essentially studied both the theory of topological double sequence spaces and the theory of summability of double sequences. Mursaleen and Edely [24] recently introduced the statistical convergence and Cauchy for double sequences and gave the relation between statistical convergent and strongly Cesàro summable double sequences. Next, Mursaleen [25] and Mursaleen and Edely [26] defined the almost strong regularity of matrices for double sequences and applied these matrices to establish a core theorem and introduced the  $M$ -core for double sequences and determined those four dimensional matrices transforming every




Go back

Full Screen

Close

Quit



bounded double sequence  $x = (x_{jk})$  into one whose core is a subset of the  $M$ -core of  $x$ . More recently, Altay and Basar [27] defined the spaces  $\mathcal{BS}$ ,  $\mathcal{BS}(t)$ ,  $\mathcal{CS}_p$ ,  $\mathcal{CS}_{bp}$ ,  $\mathcal{CS}_r$  and  $\mathcal{BV}$  of double sequences consisting of all double series whose sequence of partial sums are in the spaces  $\mathcal{M}_u$ ,  $\mathcal{M}_u(t)$ ,  $\mathcal{C}_p$ ,  $\mathcal{C}_{bp}$ ,  $\mathcal{C}_r$  and  $\mathcal{L}_u$ , respectively, and also examined some properties of those sequence spaces and determined the  $\alpha$ -duals of the spaces  $\mathcal{BS}$ ,  $\mathcal{BV}$ ,  $\mathcal{CS}_{bp}$  and the  $\beta(\vartheta)$  – duals of the spaces  $\mathcal{CS}_{bp}$  and  $\mathcal{CS}_r$  of double series. Further Basar and Sever [28] introduced the Banach space  $\mathcal{L}_q$  of double sequences corresponding to the well-known space  $\ell_q$  of single sequences and examined some properties of the space  $\mathcal{L}_q$ . Quite recently Subramanian and Misra [29] studied the space  $\chi_M^2(p, q, u)$  of double sequences and gave some inclusion relations.

Spaces that are strongly summable sequences were discussed by Kuttner [31], Maddox [32], and others. The class of sequences which are strongly Cesàro summable with respect to a modulus was introduced by Maddox [8] as an extension of the definition of strongly Cesàro summable sequences. Connor [33] further extended this definition to a definition of strong  $A$ -summability with respect to a modulus where  $A = (a_{n,k})$  is a nonnegative regular matrix and established some connections among strong  $A$ -summability, strong  $A$ -summability with respect to a modulus, and  $A$ -statistical convergence. In [34] the notion of convergence of double sequences was presented by A. Pringsheim. Also, in [35]–[38] and [39] the four dimensional matrix transformation  $(Ax)_{k,\ell} = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} a_{k\ell}^{mn} x_{mn}$  was studied extensively by Robison and Hamilton. This will be accomplished by presenting the following sequence spaces:

$$\begin{aligned} \text{Ces}_p^q(\chi_f^2) &= d(x, 0) \\ &= \left\{ x \in \chi^2 := \lim_{m,n \rightarrow \infty} \left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( ((m+n) |x_{mn}|)^{\frac{1}{m+n}} \right) \right)^{p_{mn}} \right)^{\frac{1}{p_{mn}}} = 0 \right\} \end{aligned}$$

and




Go back

Full Screen

Close

Quit



$$\text{Ces}_p^q(\Lambda_f^2) = d(x, 0) = \left\{ x \in \chi^2 := \sup \left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( (|x_{mn}|)^{\frac{1}{m+n}} \right) \right)^{p_{mn}} \right)^{\frac{1}{p_{mn}}} < \infty \right\}$$

where  $f$  is a modulus function. Other implications, general properties and variations will also be presented.

In the sequel of the paper we need the following inequality

$$(1.1) \quad (a + b)^p \leq a^p + b^p$$

for  $a, b, \geq 0$  and  $0 < p < 1$ . The double series  $\sum_{m,n=1}^{\infty} x_{mn}$  is called convergent if and only if the double sequence  $(s_{mn})$  is convergent, where  $s_{mn} = \sum_{i,j=1}^{m,n} x_{ij}$  ( $m, n \in \mathbb{N}$ ) (see [1]).

A sequence  $x = (x_{mn})$  is said to be double analytic if  $\sup_{m,n} |x_{mn}|^{1/m+n} < \infty$ . The vector space of all double analytic sequences will be denoted by  $\Lambda^2$ . A sequence  $x = (x_{mn})$  is called double gai sequence if  $((m+n)! |x_{mn}|)^{1/m+n} \rightarrow 0$  as  $m, n \rightarrow \infty$ . The double gai sequences will be denoted by  $\chi^2$ . Let  $\phi = \{\text{all finite sequences}\}$ .

Consider a double sequence  $x = (x_{ij})$ . The  $(m, n)^{th}$  section  $x^{[m,n]}$  of the sequence is defined by  $x^{[m,n]} = \sum_{i,j=0}^{m,n} x_{ij} \mathfrak{S}_{ij}$  for all  $m, n \in \mathbb{N}$ ; where  $\mathfrak{S}_{ij}$  denotes the double sequence whose only non zero term is a  $\frac{1}{(i+j)!}$  in the  $(i, j)^{th}$  place for each  $i, j \in \mathbb{N}$ .

An FK-space (or a metric space)  $X$  is said to have AK property if  $(\mathfrak{S}_{mn})$  is a Schauder basis for  $X$ . Or equivalently  $x^{[m,n]} \rightarrow x$ .

An FDK-space is a double sequence space endowed with a complete metrizable; locally convex topology under which the coordinate mappings  $x = (x_k) \rightarrow (x_{mn})$  ( $m, n \in \mathbb{N}$ ) are also continuous.

Orlicz [13] used the idea of Orlicz function to construct the space  $(L^M)$ . Lindenstrauss and Tzafriri [7] investigated Orlicz sequence spaces in more detail and proved that every Orlicz sequence space  $\ell_M$  contains a subspace isomorphic to  $\ell_p$  ( $1 \leq p < \infty$ ). Subsequently, different classes of




Go back

Full Screen

Close

Quit



sequence spaces were defined by Parashar and Choudhary [14], Mursaleen et al. [11], Bektas and Altin [3], Tripathy et al. [18], Rao and Subramanian [15] and many others. The Orlicz sequence spaces are the special cases of Orlicz spaces studied in [6].

Recalling [13] and [6], an Orlicz function is a function  $M: [0, \infty) \rightarrow [0, \infty)$  which is continuous, non-decreasing and convex with  $M(0) = 0$ ,  $M(x) > 0$  for  $x > 0$  and  $M(x) \rightarrow \infty$  as  $x \rightarrow \infty$ . If convexity of Orlicz function  $M$  is replaced by subadditivity of  $M$ , then this function is called modulus function, defined by Nakano [12] and further discussed by Ruckle [16] and Maddox [8], and many others.

An Orlicz function  $M$  is said to satisfy the  $\Delta_2$ -condition for all values of  $u$  if there exists a constant  $K > 0$  such that  $M(2u) \leq KM(u)$  ( $u \geq 0$ ). The  $\Delta_2$ -condition is equivalent to  $M(\ell u) \leq K\ell M(u)$  for all values of  $u$  and for  $\ell > 1$ .

Lindenstrauss and Tzafriri [7] used the idea of Orlicz function to construct Orlicz sequence space

$$\ell_M = \left\{ x \in w : \sum_{k=1}^{\infty} M\left(\frac{|x_k|}{\rho}\right) < \infty, \text{ for some } \rho > 0 \right\}.$$

The space  $\ell_M$  with the norm

$$\|x\| = \inf \left\{ \rho > 0 : \sum_{k=1}^{\infty} M\left(\frac{|x_k|}{\rho}\right) \leq 1 \right\}$$

becomes a Banach space which is called an Orlicz sequence space. For  $M(t) = t^p$ , ( $1 \leq p < \infty$ ), the spaces  $\ell_M$  coincide with the classical sequence space  $\ell_p$ .

If  $X$  is a sequence space, we give the following definitions:

- (i)  $X'$  is the continuous dual of  $X$ ;



Go back

Full Screen

Close

Quit

- (ii)  $X^\alpha = \left\{ a = (a_{mn}) : \sum_{m,n=1}^{\infty} |a_{mn}x_{mn}| < \infty, \text{ for each } x \in X \right\};$
- (iii)  $X^\beta = \left\{ a = (a_{mn}) : \sum_{m,n=1}^{\infty} a_{mn}x_{mn} \text{ is convergent, for each } x \in X \right\};$
- (iv)  $X^\gamma = \left\{ a = (a_{mn}) : \sup_{mn} \geq 1, \left| \sum_{m,n=1}^{M,N} a_{mn}x_{mn} \right| < \infty, \text{ for each } x \in X \right\};$
- (v) let  $X$  be an FK-space  $\supset \phi$ ; then  $X^f = \left\{ f(\mathfrak{I}_{mn}) : f \in X' \right\};$
- (vi)  $X^\delta = \left\{ a = (a_{mn}) : \sup_{mn} |a_{mn}x_{mn}|^{1/m+n} < \infty, \text{ for each } x \in X \right\};$

$X^\alpha, X^\beta, X^\gamma$  are called  $\alpha$ - (or Köthe-Toeplitz)-dual of  $X$ ,  $\beta$ - (or generalized- Köthe-Toeplitz)-dual of  $X$ ,  $\gamma$ -dual of  $X$ ,  $\delta$ -dual of  $X$ , respectively.  $X^\alpha$  was defined by Gupta and Kamptan [20]. It is clear that  $x^\alpha \subset X^\beta$  and  $X^\alpha \subset X^\gamma$ , but  $X^\beta \subset X^\gamma$  does not hold, since the sequence of partial sums of a double convergent series need not to be bounded.

The notion of difference sequence spaces (for single sequences) was introduced by Kizmaz [30] as follows

$$Z(\Delta) = \{x = (x_k) \in w : (\Delta x_k) \in Z\}$$

for  $Z = c, c_0$  and  $\ell_\infty$ , where  $\Delta x_k = x_k - x_{k+1}$  for all  $k \in \mathbb{N}$ .

Here  $c, c_0$  and  $\ell_\infty$  denote the classes of convergent, null and bounded scalar valued single sequences, respectively. The difference space  $bv_p$  of the classical space  $\ell_p$  was introduced and studied in the case  $1 \leq p \leq \infty$  by Basar and Altay in [42] and in the case  $0 < p < 1$  by Altay and Basar




Go back

Full Screen

Close

Quit



in [43]. The spaces  $c(\Delta)$ ,  $c_0(\Delta)$ ,  $\ell_\infty(\Delta)$  and  $bv_p$  are Banach spaces normed by

$$\|x\| = |x_1| + \sup_{k \geq 1} |\Delta x_k| \quad \text{and} \quad \|x\|_{bv_p} = \left( \sum_{k=1}^{\infty} |x_k|^p \right)^{1/p}, \quad (1 \leq p < \infty).$$

Later on the notion was further investigated by many others. We now introduce the following difference double sequence spaces defined by

$$Z(\Delta) = \{x = (x_{mn}) \in w^2 : (\Delta x_{mn}) \in Z\}$$

where  $Z = \Lambda^2$ ,  $\chi^2$  and  $\Delta x_{mn} = (x_{mn} - x_{mn+1}) - (x_{m+1n} - x_{m+1n+1}) = x_{mn} - x_{mn+1} - x_{m+1n} + x_{m+1n+1}$  for all  $m, n \in \mathbb{N}$ .

## 2. DEFINITIONS AND PRELIMINARIES

$\text{Ces}_p^q(\chi_f^2)$  and  $\text{Ces}_p^q(\Lambda_f^2)$  denote the Pringscheims sense of Cesàro double gai sequence space of modulus and Pringscheims sense of Cesàro double analytic sequence space of modulus, respectively.

**Definition 2.1.** A modulus function was introduced by Nakano [12]. We recall that a modulus  $f$  is a function from  $[0, \infty) \rightarrow [0, \infty)$  such that

- (1)  $f(x) = 0$  if and only if  $x = 0$ ,
- (2)  $f(x + y) \leq f(x) + f(y)$  for all  $x \geq 0, y \geq 0$ ,
- (3)  $f$  is increasing,
- (4)  $f$  is continuous from the right at 0. Since  $|f(x) - f(y)| \leq f(|x - y|)$ , it follows from here that  $f$  is continuous on  $[0, \infty)$ .




Go back

Full Screen

Close

Quit



**Definition 2.2.** Let  $A = (a_{k\ell}^{mn})$  denote a four dimensional summability method that maps the complex double sequences  $x$  into the double sequence  $Ax$  where the  $k, \ell$ -th term to  $Ax$  is as follows:

$$(Ax)_{k\ell} = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} a_{k\ell}^{mn} x_{mn}.$$

Such transformation is said to be nonnegative if  $a_{k\ell}^{mn}$  is nonnegative.

The notion of regularity for two dimensional matrix transformations was presented by Silverman [40] and Toeplitz [41]. Following Silverman and Toeplitz, Robison and Hamilton presented the following four dimensional analog of regularity for double sequences in which they both added an additional assumption of boundedness. This assumption was made because a double sequence which is  $P$ -convergent is not necessarily bounded.

**Definition 2.3.** Let  $p \in [1, \infty)$  and  $q$  be a double gai sequence of positive real numbers such that

$$Q_{ij} = \sum_{m=0}^i \sum_{n=0}^j q_{mn}, \quad i, j \in \mathbb{N},$$

$$\text{Ces}_p^q(\chi_f^2) = d(x, 0)$$

$$= \left\{ x \in \chi^2 := \lim_{m, n \rightarrow \infty} \left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( ((m+n) |x_{mn}|)^{\frac{1}{m+n}} \right) \right)^{p_{mn}} \right)^{\frac{1}{p_{mn}}} = 0 \right\}$$

If  $q_{mn} = 1$  for all  $m, n \in \mathbb{N}$ , then  $\text{Ces}_p^q(\chi_f^2)$  reduces to  $\text{Ces}_p(\chi_f^2)$ , and if  $f(x) = x$ , then  $\text{Ces}_p^q(\chi_f^2)$  reduces to  $\text{Ces}_p^q(\chi^2)$ .




Go back

Full Screen

Close

Quit



**Definition 2.4.** Let  $p \in [1, \infty)$  and  $q$  be a double analytic sequence of positive real numbers such that

$$Q_{ij} = \sum_{m=0}^i \sum_{n=0}^j q_{mn}, \quad i, j \in \mathbb{N},$$

$$\begin{aligned} \text{Ces}_p^q(\Lambda_f^2) &= d(x, 0) \\ &= \left\{ x \in \chi^2 := \sup \left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( (|x_{mn}|)^{\frac{1}{m+n}} \right) \right)^{p_{mn}} \right)^{\frac{1}{p_{mn}}} < \infty \right\}. \end{aligned}$$

If  $q_{mn} = 1$  for all  $m, n \in \mathbb{N}$ , then  $\text{Ces}_p^q(\Lambda_f^2)$  reduces to  $\text{Ces}_p(\Lambda_f^2)$ , and if  $f(x) = x$ , then  $\text{Ces}_p^q(\Lambda_f^2)$  reduces to  $\text{Ces}_p^q(\Lambda^2)$ .

The space  $\text{Ces}_p^q(\chi_f^2)$  is a metric space with the metric

$$\begin{aligned} d(x, y) &= \inf \left\{ \sup \left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( ((m+n)! |x_{mn} - y_{mn}|)^{\frac{1}{m+n}} \right) \right)^{p_{mn}} \right)^{\frac{1}{p_{mn}}} \leq 1 \right\} \end{aligned}$$



Go back

Full Screen

Close

Quit



The space  $Ces_p^q(\Lambda_f^2)$  is a metric space with the metric

$$d(x, y) = \inf \left\{ \sup \left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( (|x_{mn} - y_{mn}|)^{\frac{1}{m+n}} \right) \right)^{p_{mn}} \right)^{\frac{1}{p_{mn}}} \leq 1 \right\}.$$

### 3. MAIN RESULTS

**Proposition 3.1.** *Let  $x, y \in Ces_p^q(\chi_f^2)$ . Then for any  $\varepsilon > 0$  and  $L > 0$ , there exists  $\delta > 0$  such that  $(d(x + y, 0), 0)^{p_{mn}} = d(x, 0)^{p_{mn}} + \varepsilon$ , whenever  $d(x, 0)^p \leq L$  and  $d(y, 0)^{p_{mn}} \leq \delta$ .*

*Proof.* For any fix  $\varepsilon > 0$ ,

$$\begin{aligned} d(x + y, 0)^{p_{mn}} &= \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( ((m+n)! |x_{mn} + y_{mn}|)^{\frac{1}{m+n}} \right) \right)^{p_{mn}} \\ &\leq \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( ((m+n)! |x_{mn}|)^{\frac{1}{m+n}} \right) \right)^{p_{mn}} \\ &\quad + \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( ((m+n)! |y_{mn}|)^{\frac{1}{m+n}} \right) \right)^{p_{mn}} \end{aligned}$$



Go back

Full Screen

Close

Quit

$$\begin{aligned}
&\leq (1 - \beta) \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( ((m+n)! |x_{mn}|)^{\frac{1}{m+n}} \right) \right)^{P_{mn}} \\
&\quad + (\beta) \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( ((m+n)! |x_{mn}|)^{\frac{1}{m+n}} \right) \right)^{P_{mn}} \\
&\quad \cdot (\beta) \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j \frac{q_{mn} f \left( ((m+n)! |y_{mn}|)^{\frac{1}{m+n}} \right)}{\beta} \right)^{P_{mn}} \\
&\leq \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( ((m+n)! |x_{mn}|)^{\frac{1}{m+n}} \right) \right)^{P_{mn}} \\
&\quad + \frac{\beta}{2} \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j 2q_{mn} f \left( ((m+n)! |x_{mn}|)^{\frac{1}{m+n}} \right) \right)^{P_{mn}} \\
&\quad \cdot \frac{\beta}{2} \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j \frac{2q_{mn} f \left( ((m+n)! |y_{mn}|)^{\frac{1}{m+n}} \right)}{\beta} \right)^{P_{mn}} \\
&\leq d(x, 0)^{P_{mn}} + \frac{\varepsilon}{2} \\
&\quad + \left( \frac{2}{\beta} \right)^{P_{mn}-1} \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( ((m+n)! |y_{mn}|)^{\frac{1}{m+n}} \right) \right)^{P_{mn}} \\
&\leq d(x, 0)^{P_{mn}} + \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \\
&\leq d(x, 0)^{P_{mn}} + \varepsilon.
\end{aligned}$$



Go back

Full Screen

Close

Quit

**Proposition 3.2.** For every  $p = (p_{mn})$ ,

$$[Ces_p^q(\Lambda_f^2)]^\beta = [Ces_p^q(\Lambda_f^2)]^\alpha = [Ces_p^q(\Lambda_f^2)]^\gamma = [Ces_p^q(\eta_f^2)]^\beta,$$

where

$$[Ces_p^q(\eta_f^2)] = \bigcap_{N \in \mathbb{N} - \{1\}} \left\{ x = x_{mn} : \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f(|x_{mn}| N^{\frac{m+n}{p_{mn}}}) \right)^{\frac{1}{p_{mn}}} < \infty \right\}.$$

*Proof.* First we show that  $[Ces_p^q(\eta_f^2)] \subset [Ces_p^q(\Lambda_f^2)]^\beta$ .

Let  $x \in [Ces_p^q(\eta_f^2)]$  and  $y \in [Ces_p^q(\Lambda_f^2)]^\beta$ . Then we can find a positive integer  $N$  such that

$$\begin{aligned} & (|y_{mn}|^{1/m+n}) \\ & < \max \left( 1, \sup_{m,n \geq 1} \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f(|y_{mn}|^{\frac{1}{m+n}})^{p_{mn}} \right)^{\frac{1}{p_{mn}}} \right) < N \end{aligned}$$

for all  $m, n$ .

Hence we may write

$$\left| \sum_{m,n} x_{mn} y_{mn} \right| \leq \sum_{m,n} |x_{mn} y_{mn}| \leq \sum_{m,n} (f(|x_{mn} y_{mn}|)) \leq \sum_{m,n} (f(|x_{mn}| N^{m+n})).$$



Go back

Full Screen

Close

Quit

Since  $x \in Ces_p^q(\eta_f^2)$ . The series on the right side of the above inequality is convergent, whence  $x \in Ces_p^q(\Lambda_f^2)$ . Hence  $[Ces_p^q(\eta_f^2)] \subset [Ces_p^q(\Lambda_f^2)]^\beta$ .

Now we show that  $[Ces_p^q(\Lambda_f^2)]^\beta \subset [Ces_p^q(\eta_f^2)]$ .

For this, let  $x \in [Ces_p^q(\Lambda_f^2)]^\beta$  and suppose that  $x \notin [Ces_p^q(\eta_f^2)]$ . Then there exists a positive integer  $N > 1$  such that  $\sum_{m,n} (f(|x_{mn}| N^{m+n})) = \infty$ .

If we define  $y_{mn} = N^{m+n} \text{Sgn} x_{mn}$   $m, n = 1, 2, \dots$ , then  $y \in [Ces_p^q(\Lambda_f^2)]$ .

But, since

$$\left| \sum_{m,n} x_{mn} y_{mn} \right| = \sum_{m,n} (f(|x_{mn} y_{mn}|)) = \sum_{m,n} (f(|x_{mn}| N^{m+n})) = \infty,$$

we get  $x \notin [Ces_p^q(\Lambda_f^2)]^\beta$ , which contradicts the assumption  $x \in [Ces_p^q(\Lambda_f^2)]^\beta$ . Therefore  $x \in [Ces_p^q(\eta_f^2)]$  and  $[Ces_p^q(\Lambda_f^2)]^\beta = [Ces_p^q(\eta_f^2)]$ .

(ii) and (iii) can be shown in a similar way of (i). Therefore, we omit it.  $\square$

**Proposition 3.3.** *Let  $p = (p_{mn})$  be a Cesàro space of double analytic modulus sequence of strictly positive real numbers  $p_{mn}$ . Then*

(i)  $Ces_p^q(\Lambda_f^2)$  is a paranormed space with

$$(3.1) \quad g(x) = \sup \left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f(|x_{mn}|^{\frac{1}{m+n}}) \right)^{\frac{p_{mn}}{M}} \right)^{\frac{1}{p_{mn}}}$$




Go back

Full Screen

Close

Quit



if and only if  $h = \inf p_{mn} > 0$ , where  $M = \max(1, H)$  and  $H = \sup p_{mn}$ .

(ii)  $\text{Ces}_p^q(\Lambda_f^2)$  is a complete paranormed linear metric space if the condition  $p$  in (3.1) is satisfied.

*Proof.* The proof of (i). *Sufficiency.* Let  $h > 0$ . It is trivial that  $g(\theta) = 0$  and  $g(-x) = g(x)$ .

The inequality  $g(x + y) \leq g(x) + g(y)$  follows from the inequality (3.1), since  $p_{mn}/M \leq 1$  for all positive integers  $m, n$ . We also may write  $g(\lambda x) \leq \max(|\lambda|, |\lambda|^{h/M}) g(x)$ , since  $|\lambda|^{p_{mn}} \leq \max(|\lambda|^h, |\lambda|^M)$  for all positive integers  $m, n$  and for any  $\lambda \in \mathbb{C}$ , the set of complex numbers. Using this inequality, it can be proved that  $\lambda x \rightarrow \theta$ , when  $x$  is fixed and  $\lambda \rightarrow 0$ , or  $\lambda \rightarrow 0$  and  $x \rightarrow \theta$ , or  $\lambda$  is fixed and  $x \rightarrow \theta$ .

*Necessity.* Let  $\text{Ces}_p^q(\Lambda_f^2)$  be a paranormed space with the paranorm

$$g(x) = \sup \left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( |x_{mn}|^{\frac{1}{m+n}} \right) \right)^{\frac{p_{mn}}{M}} \right)^{\frac{1}{p_{mn}}}$$

and suppose that  $h = 0$ . Since  $|\lambda|^{p_{mn}/M} \leq |\lambda|^{h/M} = 1$  for all positive integers  $m, n$  and  $\lambda \in \mathbb{C}$  such that  $0 < |\lambda| \leq 1$ , we have

$$\sup \left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f(|\lambda|) \right)^{\frac{p_{mn}}{M}} \right)^{\frac{1}{p_{mn}}} = 1.$$




Go back

Full Screen

Close

Quit



Hence it follows that

$$g(\lambda x) = \sup \left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f(|\lambda|) \right)^{\frac{p_{mn}}{M}} \right)^{\frac{1}{p_{mn}}} = 1$$

for  $x = (\alpha) \in \text{Ces}_p^q(\Lambda_f^2)$  as  $\lambda \rightarrow 0$ . But this contradicts the assumption  $\text{Ces}_p^q(\Lambda_f^2)$  is a paranormed space with  $g(x)$ .

The proof of (ii) is clear. □

**Corollary 3.4.**  $\text{Ces}_p^q(\Lambda_f^2)$  is a complete paranormed space with the natural paranorm if and only if  $\text{Ces}_p^q(\Lambda_f^2) = \text{Ces}^q(\Lambda_f^2)$ .

**Proposition 3.5.** For every  $p = (p_{mn})$ ,  $\text{Ces}_p^q(\eta_f^2) \subset [\text{Ces}_p^q(\chi_f^2)]^\beta \subsetneq \text{Ces}_p^q(\Lambda_f^2)$ .

*Proof.* The proof of (i). First, we show that  $\text{Ces}_p^q(\eta_f^2) \subset [\text{Ces}_p^q(\chi_f^2)]^\beta$ . We know that  $[\text{Ces}_p^q(\chi_f^2)] \subset \text{Ces}_p^q(\Lambda_f^2)$ .  $[\text{Ces}_p^q(\Lambda_f^2)]^\beta \subset [\text{Ces}_p^q(\chi_f^2)]^\beta$ . But  $[\text{Ces}_p^q(\Lambda_f^2)]^\beta = \text{Ces}_p^q(\eta_f^2)$  by Proposition 3.2. Therefore,

$$(3.2) \quad \text{Ces}_p^q(\eta_f^2) \subset [\text{Ces}_p^q(\chi_f^2)]^\beta.$$

The proof of (ii). Now we show that  $[\text{Ces}_p^q(\chi_f^2)]^\beta \subsetneq \text{Ces}_p^q(\Lambda_f^2)$ .




Go back

Full Screen

Close

Quit



Let  $y = (y_{mn})$  be an arbitrary point  $[\text{Ces}_p^q(\chi_f^2)]^\beta$ . If  $y$  is not  $\text{Ces}_p^q(\Lambda_f^2)$ , then for each natural number  $q$ , we can find an index  $m_p n_q$  such that

$$\left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{m_q n_q} f \left( ((m_q + n_q)! |y_{m_q n_q}|)^{\frac{1}{m_q + n_q}} \right) \right)^{p_{mn}} \right)^{\frac{1}{p_{mn}}} > q$$

for  $(1, 2, 3, \dots)$ . Define  $x = \{x_{mn}\}$  by

$$\left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( ((m + n)! |x_{mn}|)^{p_{mn}} \right)^{\frac{1}{p_{mn}}} \right) \right) q^{\frac{1}{m+n}}$$

for  $(mn) = (m_q n_q)$  and some  $q \in \mathbb{N}$ ; and

$$\left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( ((m + n)! |x_{mn}|)^{p_{mn}} \right)^{\frac{1}{p_{mn}}} \right) \right) = 0, \text{ otherwise.}$$

Then  $x$  is  $\text{Ces}_p^q(\chi_f^2)$ , but for infinitely  $mn$ ,

$$(3.3) \quad \left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( ((m + n)! |y_{mn} x_{mn}|)^{p_{mn}} \right)^{\frac{1}{p_{mn}}} \right) \right) > 1.$$

Consider the sequence  $z = \{z_{mn}\}$ , where

$$Q_{11} (q_{11} f(2! z_{11})^{p_{mn}})^{p_{mn}} = Q_{11} (q_{11} f(2! x_{11})^{p_{mn}})^{p_{mn}} - s$$




Go back

Full Screen

Close

Quit



with

$$\begin{aligned}
 s &= \left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( ((m+n)! |x_{mn}|)^{p_{mn}} \frac{1}{p_{mn}} \right) \right) \right); \\
 &\left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( ((m+n)! |z_{mn}|)^{p_{mn}} \frac{1}{p_{mn}} \right) \right) \right) \\
 &= \left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( ((m+n)! |x_{mn}|)^{p_{mn}} \frac{1}{p_{mn}} \right) \right) \right).
 \end{aligned}$$

The  $z$  is a point of  $\text{Ces}_p^q(\chi_f^2)$ . Also

$$\left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( ((m+n)! |z_{mn}|)^{p_{mn}} \frac{1}{p_{mn}} \right) \right) \right) = 0.$$

Hence  $z$  is in  $\text{Ces}_p^q(\chi_f^2)$ . But, by the equation (3.3),

$$\left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( ((m+n)! |z_{mn} y_{mn}|)^{p_{mn}} \frac{1}{p_{mn}} \right) \right) \right)$$

does not converge and so  $\sum \sum x_{mn} y_{mn}$  diverges. Thus, the sequence  $y$  would not be  $[\text{Ces}_p^q(\chi_f^2)]^\beta$ .

This contradiction proves that




Go back

Full Screen

Close

Quit



$$(3.4) \quad [\text{Ces}_p^q(\chi_f^2)]^\beta \subset \text{Ces}_p^q(\Lambda_f^2).$$

If we now choose  $f = \text{id}$ , where  $\text{id}$  is the identity and

$$\frac{1}{Q_{1j}}(q_{1n}((m+n)!y_{1n})) = \frac{1}{Q_{1j}}(q_{1n}((m+n)!x_{1n}))$$

and

$$\frac{1}{Q_{ij}}(q_{mn}((m+n)!y_{mn})) = \frac{1}{Q_{ij}}(q_{mn}((m+n)!x_{mn})) = 0, \quad (m, i > 1)$$

for all  $n, j$ , then obviously  $x \in \text{Ces}_p^q(\chi_f^2)$  and  $y \in \text{Ces}_p^q(\Lambda_f^2)$ , but

$$(3.5) \quad \sum \sum x_{mn}y_{mn} = \infty.$$

Hence  $y \notin [\text{Ces}_p^q(\chi_f^2)]^\beta$ .

From (3.4) and (3.5), we are granted  $[\text{Ces}_p^q(\chi_f^2)]^\beta \not\subseteq \text{Ces}_p^q(\Lambda_f^2)$ . □

**Proposition 3.6.** *In  $\text{Ces}_p^q(\chi_f^2)$  weak convergence does not imply strong convergence.*

*Proof.* Assume that weak convergence implies strong convergence  $\text{Ces}_p^q(\chi_f^2)$ . Then, we would have  $[\text{Ces}_p^q(\chi_f^2)]^{\beta\beta} = \text{Ces}_p^q(\chi_f^2)$  [see Wilansky]. But

$$[\text{Ces}_p^q(\chi_f^2)]^{\beta\beta} \not\subseteq [\text{Ces}_p^q(\Lambda_f^2)]^\beta = \text{Ces}_p^q(\eta_f^2).$$




Go back

Full Screen

Close

Quit



Thus  $[\text{Ces}_p^q(\chi_f^2)]^{\beta\beta} \neq \text{Ces}_p^q(\chi_f^2)$ . Hence weak convergence does not imply strong convergence in  $\text{Ces}_p^q(\chi_f^2)$ .  $\square$

**Proposition 3.7.** *Let  $f$  be an modulus function which satisfies the  $\Delta_2$ -condition. Then  $\text{Ces}_p^q(\chi^2) \subset \text{Ces}_p^q(\chi_f^2)$ .*

*Proof.* Let

$$(3.6) \quad x \in \text{Ces}_p^q(\chi^2).$$

Then

$$\left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} ((m+n)! |x_{mn}|)^{p_{mn}} \right)^{\frac{1}{p_{mn}}} \right) \leq \varepsilon$$

for sufficiently large  $m, n$  and every  $\varepsilon > 0$ .

$$\left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f(((m+n)! |x_{mn}|)^{p_{mn}})^{\frac{1}{p_{mn}}} \right) \right) \leq f(\varepsilon)$$

(because  $f$  is non-decreasing). This implies

$$\left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f(((m+n)! |x_{mn}|)^{p_{mn}})^{\frac{1}{p_{mn}}} \right) \right) \leq Kf(\varepsilon) < (\varepsilon)$$




Go back

Full Screen

Close

Quit



(by the  $\Delta_2$ -condition, for some  $K > 0$  and by defining  $f(\varepsilon) < \frac{\varepsilon}{K}$ ).

$$(3.7) \quad \lim_{m,n \rightarrow \infty} \left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( ((m+n)! |x_{mn}|)^{\frac{1}{m+n}} \right) \right)^{p_{mn}} \right)^{\frac{1}{p_{mn}}} = 0.$$

Hence

$$(3.8) \quad x \in \text{Ces}_p^q(\chi_f^2).$$

From (3.6) and (3.8), we get  $\text{Ces}_p^q(\chi^2) \subset \text{Ces}_p^q(\chi_f^2)$ . □

**Proposition 3.8.**  $[\text{Ces}_p^q(\Lambda_f^2)]^\beta \subsetneq \text{Ces}_p^q(\chi_f^2)$ .

*Proof.* Let  $(x_{mn}) \in [\text{Ces}_p^q(\Lambda_f^2)]^\beta$

$$(3.9) \quad \sum \sum |x_{mn} y_{mn}| < \infty$$

for all  $(y_{mn}) \in [\text{Ces}_p^q(\Lambda_f^2)]^\beta$ . Assume that  $(x_{mn}) \notin \text{Ces}_p^q(\chi_f^2)$ . Then there exists a sequence of positive integers

$$f(|x_{m_r+n_r}|) > \frac{1}{((m_r+n_r)! 2^{(m_r+n_r)})}, \quad (r = 1, 2, 3, \dots).$$

Take

$$y_{m_r+n_r} = \begin{cases} (2(m_r+n_r)!)^{m_r+n_r} & \text{for } r = 1, 2, 3, \dots, \\ y_{m_r+n_r} = 0 & \text{otherwise.} \end{cases}$$



Go back

Full Screen

Close

Quit



Then  $(y_{mn}) \in [\text{Ces}_p^q(\Lambda_f^2)]$ . But

$$\sum \sum |x_{mn}y_{mn}| = \sum_{r=1}^{\infty} |x_{m_r+n_r}y_{m_r+n_r}| = f \left( \sum_{r=1}^{\infty} |x_{m_r+n_r}y_{m_r+n_r}| \right) > 1 + 1 + 1 + \dots .$$

We know that the infinite series  $1 + 1 + 1 + \dots$  diverges. Hence  $\sum \sum |x_{mn}y_{mn}|$  diverges. This contradicts (3.9). Hence  $(x_{mn}) \in \text{Ces}_p^q(\chi_f^2)$ . Therefore,

$$(3.10) \quad [\text{Ces}_p^q(\Lambda_f^2)]^\beta \subset \text{Ces}_p^q(\chi_f^2).$$

If we now choose  $p = (p_{mn})$ , it is a constant  $f = \text{id}$ , where  $\text{id}$  is the identity and

$$\frac{1}{Q_{1j}} (q_{1n} ((1+n)!y_{1n})) = \frac{1}{Q_{1j}} (q_{1n} ((1+n)!x_{1n})) \quad \text{and} \\ \frac{1}{Q_{ij}} (q_{mn} ((m+n)!y_{mn})) = \frac{1}{Q_{ij}} (q_{mn} ((m+n)!x_{mn})) = 0$$

where  $(m, i > 1)$  for all  $n, j$ , then obviously  $x \in \text{Ces}_p^q(\chi_f^2)$  and  $y \in \text{Ces}_p^q(\Lambda_f^2)$ , but

$$(3.11) \quad \sum \sum x_{mn}y_{mn} = \infty.$$

Hence  $y \notin [\text{Ces}_p^q(\chi_f^2)]^\beta$ .

From (3.10) and (3.1), we are granted  $[\text{Ces}_p^q(\Lambda_f^2)]^\beta \not\subseteq \text{Ces}_p^q(\chi_f^2)$ . □




Go back

Full Screen

Close

Quit



**Proposition 3.9.** Let  $(\text{Ces}_p^q(\chi_f^2))^*$  denote the dual space of  $\text{Ces}_p^q(\chi_f^2)$ . Then we have  $(\text{Ces}_p^q(\chi_f^2))^* \text{Ces}_p^q(\Lambda_f^2)$ .

*Proof.* We recall that

$$x = \mathfrak{S}_{mn} = \begin{pmatrix} 0, & 0, & \dots & 0, & 0, & \dots & 0 \\ 0, & 0, & \dots & 0, & 0, & \dots & 0 \\ \vdots & & & & & & \\ 0, & 0, & \dots & \frac{1}{(m+n)!}, & 0, & \dots & 0 \\ 0, & 0, & \dots & 0, & 0, & \dots & 0 \end{pmatrix}$$

with  $\frac{1}{(m+n)!}$  in the  $(m, n)^{th}$  position and zero otherwise, with

$$\begin{aligned} x &= \mathfrak{S}_{mn} \left\{ \left( \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \left( \frac{1}{Q_{ij}} \sum_{m=1}^i \sum_{n=1}^j q_{mn} f \left( ((m+n)! |x_{mn}|^{\frac{1}{m+n}}) \right)^{p_{mn}} \right)^{\frac{1}{p_{mn}}} \right) \right\} \\ &= \begin{pmatrix} 0, & 0, & \dots & 0, & 0, & \dots & 0 \\ \vdots & & & & & & \\ 0, & 0, & \dots & \frac{(m+n)!}{(m+n)!}, & 0, & \dots & 0 \\ 0, & 0, & \dots & 0, & 0, & \dots & 0 \end{pmatrix} = \begin{pmatrix} 0, & 0, & \dots & 0, & 0, & \dots & 0 \\ \vdots & & & & & & \\ 0, & 0, & \dots & 1^{1/m+n}, & 0, & \dots & 0 \\ 0, & 0, & \dots & 0, & 0, & \dots & 0 \end{pmatrix} \end{aligned}$$



Go back

Full Screen

Close

Quit



which is a  $\text{Ces}_p^q(\chi_f^2)$  sequence. Hence  $\mathfrak{S}_{mn} \in \text{Ces}_p^q(\chi_f^2)$ . Let us take  $f(x) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} x_{mn} y_{mn}$  with  $x \in \text{Ces}_p^q(\chi_f^2)$  and  $f \in (\text{Ces}_p^q(\chi_f^2))^*$ . Take  $x = (x_{mn}) = \mathfrak{S}_{mn} \in \text{Ces}_p^q(\chi_f^2)$ . Then

$$|y_{mn}| \leq \|f\| d(\mathfrak{S}_{mn}, 0) < \infty \quad \text{for each } m, n.$$

Thus  $(y_{mn})$  is a bounded sequence and hence an Cesàro double analytic sequence of modulus. In other words  $y \in \text{Ces}_p^q(\Lambda_f^2)$ . Therefore  $(\text{Ces}_p^q(\chi_f^2))^* = \text{Ces}_p^q(\Lambda_f^2)$ .  $\square$

1. Apostol T., *Mathematical Analysis*. Addison-wesley, London, 1978.
2. Basarir M. and Solancan O., *On some double sequence spaces*. J. Indian Acad. Math. **21(2)** (1999), 193–200.
3. Bektas C. and Altin Y., *The sequence space  $\ell_M(p, q, s)$  on seminormed spaces*. Indian J. Pure Appl. Math. **34(4)** (2003), 529–534.
4. Bromwich T. J., *An introduction to the theory of infinite series*. Macmillan and Co.Ltd., New York, 1965.
5. Hardy G. H., *On the convergence of certain multiple series*, Proc. Camb. Phil. Soc. **19** (1917), 86–95.
6. Krasnoselskii M. A. and Rutickii Y. B., *Convex functions and Orlicz spaces*. Gorningen, Netherlands 1961.
7. Lindenstrauss J. and Tzafriri L., *On Orlicz sequence spaces*. Israel J. Math. **10** (1971), 379–390.
8. Maddox I.J., *Sequence spaces defined by a modulus*. Math. Proc. Cambridge Philos. Soc **100(1)** (1986), 161–166.
9. Moricz F., *Extentions of the spaces  $c$  and  $c_0$  from single to double sequences*. Acta. Math. Hung. **57(1–2)** (1991), 129–136.
10. Moricz F. and Rhoades B. E., *Almost convergence of double sequences and strong regularity of summability matrices*. Math. Proc. Camb. Phil. Soc. **104** (1988), 283–294.
11. Mursaleen M., Khan M. A. and Qamaruddin, *Difference sequence spaces defined by Orlicz functions*. Demonstratio Math. **XXXII** (1999), 145–150.
12. Nakano H., *Concave modulars*. J. Math. Soc. Japan **5** (1953), 29–49.
13. Orlicz W., *Über Räume  $(L^M)$* , Bull. Int. Acad. Polon. Sci. **A** (1936), 93–107.
14. Parashar S. D. and Choudhary B., *Sequence spaces defined by Orlicz functions*. Indian J. Pure Appl. Math. **25(4)** (1994), 419–428.




Go back

Full Screen

Close

Quit



15. Chandrasekhara Rao K. and Subramanian N., *The Orlicz space of entire sequences*. Int. J. Math. Math. Sci. **68** (2004), 3755–3764.
16. Ruckle W. H., *FK spaces in which the sequence of coordinate vectors is bounded*. Canad. J. Math. **25** (1973), 973–978.
17. Tripathy B. C., *On statistically convergent double sequences*. Tamkang J. Math. **34**(3), (2003), 231–237.
18. Tripathy B. C., Et M. and Altin Y., *Generalized difference sequence spaces defined by Orlicz function in a locally convex space*. J. Anal. Appl., **1**(3) (2003), 175–192.
19. Turkmenoglu A., *Matrix transformation between some classes of double sequences*. J. Inst. Math. Comp. Sci. Math. Ser. **12**(1) (1999), 23–31.
20. Kamthan P. K. and Gupta M., *Sequence spaces and series*. Lecture notes, Pure and Applied Mathematics 65. Marcel Dekker, Inc., New York, 1981.
21. Gökhan A. and Çolak R., *The double sequence spaces  $c_2^P(p)$  and  $c_2^{PB}(p)$* . Appl. Math. Comput. **157**(2) (2004), 491–501.
22. ———, *Double sequence spaces  $\ell_2^\infty$* . Appl. Math. Comput. **160**(1) (2005), 147–153.
23. Zeltser M., *Investigation of Double Sequence Spaces by Soft and Hard Analytical Methods*. Dissertationes Mathematicae Universitatis Tartuensis 25, Tartu University Press, Univ. of Tartu, Faculty of Mathematics and Computer Science, Tartu, 2001.
24. Mursaleen M. and Edely O. H. H., *Statistical convergence of double sequences*. J. Math. Anal. Appl. **288**(1) (2003), 223–231.
25. ———, *Almost convergence and a core theorem for double sequences*. J. Math. Anal. Appl. **293**(2) (2004), 532–540.
26. Mursaleen M., *Almost strongly regular matrices and a core theorem for double sequences*. J. Math. Anal. Appl. **293**(2) (2004), 523–531.
27. Altay B. and Başar F., *Some new spaces of double sequences*. J. Math. Anal. Appl. **309**(1) (2005), 70–90.
28. Başar F. and Sever Y., *The space  $\mathcal{L}_p$  of double sequences*. Math. J. Okayama Univ **51** (2009), 149–157.
29. Subramanian N. and Misra U. K., *The semi normed space defined by a double gai sequence of modulus function*. Fasciculi Math. **46** (2010).
30. Kizmaz H., *On certain sequence spaces*. Cand. Math. Bull. **24**(2) (1981), 169–176.
31. Kuttner B., *Note on strong summability*. J. London Math. Soc. **21** (1946), 118–122.
32. Maddox I. J., *On strong almost convergence*. Math. Proc. Cambridge Philos. Soc. **85**(2) (1979), 345–350.




Go back

Full Screen

Close

Quit



33. Cannon J., *On strong matrix summability with respect to a modulus and statistical convergence*. Canad. Math. Bull. **32**(2) (1989), 194–198.
34. Pringsheim A., *Zurtheorie der zweifach unendlichen zahlenfolgen*. Math. Ann. **53** (1900), 289–321.
35. Hamilton H. J., *Transformations of multiple sequences*. Duke Math. J. **2** (1936), 29–60.
36. ———, *A Generalization of multiple sequences transformation*. Duke Math. J. **4** (1938), 343–358.
37. ———, *Change of Dimension in sequence transformation*. Duke Math. J. **4** (1938), 341–342.
38. ———, *Preservation of partial Limits in Multiple sequence transformations*. Duke Math. J. **4** (1939), 293–297.
39. Robison G. M., *Divergent double sequences and series*. Amer. Math. Soc. Trans. **28** (1926), 50–73.
40. Silverman L. L., *On the definition of the sum of a divergent series*. Unpublished thesis, University of Missouri studies, Mathematics series.
41. Toeplitz O., *Über allgenmeine linear mittel bridungen*. Prace Matematyczno Fizyczne **22** (1911).
42. Başar F. and Atlay B., *On the space of sequences of  $p$ -bounded variation and related matrix mappings*. Ukrainian Math. J. **55**(1) (2003), 136–147.
43. Altay B. and Başar F., *The fine spectrum and the matrix domain of the difference operator  $\Delta$  on the sequence space  $\ell_p$ , ( $0 < p < 1$ )*. Commun. Math. Anal. **2**(2) (2007), 1–11.
44. Çolak R., Et M. and Malkowsky E., *Some Topics of Sequence Spaces, Lecture Notes in Mathematics*. Firat Univ. Elazig, **2004**, 1–63.

N. Subramanian, Department of Mathematics, SASTRA University, Thanjavur-613 401, India,  
*e-mail*: [nsmaths@yahoo.com](mailto:nsmaths@yahoo.com)



Go back

Full Screen

Close

Quit