



SUMS OF SEVENTH POWERS IN THE RING OF POLYNOMIALS OVER THE FINITE FIELD WITH FOUR ELEMENTS

MIREILLE CAR

ABSTRACT. We study representations of polynomials $P \in \mathbb{F}_4[T]$ as sums $P = X_1^7 + \dots + X_s^7$.

1. INTRODUCTION

Let F be a finite field of characteristic p with $q = p^m$ elements. Analogues of the Waring's problem for the polynomial ring $F[T]$ were investigated, ([19], [12], [16], [6], [17], [8], [5], [13], [14], [10], [9], [2], [3] [4]). Let $k > 1$ be an integer. Roughly speaking, Waring's problem over $F[T]$ consists in representing a polynomial $M \in F[T]$ as a sum

$$(1.1) \quad M = M_1^k + \dots + M_s^k$$

with $M_1, \dots, M_s \in F[T]$. Some obstructions to that may occur ([15]), and lead to consider Waring's problem over the subring $\mathcal{S}(F[T], k)$ formed by the polynomials of $F[T]$ which are sums of k -th powers. Some cancellations may occur in representations (1.1), so that it is possible to have a representation (1.1) with $\deg M$ small and $\deg(M_i^k)$ large. Without degree conditions in (1.1), the problem of representing M as sum (1.1) is close to the so called easy Waring's problem

Received March 6, 2012.

2010 *Mathematics Subject Classification*. Primary 11T55; Secondary 11P05.

Key words and phrases. Waring's problem; polynomials.



Go back

Full Screen

Close

Quit



for \mathbb{Z} . In order to have a problem close to the non-easy Waring's problem, the degree conditions

$$(1.2) \quad k \deg M_i < \deg M + k$$

are required. Representations (1.1) satisfying degree conditions (1.2) are called *strict representations*, see [6, Definition 1.8] in opposition to representations without degree conditions. For the strict Waring's problem, analogue of the classical Waring numbers $g_{\mathbb{N}}(k)$ and $G_{\mathbb{N}}(k)$ have been defined as follows. Let $g(p^m, k)$ denote the least integer s (if it exists) such that every polynomial $M \in \mathcal{S}(F[T], k)$ may be written as a sum (1.1) satisfying the degree conditions (1.2); otherwise we put $g(p^m, k) = \infty$.

Similarly, $G(p^m, k)$ denotes the least integer fulfilling the above condition for each polynomial $M \in \mathcal{S}(F[T], k)$ of sufficiently large degree. This notation is possible since these numbers depend only on p^m and k . The set $\mathcal{S}(F[T], k)$ and the parameters $G(p^m, k), g(p^m, k)$ are not sufficient to describe all possible cases, see [1, Proposition 4.4], so that in [2] and [3] we introduced new parameters defined as follows.

Let $\mathcal{S}^{\times}(F[T], k)$ denote the set of polynomials in $F[T]$ which are strict sums of k -th powers. Let $g^{\times}(p^m, k)$ denote the least integer s (if it exists) such that every polynomial $M \in \mathcal{S}^{\times}(F[T], k)$ may be written as a strict sum

$$M = M_1^k + \dots + M_s^k.$$

Similarly, $G^{\times}(p^m, k)$ denotes the least integer s fulfilling the same condition for each polynomial $M \in \mathcal{S}^{\times}(F[T], k)$ of sufficiently large degree. Gallardo's method for cubes ([8] and [5]) was generalized in [1] and [11] where bounds for $g(p^m, k)$ and $G(p^m, k)$ were established when p^m and k satisfy some conditions. A bound for $g(p^m, k)$ was established in [1] in the case when $F = \mathcal{S}(F, k)$ if one of the two following conditions is satisfied:

- i) $p > k$
- ii) $p^n > k = hp^{\nu} - 1$ for some integers $\nu > 0$ and $0 < h \leq p$.



Go back

Full Screen

Close

Quit



The smallest exponent k satisfying condition ii) is $k = 3$. It gave a matter for many articles, see [8], [5], [9], [10]. In the case of even characteristic, the second smallest exponent k satisfying condition ii) is $k = 7$. The case $k = 7, q = 2^m$ with $m > 3$ is covered by [1, Theorems 1.2 and 1.3] or by [11, Theorem 1.4]. For almost all $q = 2^m$, the upper bounds obtained in these articles for the numbers $G(2^m, 7)$ are comparable with the bound $G_{\mathbb{N}}(7) \leq 33$ known for the corresponding Waring's number for the integers ([18]). The case of the numbers $g(2^m, 7)$ is different. In the case when $m \notin \{1, 2, 3\}$ [1, Theorem 1.3] as well as [11, Theorem 1.4] gives $g(2^m, 7) \leq 239\ell(2^m, 7)$ when for the integers, it is known that $g_{\mathbb{N}}(7) = 143$ ([7]). In [4] we obtained better bounds for the numbers $g(2^m, 7)$ in the case when $m \notin \{1, 2, 3\}$, the method yielding also to better bounds for some numbers $G(2^m, 7)$. The aim of this paper is the study of one of the remaining cases, namely, the case $q = 4$. The case $q = 8$ will be the subject of a separate paper. When a finite field with 8 elements is not a 7-Waring field, every field with 4^h elements is a 7-Waring field, so that, from [15], $\mathcal{S}(\mathbb{F}_4[T], 7) = \mathbb{F}_4[T]$. We will see further that T is not a strict sum of seventh powers in the ring $\mathbb{F}_4[T]$, see Proposition 3.5 below, so that $\mathcal{S}(\mathbb{F}_4[T], 7) \neq \mathcal{S}^{\times}(\mathbb{F}_4[T], 7)$.

The main results proved in this work are summarized in the following theorems.

Theorem 1.1. *We have*

$$\mathcal{S}^{\times}(\mathbb{F}_4[T], 7) = \mathcal{A}_1 \cup \mathcal{A}_2 \cup \mathcal{A}_3 \cup \mathcal{A}_{\infty},$$

where

(i) \mathcal{A}_1 is the set of polynomials $A = \sum_{n=0}^7 a_n T^n \in \mathbb{F}_4[T]$ such that $a_1 = a_4, a_2 = a_5, a_3 = a_6$;



Go back

Full Screen

Close

Quit



(ii) \mathcal{A}_2 is the set of polynomials $A = \sum_{n=0}^{14} a_n T^n \in \mathbb{F}_4[T]$ with $7 < \deg A \leq 14$ such that

$$\begin{cases} a_1 + a_4 + a_{10} + a_{13} = 0, \\ a_2 + a_5 + a_8 + a_{11} = 0, \\ a_3 + a_6 + a_9 + a_{12} = 0; \end{cases}$$

(iii) \mathcal{A}_3 is the set of polynomials $A = \sum_{n=0}^{21} a_n T^n \in \mathbb{F}_4[T]$ with $14 < \deg A \leq 21$ such that

$$a_3 + a_6 + a_9 + a_{12} + a_{15} + a_{18} = 0;$$

(iv) $\mathcal{A}_\infty = \{A \in \mathbb{F}_4[T] \mid \deg A > 21\}$.

See Proposition 6.6 below.

Theorem 1.2. *Every polynomial $P \in \mathbb{F}_4[T]$ with degree ≥ 435 is a strict sum of 33 seventh powers, so that*

$$G(4, 7) = G^\times(4, 7) \leq 33,$$

and we have

$$g(4, 7) = \infty,$$

$$g^\times(4, 7) \leq 43.$$

This theorem is given by Corollaries 3.6, 6.4 and by Theorem 6.7

Proving that polynomials of small degree are sums or strict sums of seventh powers requires some results on the solvability of systems of algebraic equations over the finite field \mathbb{F}_4 . This is done in Section 2. A characterization of polynomials of degree ≤ 21 that are strict sums of seventh powers is given in Section 3. In Section 4, using the general descent process described in [1], we obtain a first upper bound for $G(4, 7)$. In Section 5 we describe other descent processes. They are



Go back

Full Screen

Close

Quit



used in Section 6 to get a better upper bound for $G(4, 7)$ as well as a bound for $g(4, 7)$. We denote by F the field \mathbb{F}_4 and by α a root of the equation $\alpha^2 = \alpha + 1$.

2. EQUATIONS

Proposition 2.1. *For every $(a, b) \in F^2$, the system*

$$(\mathcal{A}(a, b)) \quad \begin{cases} x_1 + x_2 = a, \\ u_1x_1 + u_2x_2 = b, \end{cases}$$

has solutions $(u_1, u_2, x_1, x_2) \in F^4$ satisfying the condition $x_1x_2u_1u_2 \neq 0$.

Proof. Suppose $a = b$. Choose $x_1 \in F - \{0, a\}$. Then, $(1, 1, x_1, a + x_1)$ is a solution of $(\mathcal{A}(a, b))$. Suppose $a \neq b$. There is $u_2 \in F - \mathbb{F}_2$ such that $au_2 + b \neq 0$. Then, $(1, u_2, \frac{au_2+b}{1+u_2}, a + \frac{au_2+b}{1+u_2})$ is a solution of $(\mathcal{A}(a, b))$. Moreover, since $a \neq b$, we have $\frac{au_2+b}{1+u_2} \neq a$, so that

$$u_2 \times \frac{au_2 + b}{1 + u_2} \times \left(a + \frac{au_2 + b}{1 + u_2} \right) \neq 0.$$

□

Proposition 2.2. *For $(a, b, c) \in F^3$, let $(\mathcal{B}_s(a, b, c))$ denote the system of equations*

$$\begin{cases} x_1 + \dots + x_s = a, \\ y_1 + \dots + y_s = b, \\ x_1y_1 + \dots + x_sy_s = c. \end{cases}$$

- (I) *For every $(a, b, c) \in F^\times \times F \times F$, the system $(\mathcal{B}_2(a, b, c))$ admits solutions $(x_1, x_2, y_1, y_2) \in F^4$ satisfying the condition $x_1x_2 \neq 0$.*
- (II) *For every $(a, b, c) \in F^3$, the system $(\mathcal{B}_3(a, b, c))$ admits solutions $(x_1, x_2, x_3, y_1, y_2, y_3) \in F^6$ satisfying the condition $x_1x_2x_3y_1y_2y_3 \neq 0$.*



Go back

Full Screen

Close

Quit



(III) For every $(a, b, c) \in F^\times \times F \times F$, the system $(\mathcal{B}_3(a, b, c))$ admits solutions $(x_1, x_2, x_3, y_1, y_2, y_3) \in F^6$ satisfying the conditions

$$\begin{cases} x_1 x_2 x_3 y_1 y_2 y_3 \neq 0, \\ x_1^2 y_1 \neq x_2^2 y_2. \end{cases}$$

Proof. (I) Suppose $a \neq 0$. Let $x_1 \in F - \{0, a\}$ and let $x_2 = a + x_1$. Then, $x_2 \neq 0$ and $x_2 \neq x_1$. The matrix

$$\begin{pmatrix} 1 & 1 \\ x_1 & x_2 \end{pmatrix}$$

is invertible. Thus, for each $(b, c) \in F^2$, there exists $(y_1, y_2) \in F^2$ such that

$$\begin{cases} y_1 + y_2 = b, \\ x_1 y_1 + x_2 y_2 = c. \end{cases}$$

(II) Let $\mathbf{E}(a, b, c)$ denote the set of $(x_1, x_2, x_3, y_1, y_2, y_3) \in F^6$ solutions of $(\mathcal{B}_3(a, b, c))$ satisfying $x_1 x_2 x_3 y_1 y_2 y_3 \neq 0$, and satisfying

$$\begin{cases} x_1 x_2 x_3 y_1 y_2 y_3 \neq 0, \\ x_1^2 y_1 \neq x_2^2 y_2, \end{cases}$$

respectively. For $(x_1, x_2, x_3, y_1, y_2, y_3) \in F^6$, the three following statements are equivalent:

- (i) $(x_1, x_2, x_3, y_1, y_2, y_3) \in \mathbf{E}(a, b, c)$,
- (ii) $(y_1, y_2, y_3, x_1, x_2, x_3) \in \mathbf{E}(b, a, c)$,
- (iii) $(x_1 y_1, x_2 y_2, x_3 y_3, (y_1)^2, (y_2)^2, (y_3)^2) \in \mathbf{E}(c, b^2, a)$. Thus, it suffices to deal with the cases $(a, b, c) = (0, 0, 0)$, $(a, b, c) = (a, 0, 0)$ with $a \neq 0$, $(a, b, c) = (a, b, 0)$ with $ab \neq 0$, and (a, b, c) with $abc \neq 0$. Firstly, we observe that if $x \in F - \mathbb{F}_2$, then $(1, x, x+1, 1, x, x+1) \in \mathbf{E}(0, 0, 0)$. Now, we consider the systems with $a \neq 0$. Up to the automorphism $x \mapsto ax$, and the \mathbb{F}_2 -automorphism $\alpha \mapsto \alpha + 1$, it suffices to consider the cases $(a, b, c) = (1, 0, 0)$,



Go back

Full Screen

Close

Quit



$$\begin{aligned}
 (a, b, c) &= (1, 1, 0), \quad (a, b, c) = (1, 1, 1), \quad (a, b, c) = (1, 1, \alpha). \text{ Observe that} \\
 (1, 1, 1, 1, \alpha, \alpha + 1) &\in \mathbf{E}(1, 0, 0), \\
 (\alpha + 1, \alpha + 1, 1, 1, \alpha, \alpha) &\in \mathbf{E}(1, 1, 0), \\
 (1, \alpha, \alpha, 1, 1, 1) &\in \mathbf{E}(1, 1, 1), \\
 (1, \alpha + 1, \alpha + 1, \alpha + 1, \alpha + 1, 1) &\in \mathbf{E}(1, 1, \alpha).
 \end{aligned}$$

□

Proposition 2.3. For every $(a, b, c) \in F^3$, the system

$$(\mathcal{C}(a, b, c)) \quad \begin{cases} x_1 + x_2 = a, \\ y_1 + y_2 = b, \\ x_1 y_1 z_1^2 + x_1^2 y_1^2 + x_2 y_2 z_2^2 + x_2^2 y_2^2 = c \end{cases}$$

admits solutions $(x_1, x_2, y_1, y_2, z_1, z_2) \in F^6$ satisfying the condition $x_1 x_2 y_1 y_2 \neq 0$.

Proof. Let $x_1 \in F$ be such that $x_1 \neq 0, a$, let $y_1 \in F$ be such that $y_1 \neq 0, b$ and let $z_1 \in F$. Let $x_2 = a - x_1$ and $y_2 = b - y_1$. Then, $x_1 x_2 y_1 y_2 \neq 0$. Let $z_2 \in F$ be defined by the relation $x_2^2 y_2^2 z_2 = c^2 + x_1^2 y_1^2 z_1 + x_1 y_1 + x_2 y_2$. Then, $(x_1, x_2, y_1, y_2, z_1, z_2)$ is a solution of $(\mathcal{C}(a, b, c))$. □

Lemma 2.4. For every $(a, b, c) \in F^\times \times F \times F$, the system of equations

$$(\mathcal{S}_1(a, b, c)) \quad \begin{cases} z_1 + \alpha z_2 + (\alpha + 1) z_3 = b, \\ \alpha z_1 + z_1^2 + (\alpha + 1) \alpha z_2 + z_2^2 + \alpha \alpha z_3 + z_3^2 = c, \end{cases}$$

admits solutions $(z_1, z_2, z_3) \in F^3$.

Proof. Let $\nu = \nu(a, b, c)$ denote the number of $(z_1, z_2, z_3) \in F^3$ solutions of $(\mathcal{S}_1(a, b, c))$. For $t \in F$ let

$$\Psi(t) = (-1)^{\text{tr}(t)}$$




Go back

Full Screen

Close

Quit



where $\text{tr}: F \rightarrow \mathbb{F}_2$ is the absolute trace map. Then Ψ is a non-trivial character, so that by orthogonality,

$$\begin{aligned} \nu &= \sum_{(z_1, z_2, z_3) \in F^3} \frac{1}{4} \sum_{t \in F} \Psi(t(b + z_1 + \alpha z_2 + (\alpha + 1)z_3)) \\ &\times \frac{1}{4} \sum_{u \in F} \Psi(u(c + az_1 + z_1^2 + (\alpha + 1)az_2 + z_2^2 + \alpha az_3 + z_3^2)). \end{aligned}$$

Thus,

$$\begin{aligned} 16\nu &= \sum_{(t, u) \in F^2} \Psi(bt + cu) \left(\sum_{z \in F} \Psi((t + au)z + uz^2) \right) \\ &\times \left(\sum_{z \in F} \Psi((\alpha t + a(\alpha + 1)u)z + uz^2) \right) \\ &\times \left(\sum_{z \in F} \Psi(((\alpha + 1)t + \alpha au)z + uz^2) \right). \end{aligned}$$

From [2, Proposition 2.3], for $(v, w) \in F^2$, we have

$$\sum_{z \in F} \Psi(vz + wz^2) = \begin{cases} 4 & \text{if } w = v^2, \\ 0 & \text{if } w \neq v^2. \end{cases}$$

Therefore,

$$\nu = 4 \sum_{(t, u) \in \mathbf{E}} \Psi(bt + cu),$$



Go back

Full Screen

Close

Quit



where \mathbf{E} is the subset of F^2 formed by the pairs (t, u) satisfying the three conditions

$$\begin{cases} u = (t + au)^2, \\ u = (\alpha t + a(\alpha + 1)u)^2, \\ u = ((\alpha + 1)t + \alpha au)^2. \end{cases}$$

Obviously, $(0, 0) \in \mathbf{E}$. Conversely, let $(t, u) \in \mathbf{E}$. Then the first and second conditions give that $t + au = \alpha t + a(\alpha + 1)u$ while the first and last conditions give that $t + au = (\alpha + 1)t + \alpha au$, so that $(\alpha + 1)au = t = \alpha au$ with $a \neq 0$. Thus, $t = u = 0$, so that $\mathbf{E} = \{(0, 0)\}$ and $\nu = 4$. \square

Proposition 2.5. Let $\mathbf{b} = (a, b, c, d) \in F^4$. Then the system of equations

$$(\mathcal{D}(\mathbf{b})) \quad \begin{cases} \sum_{i=1}^3 y_i = a, \\ \sum_{i=1}^3 u_i y_i = b, \\ \sum_{i=1}^3 u_i^2 z_i^2 y_i = c, \\ \sum_{i=1}^3 (u_i^2 z_i + u_i z_i^2) y_i = d \end{cases}$$

admits solutions $(u_1, u_2, u_3, y_1, y_2, y_3, z_1, z_2, z_3) \in F^9$ such that $u_1 u_2 u_3 y_1 y_2 y_3 \neq 0$.

Proof. (I) Suppose that there exists $(y_1, y_2, y_3, u) \in F^4$ satisfying the conditions:

$$\begin{cases} y_1 + y_2 + y_3 = a, \\ y_1 + y_2 + uy_3 = b, \\ y_1 y_2 y_3 u \neq 0, \\ y_1 \neq y_2, \end{cases}$$




Go back

Full Screen

Close

Quit



and denote (H) this hypothesis. Then the matrix

$$\begin{pmatrix} y_1 & y_2 \\ y_1^2 & y_2^2 \end{pmatrix}$$

is invertible. Let $z_3 \in F$. There is $(z_1, z_2) \in F^2$ such that

$$\begin{aligned} y_1 z_1 + y_2 z_2 &= c + d + (u^2 z_3 + u^2 z_3^2 + u z_3^2) y_3, \\ y_1^2 z_1 + y_2^2 z_2 &= c^2 + u z_3 y_3^2. \end{aligned}$$

Then, we have

$$\begin{aligned} z_1^2 y_1 + z_2^2 y_2 + u^2 z_3^2 y_3 &= c, \\ (z_1 + z_1^2) y_1 + (z_2 + z_2^2) y_2 + (u^2 z_3 + u z_3^2) y_3 &= d, \end{aligned}$$

so that $(1, 1, u, y_1, y_2, y_3, z_1, z_2, z_3)$ is a solution of $(\mathcal{D}(\mathbf{b}))$ such that $u y_1 y_2 y_3 \neq 0$.

(II) We prove that if one of the three following conditions:

- (i) $a = b$,
- (i) $a \notin \{0, b, (\alpha + 1)b\}$,
- (iii) $a = (\alpha + 1)b \neq 0$, (so that $a \neq b$),

is satisfied, then hypothesis (H) is satisfied, so that the conclusion of the proposition holds.



Go back

Full Screen

Close

Quit



(i) Suppose $a = b$. If $a = 0$, then $(1, \alpha, \alpha + 1)$ is a solution of (e_1) . If $a \neq 0$, then (a, y, y) with $y \notin \{0, a\}$ is a solution of (e_1) . Thus, in the two cases, (e_1) admits solutions $(y_1, y_2, y_3) \in F^3$ such that $y_1 y_2 y_3 \neq 0$ and $y_1 \neq y_2$. Hypothesis (H) is satisfied with $u = 1$.

(ii) Suppose $a \notin \{0, b, (\alpha + 1)b\}$. Then $a + \alpha(a + b) \neq 0$. Let $u = \alpha$, $y_3 = \alpha(a + b)$. Choose $y_1 \in F - \{0, a + \alpha(a + b)\}$ and $y_2 = y_1 + a + \alpha(a + b)$. Then, $y_1 \neq y_2$ and $y_1 y_2 y_3 \neq 0$, so that (H) is satisfied.

(iii) Suppose $a = (\alpha + 1)b \neq 0$. Let $u = (\alpha + 1)$, $y_3 = b$. Choose $y_1 \in F - \{0, \alpha b\}$ and $y_2 = y_1 + \alpha b$. Then, $y_1 \neq y_2$, $y_1 y_2 y_3 \neq 0$ and $y_1 + y_2 + y_3 = (\alpha + 1)b = a$, $y_1 + y_2 + u y_3 = \alpha b + (\alpha + 1)b = b$, so that (H) is satisfied.

(III) We examine the remaining case, that is the case $a = 0$, $b \neq 0$. Lemma 2.4 gives the existence of $(z_1, z_2, z_3) \in F^3$, a solution of $(\mathcal{S}_1(b, c^2/b, d/b))$ such that

$$\begin{aligned} b^2 z_1^2 + (\alpha + 1)b^2 z_2^2 + \alpha b^2 z_3^2 &= c, \\ b^2 z_1 + b z_1^2 + (\alpha + 1)b^2 z_2 + b z_2^2 + \alpha b^2 z_3 + b z_3^2 &= d. \end{aligned}$$

Let

$$u_1 = b, u_2 = (\alpha + 1)b, u_3 = \alpha b, y_1 = 1, y_2 = \alpha, y_3 = \alpha + 1.$$

Then, $(u_1, u_2, u_3, y_1, y_2, y_3, z_1, z_2, z_3)$ is a solution of $(\mathcal{D}(\mathbf{b}))$ such that $u_1 u_2 u_3 y_1 y_2 y_3 \neq 0$. \square

Lemma 2.6. Let $(a, b) \in F^2$. Then the system of equations

$$(\mathcal{S}_2(a, b)) \quad \begin{cases} u_1 + u_2 + u_3 = a, \\ x_1 + x_2 + x_3 = b, \end{cases}$$

admits solutions $(u_1, u_2, u_3, x_1, x_2, x_3) \in F^6$ satisfying the conditions

$$(2.1) \quad u_1 u_2 u_3 \neq 0,$$




Go back

Full Screen

Close

Quit



$$(2.2) \quad \det \begin{pmatrix} 1 & 1 & 1 \\ u_1 & u_2 & u_3 \\ u_1x_1^2 & u_2x_2^2 & u_3x_3^2 \end{pmatrix} \neq 0.$$

Proof. If $(u_1, u_2, u_3, x_1, x_2, x_3) \in F^6$ is a solution of $(\mathcal{S}_2(0, 1))$ satisfying conditions (2.1) and (2.2), then for $b \in F, b \neq 0$, $(u_1, u_2, u_3, bx_1, bx_2, bx_3)$ is a solution of $(\mathcal{S}_2(0, b))$ satisfying conditions (2.1) and (2.2). If $(u_1, u_2, u_3, x_1, x_2, x_3) \in F^6$ is a solution of $(\mathcal{S}_2(1, 0))$ satisfying conditions (2.1) and (2.2), then for $a \in F, a \neq 0$, $(au_1, au_2, au_3, x_1, x_2, x_3)$ is a solution of $(\mathcal{S}_2(a, 0))$ satisfying conditions (2.1) and (2.2). If $(u_1, u_2, u_3, x_1, x_2, x_3) \in F^6$ is solution of $(\mathcal{S}_2(1, 1))$ satisfying conditions (2.1) and (2.2), then for $a, b \in F, ab \neq 0$, $(au_1, au_2, au_3, bx_1, bx_2, bx_3)$ is a solution of $(\mathcal{S}_2(a, b))$ satisfying conditions (2.1) and (2.2). It is sufficient to examine the cases $(a, b) = (0, 0), (a, b) = (0, 1), (a, b) = (1, 0), (a, b) = (1, 1)$. Observe that

$(1, \alpha, \alpha^2, \alpha, 1, \alpha^2)$ is a solution of $(\mathcal{S}_2(0, 0))$ satisfying conditions (2.1) and (2.2);

$(1, \alpha, \alpha^2, 0, 0, 1)$ is a solution of $(\mathcal{S}_2(0, 1))$ satisfying conditions (2.1) and (2.2);

$(1, \alpha, \alpha, 1, \alpha, \alpha^2)$ is a solution of $(\mathcal{S}_2(1, 0))$ satisfying conditions (2.1) and (2.2);

$(1, \alpha, \alpha, 0, 0, 1)$ is a solution of $(\mathcal{S}_2(1, 1))$ satisfying conditions (2.1) and (2.2). □

Lemma 2.7. Let $(u_1, u_2, u_3, x_1, x_2, x_3) \in F^6$ be such that

$$(2.1) \quad u_1u_2u_3 \neq 0,$$

$$(2.2) \quad \det \begin{pmatrix} 1 & 1 & 1 \\ u_1 & u_2 & u_3 \\ u_1x_1^2 & u_2x_2^2 & u_3x_3^2 \end{pmatrix} \neq 0.$$



Go back

Full Screen

Close

Quit



Then, for every $(c, d) \in F^2$, there exists $(y_1, y_2, y_3) \in F^3$ such that

$$(\mathcal{S}_3(c, d)) \quad \begin{cases} y_1 + y_2 + y_3 = c, \\ u_1^2 y_1^2 + u_1 x_1^2 y_1 + \dots + u_3^2 y_3^2 + u_3 x_3^2 y_3 = d. \end{cases}$$

Proof. Let N denote the number of $(y_1, y_2, y_3) \in F^3$ solutions of $(\mathcal{S}_3(c, d))$. With the notations used in the proof of Lemma 2.4, we have

$$\begin{aligned} N &= \sum_{(y_1, y_2, y_3) \in F^3} \frac{1}{4} \sum_{t \in F} \Psi(t(c + y_1 + y_2 + y_3)) \\ &\quad \times \frac{1}{4} \sum_{u \in F} \Psi(u(d + u_1^2 y_1^2 + u_2^2 y_2^2 + u_3^2 y_3^2)). \end{aligned}$$

Thus,

$$16N = \sum_{(t, u) \in F^2} \Psi(ct + du) \prod_{i=1}^3 \Theta_i(t, u),$$

where

$$\Theta_i(t, u) = \sum_{y \in F} \Psi(ty + u(u_i^2 y^2 + u_i x_i^2 y)).$$

From [2, Proposition 2.3], $\Theta_i(t, u) \in \{0, 4\}$ and $\Theta_i(t, u) = 4$ if and only if $uu_i^2 = (t + uu_i x_i^2)^2$. Thus,

$$N = 4 \sum_{(t, u) \in E} \Psi(ct + du),$$



Go back

Full Screen

Close

Quit



where E is the set of pairs $(t, u) \in F^2$ such that

$$\begin{cases} t + uu_1x_1^2 = u^2u_1, \\ t + uu_2x_2^2 = u^2u_2, \\ t + uu_3x_3^2 = u^2u_3. \end{cases}$$

Observe that $(0, 0) \in E$. Moreover, if $(t, 0) \in E$, then $t = 0$. Suppose that $(t, u) \in E$ with $u \neq 0$. Then,

$$t = u(u_1x_1^2 + uu_1) = u(u_2x_2^2 + uu_2) = u(u_3x_3^2 + uu_3),$$

so that

$$\begin{cases} u_1x_1^2 + uu_1 = u_2x_2^2 + uu_2, \\ u_1x_1^2 + uu_1 = u_3x_3^2 + uu_3. \end{cases}$$

Thus,

$$\begin{cases} u_1x_1^2 + u_2x_2^2 = u(u_1 + u_2), \\ u_1x_1^2 + u_3x_3^2 = u(u_1 + u_3), \end{cases}$$

so that

$$(u_1x_1^2 + u_2x_2^2)(u_1 + u_3) + (u_1x_1^2 + u_3x_3^2)(u_1 + u_2),$$

in contradiction with condition (2.2). □




Go back

Full Screen

Close

Quit



Proposition 2.8. Let $\mathbf{b} = (b_1, b_2, \dots, b_7) \in F^7$. Then the system of equations

$$(\mathcal{E}(\mathbf{b})) \quad \begin{cases} \sum_{i=1}^3 u_i = b_1, \\ \sum_{i=1}^3 x_i = b_2, \\ \sum_{i=1}^3 (y_i + u_i^2 x_i^2) = b_3, \\ \sum_{i=1}^3 (z_i + u_i x_i^3) = b_4, \\ \sum_{i=1}^3 (u_i^2 y_i^2 + u_i x_i^2 y_i) = b_5, \\ \sum_{i=1}^3 (u_i x_i^2 z_i + u_i x_i y_i^2 + u_i^2 x_i^2) = b_6, \\ \sum_{i=1}^3 (u_i^2 z_i^2 + u_i y_i^3 + u_i^2 x_i y_i + u_i x_i^3) = b_7 \end{cases}$$

admits solutions $(u_1, u_2, u_3, x_1, x_2, x_3, y_1, y_2, y_3, z_1, z_2, z_3) \in F^{12}$ with $u_1 u_2 u_3 \neq 0$.



Go back

Full Screen

Close

Quit



Proof. Lemma 2.6 gives the existence of $(u_1, u_2, u_3, x_1, x_2, x_3) \in F^6$, a solution of $\mathcal{S}_2(b_1, b_2)$ satisfying (2.1) and (2.2). Lemma 2.7 gives the existence of $(y_1, y_2, y_3) \in F^3$, a solution of $\mathcal{S}_3(b_3 + \sum_{i=1}^3 u_i^2 x_i^2, b_5)$. Condition (2.2) insures the existence of $(z_1, z_2, z_3) \in F^3$ such that

$$\left\{ \begin{array}{l} z_1 + z_2 + z_3 = b_4 + \sum_{i=1}^3 u_i x_i^3, \\ u_1 z_1 + u_2 z_2 + u_3 z_3 = b_7^2 + \sum_{i=1}^3 (u_i^2 x_i^3 + u_i^2 y_i^3 + u_i x_i^2 y_i^2), \\ u_1 x_1^2 z_1 + u_2 x_2^2 z_2 + u_3 x_3^2 z_3 = b_6 + \sum_{i=1}^3 (u_i x_i y_i^2 + u_i^2 x_i^2). \end{array} \right.$$

□

Lemma 2.9. *Let $(a, b, c) \in F^\times \times F^2$ be such that $ab + c \neq 0$. Then the system*

$$(\mathcal{S}_4(a, b, c)) \quad \left\{ \begin{array}{l} u + v = a, \\ x + y = b, \\ ux + vy = c \end{array} \right.$$

admits a solution $(u, x, v, y) \in F^4$ such that $uv \neq 0$ and $u^2x + v^2y \neq 0$.

Proof. Let $u \in F - \{0, a\}$ and $v = u + a$. Then $uv(u + v) \neq 0$, so that with $x = (bu + c + ab)/a$ and $y = (bu + c)/a$, (u, x, v, y) is a solution of $(\mathcal{S}_4(a, b, c))$. Suppose that $u^2x + v^2y = 0$. Then, $u^2b + uab + ac = 0$. If $b = 0$, then $c = 0$, in contradiction with $ab + c \neq 0$. Thus $b \neq 0$, so that $u^2 + au + \frac{ac}{b} = 0$ and $\frac{c}{ab} \in \{0, 1\}$. Thus, $c = 0$. We have $u^2x + v^2y = 0$ and $ux + vy = 0$. Since $b \neq 0$, we have $(x, y) \neq (0, 0)$. If $x = 0$, then $vy = 0$, so that $y = 0$, a contradiction. Similarly,



Go back

Full Screen

Close

Quit



$y = 0$ is impossible. Thus, $xy \neq 0$. Therefore $u = u^2x/ux = v^2y/vy = v$ in contradiction with $u + v \neq 0$. Hence, (u, x, v, y) is a solution of $(\mathcal{S}_4(a, b, c))$ such that $uv \neq 0$ and $u^2x + v^2y \neq 0$. \square

Proposition 2.10. Let $\mathbf{b} = (b_1, b_2, \dots, b_8) \in F^8$. Then the system of equations

$$(\mathcal{F}(\mathbf{b})) \quad \begin{cases} \sum_{i=1}^3 v_i = b_1, \\ \sum_{i=1}^3 u_i = b_2, \\ \sum_{i=1}^3 (x_i + v_i^2 u_i^2) = b_3, \\ \sum_{i=1}^3 (y_i + v_i u_i^3) = b_4, \\ \sum_{i=1}^3 (v_i^2 x_i^2 + v_i u_i^2 x_i + u_i + z_i) = b_5, \\ \sum_{i=1}^3 (v_i u_i^2 y_i + v_i u_i x_i^2 + v_i^2 u_i^2) = b_6, \\ \sum_{i=1}^3 (v_i u_i^3 + v_i x_i^3 + v_i^2 y_i^2 + v_i u_i^2 z_i + v_i^2 u_i x_i) = b_7, \\ \sum_{i=1}^3 (v_i^2 u_i y_i + v_i u_i y_i^2 + v_i x_i^2 y_i) = b_8 \end{cases}$$

admits solutions

$$(v_1, v_2, v_3, u_1, u_2, u_3, x_1, x_2, x_3, y_1, y_2, y_3, z_1, z_2, z_3) \in F^{15}$$

satisfying the condition $v_1 v_2 v_3 \neq 0$.



Go back

Full Screen

Close

Quit



Proof. Let $v_1 = 1$, $v_2 \in F - \{0, 1, b_1 + 1\}$ and let v_3 be defined by

$$v_1 + v_2 + v_3 = b_1.$$

Then we have

$$v_1 v_2 v_3 \neq 0, \quad v_1 \neq v_2.$$

Let $u_1 \in F - \{0, (v_1 v_2)^2\}$, $u_3 = 0$ and let u_2 be defined by

$$u_1 + u_2 + u_3 = b_2.$$

Then we have

$$(\dagger) \quad v_1 u_1^2 \neq v_3 u_3^2.$$

(I) Suppose that $v_1 u_1 + v_2 u_2 = 0$. Let $y_1 \in F$, $y_2 \in F - \{u_2 y_1 / u_1\}$ and let y_3 be defined by

$$y_1 + y_2 + y_3 = b_4 + \sum_{i=1}^3 v_i u_i^3.$$

Then we have

$$v_1 v_2 (u_1 y_2 + u_2 y_1) \neq 0,$$

so that

$$\det \begin{pmatrix} 1 & 1 & 1 \\ v_1 u_1 & v_2 u_2 & v_3 u_3 \\ v_1 y_1 & v_2 y_2 & v_3 y_3 \end{pmatrix} \neq 0.$$



Go back

Full Screen

Close

Quit



(II) Suppose that $v_1u_1 + v_2u_2 \neq 0$. Let $y_1 \in F$. Since $u_1 \neq (v_1v_2)^2$, we have $1 + v_1v_2u_1 \neq 0$. Let $y_2 \in F$ be such that

$$(1 + v_1v_2u_1)y_2 + (1 + v_1v_2u_2)y_1 \neq v_3(v_1u_1 + v_2u_2)(b_4 + \sum_{i=1}^3 v_iu_i^3),$$

and let y_3 be defined by

$$y_1 + y_2 + y_3 = b_4 + \sum_{i=1}^3 v_iu_i^3.$$

Then we have

$$v_1v_2(u_1y_2 + u_2y_1) + v_3y_3(v_1u_1 + v_2u_2) \neq 0,$$

so that

$$\det \begin{pmatrix} 1 & 1 & 1 \\ v_1u_1 & v_2u_2 & v_3u_3 \\ v_1y_1 & v_2y_2 & v_3y_3 \end{pmatrix} \neq 0.$$

In both cases we get the existence of $(y_1, y_2, y_3) \in F^3$ satisfying

$$\det \begin{pmatrix} 1 & 1 & 1 \\ v_1^2u_1^2 & v_2^2u_2^2 & v_3^2u_3^2 \\ v_1^2y_1^2 & v_2^2y_2^2 & v_3^2y_3^2 \end{pmatrix} \neq 0$$



Go back

Full Screen

Close

Quit



from which we deduce the existence of $(x_1, x_2, x_3) \in F^3$ such that

$$\begin{aligned}x_1 + x_2 + x_3 &= b_3 + \sum_{i=1}^3 v_i^2 u_i^2, \\v_1^2 u_1^2 x_1 + v_2^2 u_2^2 x_2 + v_3^2 u_3^2 x_3 &= (b_1 + b_2 + b_6)^2 \sum_{i=1}^3 v_i u_i^2 y_i, \\v_1^2 y_1^2 x_1 + v_2^2 y_2^2 x_2 + v_3^2 y_3^2 x_3 &= b_8^2 \sum_{i=1}^3 (v_i u_i^2 y_i^2 + v_i^2 u_i^2 y_i).\end{aligned}$$

From (†),

$$\det \begin{pmatrix} 1 & 1 \\ v_1 u_1^2 & v_2 u_2^2 \end{pmatrix} \neq 0.$$

Then there exists $(z_1, z_3) \in F^2$ such that

$$\begin{aligned}z_1 + z_3 &= b_2 + b_5 + \sum_{i=1}^3 (v_i^2 x_i^2 + v_i u_i^2 x_i), \\v_1 u_1^2 z_1 + v_3 u_3^2 z_3 &= b_7 + \sum_{i=1}^3 (v_i u_i^3 + v_i x_i^3 + v_i^2 y_i^2 + v_i^2 u_i x_i).\end{aligned}$$

Let $z_2 = 0$. Then, $(v_1, v_2, v_3, u_1, u_2, u_3, x_1, x_2, x_3, y_1, y_2, y_3, z_1, z_2, z_3)$ is a solution of $(\mathcal{F}(\mathbf{b}))$ satisfying $v_1 v_2 v_3 \neq 0$. □



Go back

Full Screen

Close

Quit

Proposition 2.11. Let $\mathbf{b} = (b_1, b_2, \dots, b_9) \in F^9$. Then the system of equations

$$(\mathcal{G}(\mathbf{b})) \quad \left\{ \begin{array}{l} \sum_{i=1}^8 u_i = b_1, \\ \sum_{i=1}^8 x_i = b_2, \\ \sum_{i=1}^8 (y_i + u_i^2 x_i^2) = b_3, \\ \sum_{i=1}^8 (z_i + u_i x_i^3) = b_4, \\ \sum_{i=1}^8 (u_i^2 y_i^2 + u_i x_i^2 y_i) = b_5, \\ \sum_{i=1}^8 (u_i x_i^2 z_i + u_i x_i y_i^2 + u_i^2 x_i^2) = b_6, \\ \sum_{i=1}^8 (u_i^2 z_i^2 + u_i y_i^3 + u_i^2 x_i y_i + u_i x_i^3) = b_7, \\ \sum_{i=1}^8 (u_i^2 y_i z_i + u_i x_i^2 z_i + u_i y_i z_i^2) = b_8, \\ \sum_{i=1}^8 (u_i^2 x_i z_i + u_i x_i z_i^2 + u_i y_i^2 z_i) = b_9 \end{array} \right.$$

admits solutions $(u_1, \dots, u_8, x_1, \dots, x_8, y_1, \dots, y_8, z_1, \dots, z_8) \in F^{32}$ such that $u_1 \dots u_8 \neq 0$.



Go back

Full Screen

Close

Quit



Proof. Proposition 2.1 insures the existence of a solution (x_1, x_2, u_1, u_2) of $(\mathcal{A}(b_2, b_6^2))$ such that $u_1 u_2 x_1 x_2 \neq 0$. Thus, we have

$$\begin{aligned}x_1 + x_2 &= b_2, \\u_1 x_1^2 z_1 + u_1 x_1 y_1^2 + u_1^2 x_1^2 + u_2 x_2^2 z_2 + u_2 x_2 y_2^2 + u_2^2 x_2^2 &= b_6.\end{aligned}$$

Let $y_1 = y_2 = z_1 = z_2 = 0$. Proposition 2.5 insures the existence of a solution $(u_3, u_4, u_5, y_3, y_4, y_5, z_3, z_4, z_5)$ of $(\mathcal{D}((b_3 + b_6, b_5^2, b_9^2, b_8)))$ such that $u_3 u_4 u_5 y_3 y_4 y_5 \neq 0$. Let $x_3 = x_4 = x_5 = 0$. Then, we have

$$\left\{ \begin{array}{l} \sum_{i=1}^5 x_i = b_2, \\ \sum_{i=1}^5 (y_i + u_i^2 x_i^2) = b_3, \\ \sum_{i=1}^5 (u_i^2 y_i^2 + u_i x_i^2 y_i) = b_5, \\ \sum_{i=1}^5 (u_i x_i^2 z_1 + u_i x_i y_i^2 + u_i^2 x_i^2) = b_6, \\ \sum_{i=1}^5 (u_i^2 y_i z_i + u_i x_i^2 z_i + u_i y_i z_i^2) = b_8, \\ \sum_{i=1}^5 (u_i^2 x_i z_i + u_i x_i z_i^2 + u_i y_i^2 z_i) = b_9. \end{array} \right.$$



Go back

Full Screen

Close

Quit



Let

$$\beta_1 = b_1 + \sum_{i=1}^5 u_i,$$

$$\beta_4 = b_4 + \sum_{i=1}^5 (z_i + u_i x_i^3),$$

$$\beta_7 = b_7 + \sum_{i=1}^5 (u_i^2 z_i^2 + u_i y_i^3 + u_i^2 x_i y_i + u_i x_i^3).$$

From Proposition 2.2, $(\mathcal{B}_3(\beta_1, \beta_4, \beta_7^2))$ admits a solution $(u_6, u_7, u_8, z_6, z_7, z_8) \in F^6$ such that $u_6 u_7 u_8 z_6 z_7 z_8 \neq 0$. Let $x_6 = x_7 = x_8 = y_6 = y_7 = y_8 = 0$. Then, $(u_1, \dots, u_8, x_1, \dots, x_8, y_1, \dots, y_8, z_1, \dots, z_8)$ is a solution of $(\mathcal{G}(\mathbf{b}))$ such that $u_1 \dots u_8 \neq 0$. \square

3. STRICT SUMS OF DEGREE LESS THAN 21 IN $F[T]$

The aim of this section is the proof of the three following theorems.

Theorem 3.1. *Let $A \in F[T]$ with degree ≤ 7 , say*

$$A = \sum_{i=0}^7 a_i T^i.$$

Then, A is a strict sum of seventh powers if and only if its coefficients a_i satisfy the conditions

$$(3.1) \quad \begin{cases} a_1 = a_4, \\ a_2 = a_5, \\ a_3 = a_6. \end{cases}$$



Go back

Full Screen

Close

Quit



Moreover, if A is a strict sum of seventh powers, then A is a strict sum of 5 seventh powers.

Theorem 3.2. Let $A \in F[T]$ with degree ≤ 14 , say

$$A = \sum_{i=0}^{14} a_i T^i.$$

Then, A is a strict sum of seventh powers if and only if its coefficients a_i satisfy the conditions

$$(3.2) \quad \begin{cases} a_1 + a_4 + a_{10} + a_{13} = 0, \\ a_2 + a_5 + a_8 + a_{11} = 0, \\ a_3 + a_6 + a_9 + a_{12} = 0. \end{cases}$$

Moreover, if A is a strict sum of seventh powers, then A is a sum of 11 seventh powers.

Theorem 3.3. Let $A \in F[T]$ be such that $15 \leq \deg A \leq 21$, say

$$A = \sum_{i=0}^{21} a_i T^i.$$

Then, A is a strict sum of seventh powers if and only if its coefficients a_1, \dots, a_{21} satisfy the condition

$$(3.3) \quad a_3 + a_6 + a_9 + a_{12} + a_{15} + a_{18} = 0.$$

Moreover, if A satisfies condition (3.3), then A is a strict sum of 19 seventh powers.

Theorem 3.1 is a consequence of the two following propositions.



Go back

Full Screen

Close

Quit



Proposition 3.4. For $(a, b, c) \in F^3$,

$$(3.4) \quad \begin{aligned} cT^7 + (aT^2 + bT + c)(T^4 + T) + a \\ = ((a + b + c)(T + 1))^7 + ((\alpha^2 a + \alpha b + c)(T + \alpha))^7 \\ + ((\alpha a + \alpha^2 b + c)(T + \alpha^2))^7. \end{aligned}$$

Proof. A verification. □

Proposition 3.5.

- (i) Let $A \in F[T]$ be such that $\deg A \leq 6$. If A is a strict sum of seventh powers, then its coefficients satisfy (3.1).
- (ii) Let

$$A = \sum_{i=0}^7 a_i T^i$$

in the polynomial ring $F[T]$ be such that conditions (3.1) are satisfied. Then, A is a strict sum of 5 seventh powers.

Proof. Let $A = a_0 + a_1 T + \dots + a_6 T^6 \in F[T]$. Suppose that A is a strict sum of s seventh powers. Then,

$$A = \sum_{i=1}^s (x_i T + y_i)^7$$

with $x_i, y_i \in F$ for $i = 1, \dots, s$. Thus,

$$a_1 = a_4, a_2 = a_5, a_3 = a_6.$$

Now let $(a, b, c) \in F^3$ and let $A = a_7 T^7 + (T^4 + T)(aT^2 + bT + c) + a_0$. From (3.4),

$$A + (a_7 + c)T^7 + a_0 + a = X_1^7 + X_2^7 + X_3^7,$$



Go back

Full Screen

Close

Quit



where $X_1, X_2, X_3 \in F[T]$ have degree ≤ 1 , so that

$$A = ((a_7 + c)T)^7 + (a_0 + a)^7 + X_1^7 + X_2^7 + X_3^7.$$

□

Corollary 3.6. *We have $\mathcal{S}^\times(F, 7) \neq \mathcal{S}(F, 7)$, so that $g(4, 7) = \infty$.*

Proof. Conditions (3.1) are not satisfied by T , so that $\mathcal{S}^\times(F[T], 7) \neq F[T]$. On the other hand, from Paley's theorem, [15], [6, Theorem 1.7], $\mathcal{S}(F[T], 7) = F[T]$. □

Theorem 3.2 is a consequence of the following proposition.

Proposition 3.7. *Let $A \in F[T]$ with degree ≤ 14 , say $A = a_0 + a_1T + \dots + a_{14}T^{14}$.*

(i) *If A is a sum*

$$A = \sum_{i=1}^s (X_i)^7$$

with $X_i \in F[T]$ of degree ≤ 2 , then the coefficients a_1, \dots, a_{13} satisfy (3.2).

(ii) *If $(a_1, \dots, a_{13}) \in F^{13}$ satisfies (3.2), then A is a sum*

$$A = X_1^7 + \dots + X_{11}^7$$

of 11 seventh powers of polynomials X_i with $\deg X_i \leq 2$.

Proof. (i) Suppose that A is a sum

$$A = \sum_{i=1}^s (x_iT^2 + y_iT + z_i)^7$$




Go back

Full Screen

Close

Quit



with $x_i, y_i, z_i \in F$ for $i = 1, \dots, s$. Then,

$$\begin{aligned}
 a_1 + a_4 + a_{10} + a_{13} &= \sum_{i=1}^s y_i(z_i)^3 + \sum_{i=1}^s ((x_i)^2(z_i)^2 + x_i(y_i)^2 z_i + y_i(z_i)^3) \\
 &\quad + \sum_{i=1}^s ((x_i)^2(z_i)^2 + x_i(y_i)^2 z_i + (x_i)^3(y_i)) + \sum_{i=1}^s (x_i)^3 y_i = 0.
 \end{aligned}$$

The proof of the other identities is similar.

(ii) Conversely, suppose that $(a_1, \dots, a_{13}) \in F^{13}$ satisfies (3.2). Proposition 2.3 insures the existence of $(x_1, x_2, y_1, y_2, z_1, z_2) \in F^6$ solution of $(\mathcal{C}(a_{11}, a_{13}, a_9))$ such that $x_1 x_2 y_1 y_2 \neq 0$. For such a solution, we have

$$\begin{cases}
 a_{13} = \sum_{i=1}^2 y_i = \sum_{i=1}^2 x_i^3 y_i, \\
 a_{11} = \sum_{i=1}^2 x_i = \sum_{i=1}^2 x_i y_i^3, \\
 a_9 = \sum_{i=1}^2 (x_i^2 y_i^2 + x_i y_i z_i^2).
 \end{cases}$$

Let

$$\begin{cases}
 a = a_8 + \sum_{i=1}^2 (x_i z_i^3 + x_i^2 y_i z_i + x_i y_i^3), \\
 b = a_{12} + \sum_{i=1}^2 (x_i^3 z_i + x_i^2 y_i^2), \\
 c = a_{10}^2 + \sum_{i=1}^2 (x_i z_i + x_i^2 y_i z_i^2 + x_i^3 y_i^2).
 \end{cases}$$




Go back

Full Screen

Close

Quit



Proposition 2.2 gives the existence of a solution $(x_3, x_4, x_5, z_3, z_4, z_5) \in F^6$ of $(\mathcal{B}_3(a, b, c))$ such that $x_3x_4x_5z_3z_4z_5 \neq 0$. For such a solution, we have

$$\begin{cases} a = \sum_{i=3}^5 x_i = \sum_{i=3}^5 x_i z_i^3, \\ b = \sum_{i=3}^5 z_i = \sum_{i=3}^5 x_i^3 z_i, \\ c^2 = \sum_{i=3}^5 x_i z_i. \end{cases}$$

Let

$$x_6 = a_{14} + \sum_{i=1}^5 x_i, \quad y_3 = y_4 = y_5 = y_6 = z_6 = 0.$$

Thus, we have

$$\begin{cases} a_{12} = \sum_{i=1}^6 (x_i^3 z_i + x_i^2 y_i^2), \\ a_{10} = \sum_{i=1}^6 (x_i^2 z_i^2 + x_i y_i^2 z_i + x_i^3 y_i), \\ a_8 = \sum_{i=1}^6 (x_i z_i^3 + x_i^2 y_i z_i + x_i y_i^3), \end{cases}$$



Go back

Full Screen

Close

Quit



as well as

$$\begin{cases} a_{13} = \sum_{i=1}^6 x_i^3 y_i, \\ a_{11} = \sum_{i=1}^6 x_i y_i^3, \\ a_9 = \sum_{i=1}^6 (x_i^2 y_i^2 + x_i y_i z_i^2). \end{cases}$$

Let

$$(\ddagger) \quad B = A + \sum_{i=1}^6 (x_i T^2 + y_i T + z_i)^7.$$

Then $\deg B \leq 7$. If

$$B = \sum_{i=0}^7 b_i T^i,$$

then,

$$b_4 + b_1 = a_4 + a_1 + \sum_{i=1}^6 (x_i^2 z_i^2 + x_i y_i^2 z_i),$$

$$b_5 + b_2 = a_5 + a_2 + \sum_{i=1}^6 (x_i z_i^3 + x_i^2 y_i z_i),$$

$$b_6 + b_3 = a_6 + a_3 = \sum_{i=1}^6 (x_i^3 z_i + x_i y_i z_i^2).$$



Go back

Full Screen

Close

Quit



Condition (3.2) insures that

$$b_4 + b_1 = a_{13} + a_{10} + \sum_{i=1}^6 (x_i^2 z_i^2 + x_i y_i^2 z_i) = 0,$$

$$b_5 + b_2 = a_{11} + a_8 + \sum_{i=1}^6 (x_i z_i^3 + x_i^2 y_i z_i) = 0,$$

$$b_6 + b_3 = a_{12} + a_9 = \sum_{i=1}^6 (x_i^3 z_i + x_i y_i z_i^2) = 0,$$

so that (3.1) is satisfied by (b_1, \dots, b_6) . Proposition 3.5 gives the existence of polynomials $X_1, \dots, X_5 \in F[T]$ of degree ≤ 1 such that

$$B = \sum_{i=0}^5 X_i^7.$$

We conclude with (‡). □

Theorem 3.3 is a consequence of the following proposition.

Proposition 3.8. *Let*

$$A = \sum_{i=0}^{21} a_i T^i$$

be a polynomial in $F[T]$ with $\deg A \leq 21$. Then, A may be written as a sum

$$A = \sum_i^s (X_i)^7$$




Go back

Full Screen

Close

Quit



with $X_i \in F[T]$ of degree ≤ 3 if and only if its coefficients satisfy the condition

$$(3.3) \quad a_3 + a_6 + a_9 + a_{12} + a_{15} + a_{18} = 0.$$

Moreover, if A satisfies condition (3.3), then A is a sum of 19 seventh powers of polynomials $X_i \in F[T]$ of degree ≤ 3 .

Proof. (I) Let $A = \sum_{i=0}^{21} a_i T^i \in F[T]$. Suppose that A is a sum

$$A = \sum_{i=1}^s (u_i T^3 + x_i T^2 + y_i T + z_i)^7$$

with $u_i, x_i, y_i, z_i \in F$ for $i = 1, \dots, s$. Then we have

$$a_3 + a_6 + a_9 + a_{12} + a_{15} + a_{18} = 0.$$

(II) Let $(a_0, a_1, \dots, a_{20}, a_{21}) \in F^{22}$ satisfying (3.3). We construct a representation of A as a sum of seventh powers of polynomials of degree ≤ 3 in two steps.

(i) *First step* – From Proposition 2.11, there exists

$$(u_1, \dots, u_8, x_1, \dots, x_8, y_1, \dots, y_8, z_1, \dots, z_8) \in F^{32}$$

solution of $(\mathcal{G}(\mathbf{b}))$ with

$$\mathbf{b} = (a_{21}, a_{20}, a_{19}, a_{18}, a_{17} + a_{20}, a_{16}, a_{15}, a_1 + a_4 + a_{10} + a_{13} + a_{16} + a_{19}, \\ a_2 + a_5 + a_8 + a_{11} + a_{17} + a_{20}),$$




Go back

Full Screen

Close

Quit



$u_1 \dots u_8 \neq 0$. Therefore, we have

$$\left\{ \begin{array}{l} \sum_{i=1}^8 u_i = a_{21}, \\ \sum_{i=1}^8 u_i^3 x_i = a_{20}, \\ \sum_{i=1}^8 (u_i^3 y_i + u_i^2 x_i^2) = a_{19}, \\ \sum_{i=1}^8 (u_i^3 z_i + u_i x_i^3) = a_{18}, \\ \sum_{i=1}^8 (u_i^2 y_i^2 + u_i x_i^2 y_i) = a_{17} + a_{20}, \\ \sum_{i=1}^8 (u_i x_i^2 z_i + u_i x_i y_i^2 + u_i^2 x_i^2) = a_{16}, \\ \sum_{i=1}^8 (u_i^2 z_i^2 + u_i y_i^3 + u_i^2 x_i y_i + u_i x_i^3) = a_{15}, \\ \sum_{i=1}^8 (u_i^2 y_i z_i + u_i x_i^2 z_i + u_i y_i z_i^2) = a_1 + a_4 + a_{10} + a_{13} + a_{16} + a_{19}, \\ \sum_{i=1}^8 (u_i^2 x_i z_i + u_i x_i z_i^2 + u_i y_i^2 z_i) = a_2 + a_5 + a_8 + a_{11} + a_{17} + a_{20}, \end{array} \right.$$

so that

$$a_{17} = \sum_{i=1}^8 (u_i^3 x_i + u_i^2 y_i^2 + u_i x_i^2 y_i).$$



Go back

Full Screen

Close

Quit



(ii) *Second step* – Let

$$(\star) \quad B = A + \sum_{i=1}^8 (u_i T^3 + x_i T^2 + y_i T + z_i)^7.$$

Then $\deg B \leq 14$. If

$$B = \sum_{i=0}^{14} b_i T^i,$$

then

$$b_{13} + b_{10} + b_4 + b_1 = a_{13} + a_{10} + a_4 + a_1 + \sum_{i=1}^8 (u_i x_i y_i^2 + u_i^2 y_i z_i + u_i y_i z_i^2 + u_i^3 y_i),$$

$$b_{12} + b_9 + b_6 + b_3 = a_{12} + a_9 + a_6 + a_3 + \sum_{i=1}^8 (u_i^2 x_i y_i + u_i^3 z_i + u_i^2 z_i^2 + u_i y_i^3),$$

$$b_{11} + b_8 + b_5 + b_2 = a_{11} + a_8 + a_5 + a_2 + \sum_{i=1}^8 (u_i^2 x_i z_i + u_i x_i z_i^2 + u_i x_i^2 y_i + u_i y_i^2 z_i + u_i^2 y_i^2)$$

We have

$$a_{19} + a_{16} + a_{13} + a_{10} + a_4 + a_1 = \sum_{i=1}^8 (u_i^2 y_i z_i + u_i x_i^2 z_i + u_i y_i z_i^2),$$



Go back

Full Screen

Close

Quit



and

$$a_{19} + a_{16} = \sum_{i=1}^8 (u_i^3 y_i + u_i x_i^2 z_i + u_i x_i y_i^2),$$

so that

$$a_{13} + a_{10} + a_4 + a_1 = \sum_{i=1}^8 (u_i^2 y_i z_i + u_i y_i z_i^2 + u_i x_i y_i^2 + u_i^3 y_i).$$

Thus,

$$b_{13} + b_{10} + b_4 + b_1 = 0.$$

Similarly, we prove that

$$b_{12} + b_9 + b_6 + b_3 = b_{11} + b_8 + b_5 + b_2 = 0.$$

Proposition 3.7 gives the existence of polynomials $X_1, \dots, X_{11} \in F[T]$ such that

$$B = \sum_{i=1}^{11} X_i^7, \quad \deg X_i \leq 2.$$

We conclude with (\star) . □

Remarks. Proposition 3.7 proves that T is not a sum of seventh powers of polynomials of degree ≤ 2 . From Proposition 3.8 we deduce that every $P \in F[T]$ of degree ≤ 2 may be written as a sum of 19 seventh powers of polynomials of degree ≤ 3 , so that T is a sum of 19 seventh powers. This gives another proof of the equality $\mathcal{S}(F[T], k) = F[T]$. The following proposition gives a representation of T as a sum of 12 seventh powers of polynomials of degree ≤ 3 .




Go back

Full Screen

Close

Quit



Proposition 3.9. *We have*

$$\begin{aligned} T = & (T^3 + T^2 + 1)^7 + (T^3 + T^2 + \alpha T)^7 + (T^3 + T^2 + (\alpha + 1)T)^7 \\ & + (\alpha T^3 + \alpha T^2 + \alpha T + \alpha + 1)^7 + ((\alpha + 1)T^3 + (\alpha + 1)T^2 \\ & + (\alpha + 1)T + \alpha)^7 + (T^2 + T + 1)^7 + (T^2 + T)^7 + (T^2 + \alpha)^7 \\ & + (T^2 + \alpha + 1)^7 + (T + \alpha)^7 + (T + \alpha + 1)^7 + (T + 1)^7. \end{aligned}$$

Proof. An easy verification. □

4. THE FIRST DESCENT

The process described in [1] or in [11] works when a representation of T as sum of k -th powers is known. In the case when $k = 7$ and $q = 4$, this process leads to the following.

Theorem 4.1.

- (i) *Every polynomial $P \in F[T]$ with degree divisible by 7 and ≥ 18599 is a strict sum of 32 seventh powers.*
- (ii) *Every polynomial $P \in F[T]$ with degree ≥ 18593 is a strict sum of 33 seventh powers.*

Proof. Let $P \in F[T]$ with $7(n - 1) < \deg P \leq 7n$. Let

$$\varepsilon(P) = \begin{cases} 0 & \text{if } \deg P = 7n, \\ 1 & \text{if } \deg P < 7n \end{cases}$$

and let

$$H = \varepsilon(P)T^{7n} + P.$$



Go back

Full Screen

Close

Quit



Then, $\deg H = 7n$. From [1, Lemma 5.2], there is a sequence $H_0, H_1, \dots, H_i, \dots$, of polynomials of $F[T]$ of degree $7n_0, 7n_1, \dots, 7n_i$, and a sequence X_0, X_1, \dots, X_i of polynomials of degree n_0, n_1, \dots, n_i , such that $H = H_0$ and such that for each index i ,

$$(4.1) \quad H_i = X_i^7 + H_{i+1},$$

$$(4.2) \quad 6n_i \leq 7n_{i+1} < 6n_i + 7.$$

Moreover, for each index i , there is a polynomial $Y_i \in F[T]$ of degree n_i such that

$$(4.3) \quad \deg(H_i + Y_i^7) < 6n_i.$$

We use (4.1) or (4.3) as long as the sequence (n_i) is decreasing. Let r , if it exists, be the least index such that $3(6n_r - 1) \leq n$. We use identity (1) r times, then we use identity (4.3) once. We get

$$H = X_0^7 + \dots + X_{r-1}^7 + Y_r^7 + R,$$

with $3 \deg R \leq n$. From Proposition 3.9, there exist $R_1, \dots, R_{12} \in F[T]$ of degree $\leq 3 \deg R$ such that

$$R = R_1^7 + \dots + R_{12}^7,$$

so that

$$(4.4) \quad H = X_0^7 + \dots + X_{r-1}^7 + Y_r^7 + R_1^7 + \dots + R_{12}^7$$

with $\deg X_i = n_i \leq n_0 = n$, $\deg Y_r = n_r \leq n_0 = n$, $\deg R_j \leq 3 \deg R \leq n$. Thus, (4.4) is a strict sum of $r + 13$ seventh powers. From (4.2) we get that for $i \geq 1$,

$$7^i n_i \leq 6^i n + \sum_{j=0}^{i-1} 7^j 6^{i-j}.$$



Go back

Full Screen

Close

Quit



Therefore, for any integer $r \geq 1$, we have

$$6n_r - 1 \leq 6 \left(\frac{6}{7}\right)^r n + 35 - 36 \left(\frac{6}{7}\right)^r.$$

For $r \geq 19$, we have $\left(\frac{6}{7}\right)^r < \frac{1}{18}$. Suppose $r = 19$. If $n \geq 2657$, then

$$6 \left(\frac{6}{7}\right)^{19} n + 35 - 36 \left(\frac{6}{7}\right)^{19} \leq \frac{n}{3}.$$

□

5. OTHERS DESCENTS

The second descent process is based on very simple identities.

Proposition 5.1. *The following identity holds in the ring $F[X, Y]$,*

$$(5.1) \quad X^4Y^3 + XY^6 = X^7 + (X + Y)^7 + (X + \alpha Y)^7 + (X + (\alpha + 1)Y)^7.$$

Proof. A simple verification. □

Proposition 5.2. *For a non-negative integer i and $X \in F[T]$, let*


$$(5.2) \quad L_i(X) = X^4T^{3i} + XT^{6i}.$$

Then, the map L_i is \mathbb{F}_2 -linear and we have

$$(5.3) \quad L_i(X) = X^7 + (X + T^i)^7 + (X + \alpha T^i)^7 + (X + (\alpha + 1)T^i)^7.$$

$$(5.4) \quad T^7L_i(X) = L_{i+1}(TX).$$

Proof. Immediate. □



Go back

Full Screen

Close

Quit



Corollary 5.3. *Let n be a non-negative integer and let $a \in F$. Then, we have*

$$(5.5) \quad aT^{4n} = L_0(aT^n) + aT^n,$$

$$(5.6) \quad aT^{4n+3} = L_1(aT^n) + aT^{n+6}.$$

If $n > 0$, then

$$(5.7) \quad aT^{4n+2} = L_2(aT^{n-1}) + aT^{n+11}.$$

If $n > 1$, then

$$(5.8) \quad aT^{4n+1} = L_3(aT^{n-2}) + aT^{n+16}.$$

Proof. (5.5) and (5.6) are immediate. We get (5.7) and (5.8) noting that $aT^{4n+2} = aT^{4(n-1)+6}$ and that $aT^{4n+1} = aT^{4(n-2)+9}$. \square

Roughly speaking, the second descent process uses the following idea. Let $X = x_N T^N + x_{N-1} T^{N-1} + \dots + x_1 T + x_0$ be a polynomial of $F[T]$. Making use of (5.5)–(5.8), we replace a monomial $x_k T^k$ by the sum of an appropriate $L_i(T^j)$ and a monomial of lower degree. We begin with $x_N T^N$ and we follow decreasing degrees as long as the process gives monomials of lower degree. For more details see [4, Proposition 5.4]. Mixing this process with the first descent process leads to the following proposition.

Proposition 5.4. *Let $H \in F[T]$ with degree $7n \geq 112$. Then, there exist $X_0, X_1, X_2, X_3, Y_0, Y_1, Y_2, Y_3, Z \in F[T]$ with $\deg X_i \leq n$, $\deg Y_j \leq n$ and $\deg Z \leq 21$ such that*

$$(5.9) \quad H = X_0^7 + X_1^7 + X_2^7 + X_3^7 + L_0(Y_0) + L_1(Y_1) + L_2(Y_2) + L_3(Y_3) + Z.$$

Proof. See [4, Proposition 5.5]. \square

We continue with other descent processes.



Go back

Full Screen

Close

Quit



Proposition 5.5. *Let $n \geq 3$ be an integer and let $A \in F[T]$ of degree $\leq 7n$. Then there exist $X_1, \dots, X_4 \in F[T]$ of degree $\leq n$ such that*

$$\deg \left(A + \sum_{i=1}^3 X_i^7 \right) \leq 7(n-1),$$

so that there exist $X_1, \dots, X_5 \in F[T]$ of degree $\leq n$ such that

$$\deg \left(A + \sum_{i=1}^4 X_i^7 \right) = 7(n-1).$$

Proof. Let

$$A = \sum_{i=0}^{21} a_i T^i$$

be a polynomial of degree ≤ 21 . Proposition 2.8 gives the existence of

$$(u_1, u_2, u_3, x_1, x_2, x_3, y_1, y_2, y_3, z_1, z_2, z_3) \in F^{16},$$

a solution of $(\mathcal{E}(a_{21}, a_{20}, a_{19}, a_{18}, a_{17} + a_{20}, a_{16}, a_{15}))$ such that $u_1 u_2 u_3 \neq 0$.




Go back

Full Screen

Close

Quit



Therefore, we have

$$\left\{ \begin{array}{l} \sum_{i=1}^3 u_i = a_{21}, \\ \sum_{i=1}^3 u_i^3 x_i = a_{20}, \\ \sum_{i=1}^3 (u_i^3 y_i + u_i^2 x_i^2) = a_{19}, \\ \sum_{i=1}^3 (u_i^3 z_i + u_i x_i^3) = a_{18}, \\ \sum_{i=1}^3 (u_i^3 x_i + u_i^2 y_i^2 + u_i x_i^2 y_i) = a_{17}, \\ \sum_{i=1}^3 (u_i x_i^2 z_i + u_i x_i y_i^2 + u_i^2 x_i^2) = a_{16}, \\ \sum_{i=1}^3 (u_i^2 z_i^2 + u_i y_i^3 + u_i^2 x_i y_i + u_i x_i^3) = a_{15}. \end{array} \right.$$

Let

$$B = \left(A + \sum_{i=1}^3 (u_i T^3 + x_i T^2 + y_i T + z_i)^7 \right),$$

so that

$$\deg B \leq 14.$$

This gives the first part of the proposition in the case when $n = 3$. We get the second part of the proposition in the case $n = 3$ taking

$$X_4 = \begin{cases} 0 & \text{if } \deg B = 14, \\ T^2 & \text{if } \deg B < 14. \end{cases}$$



Go back

Full Screen

Close

Quit



Let $n \geq 3$ be an integer and let $A \in F[T]$ of degree $\leq 7n$. By euclidean division, there is a pair $(Q, R) \in F[T]$ such that, respectively,

$$A = T^{7(n-3)}Q + R, \quad \deg Q \leq 21, \quad \deg R < 7(n-3).$$

There exist $X_1, \dots, X_3 \in F[T]$ and $X_1, \dots, X_4 \in F[T]$, of degree ≤ 3 such that

$$\deg \left(Q + \sum_{i=1}^3 X_i^7 \right) \leq 14,$$

and

$$\deg \left(Q + \sum_{i=1}^4 X_i^7 \right) = 14,$$

respectively. Therefore,

$$\deg \left(T^{7(n-3)} \left(Q + \sum_{i=1}^3 X_i^7 \right) \right) \leq 7(n-1),$$

and

$$\deg \left(T^{7(n-3)} \left(Q + \sum_{i=1}^4 X_i^7 \right) \right) = 7(n-1),$$

respectively, so that,

$$\deg \left(R + T^{7(n-3)}Q + \sum_{i=1}^3 X_i^7 \right) \leq 7(n-1),$$




Go back

Full Screen

Close

Quit



and

$$\deg \left(R + T^{7(n-3)}Q + \sum_{i=1}^4 X_i^7 \right) = 7(n-1),$$

respectively. □

6. END OF THE PROOF

Proposition 6.1. *Let*

$$A = \sum_{i=0}^{28} a_i T^i$$

be a polynomial in $F[T]$ with $\deg A \leq 28$. Then A is a sum

$$(6.1) \quad A = \sum_{i=1}^3 X_i^7 + \sum_{i=1}^{19} Y_i^7,$$

where $X_1, \dots, X_3, Y_1, \dots, Y_{19}$ are polynomials of $F[T]$ such that $\deg X_i \leq 4$ and $\deg Y_i \leq 3$.

Proof. Set

$$(6.2) \quad \sigma = a_{27} + a_{24} + a_{18} + a_{15} + a_{12} + a_9 + a_6 + a_3.$$

Proposition 2.10 gives the existence of

$$(v_1, \dots, v_3, u_1, \dots, u_3, x_1, \dots, x_3, y_1, \dots, y_3, z_1, \dots, z_3) \in F^{15},$$



Go back

Full Screen

Close

Quit

a solution of $(\mathcal{F}(a_{28}, a_{27}, a_{26}, a_{25}, a_{24}, a_{23}, a_{22}, \sigma))$ such that $v_1 \dots v_4 \neq 0$. For such a solution, we have

$$(6.3) \quad \left\{ \begin{array}{l} a_{28} = \sum_{i=1}^3 v_i, \\ a_{27} = \sum_{i=1}^3 u_i = \sum_{i=1}^3 v_i^3 u_i, \\ a_{26} = \sum_{i=1}^3 (x_i + v_i^2 u_i^2) = \sum_{i=1}^3 (v_i^3 x_i + v_i^2 u_i^2), \\ a_{25} = \sum_{i=1}^3 (y_i + v_i u_i^3) = \sum_{i=1}^3 (v_i^3 y_i + v_i u_i^3), \\ a_{24} = \sum_{i=1}^3 (v_i^2 x_i^2 + v_i u_i^2 x_i + u_i + z_i) \\ \quad = \sum_{i=1}^3 (v_i^2 x_i^2 + v_i u_i^2 x_i + v_i^3 u_i + v_i^3 z_i), \\ a_{23} = \sum_{i=1}^3 (v_i u_i^2 y_i + v_i u_i x_i^2 + v_i^2 u_i^2), \\ a_{22} = \sum_{i=1}^3 (v_i u_i^3 + v_i x_i^3 + v_i^2 y_i^2 + v_i u_i^2 z_i + v_i^2 u_i x_i), \end{array} \right.$$

and

$$(6.4) \quad \sigma = \sum_{i=1}^3 (v_i^2 u_i y_i + v_i u_i y_i^2 + v_i x_i^2 y_i).$$

For $i = 1, 2, 3$, let

$$X_i = v_i T^4 + u_i T^3 + x_i T^2 + y_i T + z_i$$



Go back

Full Screen

Close

Quit



and let

$$B = A + \sum_{i=1}^3 X_i^7.$$

Identities (6.3) show that $\deg B \leq 21$. Set

$$B = \sum_{i=1}^{21} b_i T^i.$$

From (6.2), (6.3) and (6.4),

$$b_{18} + b_{15} + b_{12} + b_9 + b_6 + b_3 = 0,$$

so that from Theorem 3.3, there exist polynomials $Y_1, \dots, Y_{19} \in F[T]$ with degree ≤ 3 such that

$$B = \sum_{i=1}^{19} Y_i^7.$$

□

Corollary 6.2. *Let $A \in F[T]$ be such that $21 < \deg A \leq 28$. Then A is a strict sum of 22 seventh powers.*

Theorem 6.3.

- (i) *Every polynomial $P \in F[T]$ whose degree ≥ 441 is divisible by 7 is a strict sum of 32 seventh powers.*
- (ii) *Every polynomial $P \in F[T]$ with degree ≥ 435 is a strict sum of 33 seventh powers.*
- (iii) *Every polynomial $P \in F[T]$ such that $\deg P \geq 112$ and $\deg P$ is divisible by 7 is a strict sum of 42 seventh powers.*
- (iv) *Every polynomial $P \in F[T]$ with degree ≥ 106 is a strict sum of 43 seventh powers.*



Go back

Full Screen

Close

Quit



Proof. As for the proof of Theorem 4.1, it is sufficient to prove (i) and (iii). Let $H \in F[T]$ of degree $7n$ with $n \geq 16$. From (5.9) and (5.3), we get that there exists $Z \in F[T]$ with $\deg Z \leq 21$ such that $H + Z$ is sum of 20 seventh powers of polynomials of degree $\leq n$. From Proposition 3.9, there exist Z_1, \dots, Z_{12} with $\deg Z_i \leq 63$ such that

$$Z = \sum_{i=1}^{12} Z_i^7.$$

If $n \geq 63$, then H is a strict sum of 32 seventh powers. This proves (i).

From Proposition 6.1, there exist $V_1, \dots, V_{22} \in F[T]$ with $\deg V_i \leq 4 < n$ such that

$$Z = \sum_{i=1}^{22} V_i^7,$$

so that H is a strict sum of 42 seventh powers. This proves (iii). □


Corollary 6.4. *We have*

$$G(4, 7) = G^\times(4, 7) \leq 33.$$

We end the study of the set $\mathcal{S}^\times(F, T)$ dealing with polynomials P such that $29 \leq \deg P \leq 105$.

Proposition 6.5. *Let $A \in F[T]$.*

- (i) *If $29 \leq \deg A \leq 35$, then A is a strict sum of 25 seventh powers.*
- (ii) *If $\deg A = 42$, then A is a strict sum of 26 seventh powers.*
- (iii) *If $35 < \deg A < 42$, then A is a strict sum of 27 seventh powers.*
- (iv) *If $\deg A = 7n$ with $7 \leq n < 14$, then A is a strict sum of $n + 20$ seventh powers.*
- (v) *If $7n - 7 < \deg A < 7n$ with $7 \leq n < 14$, then A is a strict sum of $n + 21$ seventh powers.*
- (vi) *If $\deg A = 7n$ with $14 \leq n < 21$, then A is a strict sum of $n + 19$ seventh powers.*



Go back

Full Screen

Close

Quit



- (vii) If $7n - 7 < \deg A < 7n$ with $14 \leq n < 21$, then A is a strict sum of $n + 20$ seventh powers.
(viii) If $\deg A = 7n$ with $21 \leq n < 28$, then A is a strict sum of $n + 18$ seventh powers.
(ix) If $7n - 7 < \deg A < 7n$ with $21 \leq n < 28$, then A is a strict sum of $n + 19$ seventh powers.

Proof. As observed before, it suffices to prove (i), (ii), (iv), (vi) and (viii).

1. Suppose that $29 \leq \deg A \leq 35$. From Proposition 5.5, there exist $X_1, X_2, X_3 \in F[T]$ of degree ≤ 5 such that $\deg(A + \sum_{i=1}^3 X_i^7) \leq 28$. From Proposition 6.1, there exist $Y_1, \dots, Y_{22} \in F[T]$ of degree ≤ 4 such that

$$A + \sum_{i=1}^3 X_i^7 = \sum_{j=1}^{22} Y_j^7.$$

2. Suppose that $\deg A = 42$. From [1, Lemma 5.2-(i)], there is a polynomial $X \in F[T]$ of degree 6 such that $\deg(A + X^7) \leq 35$. From above, there exist $Y_1, \dots, Y_{25} \in F[T]$ of degree ≤ 5 such that

$$A + X^7 = \sum_{j=1}^{25} Y_j^7.$$

3. We prove (iv), (vi) and (viii) by induction. Suppose that for $n \geq 7$, every polynomial of degree $7k$ with $k < n$ is a strict sum of $s(k)$ seventh powers. Let $A \in F[T]$ of degree $7n$. From [1, Lemma 5.2-(ii)], there is a polynomial $X \in F[T]$ of degree n such that $\deg(A + X^7) = 7m(n)$ with $m(n)$ defined by the condition $6n \leq 7m(n) < 6n + 7$. We have

$$m(n) = \begin{cases} n - 1 & \text{if } 7 \leq n \leq 13, \\ n - 2 & \text{if } 14 \leq n \leq 20 \\ n - 3 & \text{if } 21 \leq n \leq 27. \end{cases}$$



Go back

Full Screen

Close

Quit



The induction hypothesis gives that $A + X^7$ is a strict sum of $s(m(n))$ seventh powers, so that A is a strict sum of $s(m(n)) + 1$ seventh powers. We have $s(6) = 26$. Thus,

$$s(n) = \begin{cases} n + 20 & \text{if } 7 \leq n \leq 13, \\ n + 19 & \text{if } 14 \leq n \leq 20, \\ n + 18 & \text{if } 21 \leq n \leq 27. \end{cases}$$

□

Proposition 6.6. *We have*

$$\mathcal{S}^\times(F[T], 7) = \mathcal{A}_1 \cup \mathcal{A}_2 \cup \mathcal{A}_3 \cup \mathcal{A}_\infty,$$

where

- (i) \mathcal{A}_1 is the set of polynomials $A = \sum_{n=0}^7 a_n T^n \in F[T]$ such that $a_1 = a_4, a_2 = a_5, a_3 = a_6$,
- (ii) \mathcal{A}_2 is the set of polynomials $A = \sum_{n=0}^{14} a_n T^n \in F[T]$ with $7 < \deg A \leq 14$ such that

$$\begin{cases} a_1 + a_4 + a_{10} + a_{13} = 0, \\ a_2 + a_5 + a_8 + a_{11} = 0, \\ a_3 + a_6 + a_9 + a_{12} = 0, \end{cases}$$


- (iii) \mathcal{A}_3 is the set of polynomials $A = \sum_{n=0}^{21} a_n T^n \in F[T]$ with $14 < \deg A \leq 21$ such that

$$a_3 + a_6 + a_9 + a_{12} + a_{15} + a_{18} = 0,$$

- (iv) $\mathcal{A}_\infty = \{A \in F[T] \mid \deg A > 21\}$.

Proof. With Theorems 3.1, 3.2, 3.3, Corollary 6.2 and Theorem 6.3.

□




Go back

Full Screen

Close

Quit



Theorem 6.7. *We have*

$$g^{\times}(4, 7) \leq 43.$$

Proof. From Theorems 3.1, 3.2, 3.3, every polynomial $A \in \mathcal{S}^{\times}(F[T], 7)$ of degree ≤ 21 is a strict sum of 19 seventh powers. From Corollary 6.2 and Proposition 6.5, every polynomial $A \in \mathcal{S}^{\times}(F[T], 7)$ such that $21 < \deg A \leq 175$ is a strict sum of 43 seventh powers. From Theorem 6.3, every polynomial $A \in \mathcal{S}^{\times}(F[T], 7)$ such that $\deg A \geq 106$ is a strict sum of 43 seventh powers. \square

1. Car M., *New Bounds on Some Parameters in the Waring Problem for polynomials over a finite field*, Contemporary Mathematics **461** (2008), 59–77.
2. ———, *Sums of fourth powers of polynomials over a finite field of characteristic 3*, Func. and Approx., **38(2)** (2008), 195–220.
3. ———, *Sums of $(2^r + 1)$ -th powers in the polynomial ring $\mathbb{F}_{2^m}[T]$* , Port. Math. (N.S), **67(1)** (2010), 13–56.
4. ———, *Sums of seventh powers in the polynomial ring $\mathbb{F}_{2^m}[T]$* , Port. Math. (N.S), **68(3)** (2011), 297–316.
5. Car M. and Gallardo L. H., *Sums of cubes of polynomials*, Acta Arith. **112** (2004), 41–50.
6. Effinger G. and Hayes D. R., *Additive Number Theory of Polynomials Over a Finite Field*, Oxford Mathematical Monographs, Clarendon Press, Oxford 1991.
7. Ellison W. J., *Waring's problem*, Amer. Math. Monthly **78(1)** (1971), 10–36.
8. Gallardo L. H., *On the restricted Waring problem over $\mathbb{F}_{2^n}[t]$* , Acta Arith. **42** (2000), 109–113.
9. ———, *Every sum of cubes in $\mathbb{F}_4[t]$ is a strict sum of 6 cubes*, Port. Math. **65(2)** (2008), 227–236.
10. Gallardo L. H. and Heath-Brown T.R., *Every sum of cubes in $\mathbb{F}_2[t]$ is a strict sum of 6 cubes*, Finite Fields and App., **13(4)** (2007), 977–980.
11. Gallardo L. H. and Vaserstein L. N., *The strict Waring problem for polynomials rings*, J. Number Theory, **128** 2008, 2963–2972.
12. Kubota R. M., *Waring's problem for $\mathbb{F}_q[x]$* , Dissertationes Math. (Rozprawy Mat.) **117** 1974.
13. Liu Y.-R. and Wooley T., *The unrestricted variant of Waring's problem in function fields*. Funct. Approx. Comment. Math., **37** (2007), part 2, 285–291.
14. ———, *Waring's problem in function fields*, J. Reine Angew. Math., **638** (2010), 1–67.



Go back

Full Screen

Close

Quit



15. Paley R. E. A. C., *Theorems on polynomials in a Galois field*, Quarterly J. of Math., **4** (1933), 52–63.
16. Vaserstein L. N., *Waring's problem for algebras over fields*, J. Number Theory **26** (1987), 286–298.
17. ———, *Ramsey's Theorem and Waring's Problem*, In The Arithmetic of Function Fields (eds. D. Goss and al.), de Gruyter, NewYork-Berlin, 1992.
18. Vaughan R. C. and Wooley T. D., *Waring's problem: a survey*. In Number Theory for the millenium, III Urbana, IL, 2000, 301–240.
19. Webb W. A., *Waring's problem in $GF[q, x]$* , Acta Arith., **22**, (1973), 207–220.

Mireille CAR, Aix-Marseille Université Faculté des Sciences et Techniques, Case cour A Avenue Escadrille Normandie-Niemen, F-13397 Marseille Cedex 20, France,
e-mail: mireille.car@univ-amu.fr



Go back

Full Screen

Close

Quit