

## Review Article

# Solution and Attractivity for a Rational Recursive Sequence

**E. M. Elsayed<sup>1,2</sup>**

<sup>1</sup> *Mathematics Department, Faculty of Science, King Abdulaziz University, P. O. Box 80203, Jeddah 21589, Saudi Arabia*

<sup>2</sup> *Department of Mathematics, Faculty of Science, Mansoura University, Mansoura 35516, Egypt*

Correspondence should be addressed to E. M. Elsayed, [emmelsayed@yahoo.com](mailto:emmelsayed@yahoo.com)

Received 15 February 2011; Accepted 26 March 2011

Academic Editor: Ibrahim Yalcinkaya

Copyright © 2011 E. M. Elsayed. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper is concerned with the behavior of solution of the nonlinear difference equation  $x_{n+1} = ax_{n-1} + bx_n x_{n-1} / (cx_n + dx_{n-2})$ ,  $n = 0, 1, \dots$ , where the initial conditions  $x_{-2}, x_{-1}, x_0$  are arbitrary positive real numbers and  $a, b, c, d$  are positive constants. Also, we give specific form of the solution of four special cases of this equation.

## 1. Introduction

In this paper we deal with the behavior of the solution of the following difference equation:

$$x_{n+1} = ax_{n-1} + \frac{bx_n x_{n-1}}{cx_n + dx_{n-2}}, \quad n = 0, 1, \dots, \quad (1.1)$$

where the initial conditions  $x_{-2}, x_{-1}, x_0$  are arbitrary positive real numbers and  $a, b, c, d$  are positive constants. Also, we obtain the solution of some special cases of (1.1).

Let us introduce some basic definitions and some theorems that we need in the sequel.

Let  $I$  be some interval of real numbers and let

$$f : I^{k+1} \longrightarrow I, \quad (1.2)$$

be a continuously differentiable function. Then for every set of initial conditions  $x_{-k}, x_{-k+1}, \dots, x_0 \in I$ , the difference equation

$$x_{n+1} = f(x_n, x_{n-1}, \dots, x_{n-k}), \quad n = 0, 1, \dots, \quad (1.3)$$

has a unique solution  $\{x_n\}_{n=-k}^{\infty}$  [1].

*Definition 1.1* (equilibrium point). A point  $\bar{x} \in I$  is called an equilibrium point of (1.3) if

$$\bar{x} = f(\bar{x}, \bar{x}, \dots, \bar{x}). \quad (1.4)$$

That is,  $x_n = \bar{x}$  for  $n \geq 0$ , is a solution of (1.3), or equivalently,  $\bar{x}$  is a fixed point of  $f$ .

*Definition 1.2* (stability). (i) The equilibrium point  $\bar{x}$  of (1.3) is locally stable if for every  $\epsilon > 0$ , there exists  $\delta > 0$  such that for all  $x_{-k}, x_{-k+1}, \dots, x_{-1}, x_0 \in I$  with

$$|x_{-k} - \bar{x}| + |x_{-k+1} - \bar{x}| + \dots + |x_0 - \bar{x}| < \delta, \quad (1.5)$$

we have

$$|x_n - \bar{x}| < \epsilon \quad \forall n \geq -k. \quad (1.6)$$

(ii) The equilibrium point  $\bar{x}$  of (1.3) is locally asymptotically stable if  $\bar{x}$  is locally stable solution of (1.3) and there exists  $\gamma > 0$ , such that for all  $x_{-k}, x_{-k+1}, \dots, x_{-1}, x_0 \in I$  with

$$|x_{-k} - \bar{x}| + |x_{-k+1} - \bar{x}| + \dots + |x_0 - \bar{x}| < \gamma, \quad (1.7)$$

we have

$$\lim_{n \rightarrow \infty} x_n = \bar{x}. \quad (1.8)$$

(iii) The equilibrium point  $\bar{x}$  of (1.3) is global attractor if for all  $x_{-k}, x_{-k+1}, \dots, x_{-1}, x_0 \in I$ , we have

$$\lim_{n \rightarrow \infty} x_n = \bar{x}. \quad (1.9)$$

(iv) The equilibrium point  $\bar{x}$  of (1.3) is globally asymptotically stable if  $\bar{x}$  is locally stable, and  $\bar{x}$  is also a global attractor of (1.3).

(v) The equilibrium point  $\bar{x}$  of (1.3) is unstable if  $\bar{x}$  is not locally stable. The linearized equation of (1.3) about the equilibrium  $\bar{x}$  is the linear difference equation

$$y_{n+1} = \sum_{i=0}^k \frac{\partial f(\bar{x}, \bar{x}, \dots, \bar{x})}{\partial x_{n-i}} y_{n-i}. \quad (1.10)$$

**Theorem A** (see [2]). Assume that  $p, q \in \mathbb{R}$  and  $k \in \{0, 1, 2, \dots\}$ . Then

$$|p| + |q| < 1, \quad (1.11)$$

is a sufficient condition for the asymptotic stability of the difference equation

$$x_{n+1} + px_n + qx_{n-k} = 0, \quad n = 0, 1, \dots \quad (1.12)$$

*Remark 1.3.* Theorem A can be easily extended to a general linear equations of the form

$$x_{n+k} + p_1x_{n+k-1} + \dots + p_kx_n = 0, \quad n = 0, 1, \dots, \quad (1.13)$$

where  $p_1, p_2, \dots, p_k \in \mathbb{R}$  and  $k \in \{1, 2, \dots\}$ . Then (1.13) is asymptotically stable provided that

$$\sum_{i=1}^k |p_i| < 1. \quad (1.14)$$

Consider the following equation

$$x_{n+1} = g(x_n, x_{n-1}, x_{n-2}). \quad (1.15)$$

The following theorem will be useful for the proof of our results in this paper.

**Theorem B** (see [1]). Let  $[a, b]$  be an interval of real numbers and assume that

$$g : [a, b]^3 \longrightarrow [a, b], \quad (1.16)$$

is a continuous function satisfying the following properties:

- (a)  $g(x, y, z)$  is nondecreasing in  $x$  and  $y$  in  $[a, b]$  for each  $z \in [a, b]$ , and is nonincreasing in  $z \in [a, b]$  for each  $x$  and  $y$  in  $[a, b]$ ;
- (b) if  $(m, M) \in [a, b] \times [a, b]$  is a solution of the system

$$M = g(M, M, m), \quad m = g(m, m, M), \quad (1.17)$$

then

$$m = M. \quad (1.18)$$

Then (1.15) has a unique equilibrium  $\bar{x} \in [a, b]$  and every solution of (1.15) converges to  $\bar{x}$ .

**Definition 1.4** (periodicity). A sequence  $\{x_n\}_{n=-k}^{\infty}$  is said to be periodic with period  $p$  if  $x_{n+p} = x_n$  for all  $n \geq -k$ .

*Definition 1.5* (Fibonacci sequence). The sequence  $\{F_m\}_{m=0}^{\infty} = \{1, 2, 3, 5, 8, 13, \dots\}$ , that is,  $F_m = F_{m-1} + F_{m-2}$ ,  $m \geq 0$ ,  $F_{-2} = 0$ ,  $F_{-1} = 1$  is called Fibonacci sequence.

Recently there has been a great interest in studying the qualitative properties of rational difference equations. Some prototypes for the development of the basic theory of the global behavior of nonlinear difference equations of order greater than one come from the results for rational difference equations.

However, there have not been any effective general methods to deal with the global behavior of rational difference equations of order greater than one so far. From the known work, one can see that it is extremely difficult to understand thoroughly the global behaviors of solutions of rational difference equations although they have simple forms (or expressions). One can refer to [3–23] for examples to illustrate this. Therefore, the study of rational difference equations of order greater than one is worth further consideration.

Many researchers have investigated the behavior of the solution of difference equations, for example, Aloqeili [24] has obtained the solutions of the difference equation

$$x_{n+1} = \frac{x_{n-1}}{a - x_n x_{n-1}}. \quad (1.19)$$

Amleh et al. [25] studied the dynamics of the difference equation

$$x_{n+1} = \frac{a + bx_{n-1}}{A + Bx_{n-2}}. \quad (1.20)$$

Çinar [26, 27] got the solutions of the following difference equation

$$x_{n+1} = \frac{x_{n-1}}{\pm 1 + ax_n x_{n-1}}. \quad (1.21)$$

In [28], Elabbasy et al. investigated the global stability and periodicity character and gave the solution of special case of the following recursive sequence

$$x_{n+1} = ax_n - \frac{bx_n}{cx_n - dx_{n-1}}. \quad (1.22)$$

Elabbasy et al. [29] investigated the global stability, boundedness, and periodicity character and gave the solution of some special cases of the difference equation

$$x_{n+1} = \frac{\alpha x_{n-k}}{\beta + \gamma \prod_{i=0}^k x_{n-i}}. \quad (1.23)$$

In [30], Ibrahim got the form of the solution of the rational difference equation

$$x_{n+1} = \frac{x_n x_{n-2}}{x_{n-1}(a + b x_n x_{n-2})}. \quad (1.24)$$

Karatas et al. [31] got the solution of the difference equation

$$x_{n+1} = \frac{x_{n-5}}{1 + x_{n-2} x_{n-5}}. \quad (1.25)$$

Yalçinkaya and Çinar [32] considered the dynamics of the difference equation

$$x_{n+1} = \frac{a x_{n-k}}{b + c x_n^p}. \quad (1.26)$$

Yang [33] investigated the global asymptotic stability of the difference equation

$$x_{n+1} = \frac{x_{n-1} x_{n-2} + x_{n-3} + a}{x_{n-1} + x_{n-2} x_{n-3} + a}. \quad (1.27)$$

See also [1, 2, 30, 31, 34–40]. Other related results on rational difference equations can be found in [32, 33, 41–48].

## 2. Local Stability of (1.1)

In this section we investigate the local stability character of the solutions of (1.1). Equation (1.1) has a unique equilibrium point and is given by

$$\bar{x} = a\bar{x} + \frac{b\bar{x}^2}{c\bar{x} + d\bar{x}}, \quad (2.1)$$

or

$$\bar{x}^2(1 - a)(c + d) = b\bar{x}^2, \quad (2.2)$$

if  $(c + d)(1 - a) \neq b$ , then the unique equilibrium point is  $\bar{x} = 0$ .

Let  $f : (0, \infty)^3 \rightarrow (0, \infty)$  be a function defined by

$$f(u, v, w) = av + \frac{buw}{cu + dw}. \quad (2.3)$$

Therefore it follows that

$$f_u(u, v, w) = \frac{bdvw}{(cu + dw)^2}, \quad f_v(u, v, w) = a + \frac{bu}{cu + dw}, \quad f_w(u, v, w) = \frac{-bdvw}{(cu + dw)^2}, \quad (2.4)$$

we see that

$$f_u(\bar{x}, \bar{x}, \bar{x}) = \frac{bd}{(c+d)^2}, \quad f_v(\bar{x}, \bar{x}, \bar{x}) = a + \frac{b}{c+d}, \quad f_w(\bar{x}, \bar{x}, \bar{x}) = \frac{-bd}{(c+d)^2}. \quad (2.5)$$

The linearized equation of (1.1) about  $\bar{x}$  is

$$y_{n+1} - \frac{bd}{(c+d)^2}y_n - \left(a + \frac{b}{c+d}\right)y_{n-1} + \frac{bd}{(c+d)^2}y_{n-1} = 0. \quad (2.6)$$

**Theorem 2.1.** *Assume that*

$$b(c+3d) < (1-a)(c+d)^2. \quad (2.7)$$

*Then the equilibrium point of (1.1) is locally asymptotically stable.*

*Proof.* It follows from Theorem A that (2.6) is asymptotically stable if

$$\left| \frac{bd}{(c+d)^2} \right| + \left| a + \frac{b}{c+d} \right| + \left| \frac{bd}{(c+d)^2} \right| < 1, \quad (2.8)$$

or

$$a + \frac{b}{c+d} + \frac{2bd}{(c+d)^2} < 1, \quad (2.9)$$

and so,

$$\frac{b(c+3d)}{(c+d)^2} < (1-a). \quad (2.10)$$

The proof is complete.  $\square$

### 3. Global Attractor of the Equilibrium Point of (1.1)

In this section we investigate the global attractivity character of solutions of (1.1).

**Theorem 3.1.** *The equilibrium point  $\bar{x}$  of (1.1) is global attractor if  $c(1-a) \neq b$ .*

*Proof.* Let  $p, q$  be real numbers and assume that  $g : [p, q]^3 \rightarrow [p, q]$  is a function defined by  $g(u, v, w) = av + buv/(cu + dw)$ , then we can easily see that the function  $g(u, v, w)$  is increasing in  $u, v$  and decreasing in  $w$ . Suppose that  $(m, M)$  is a solution of the system

$$M = g(M, M, m), \quad m = g(m, m, M). \quad (3.1)$$

Then from (1.1), we see that

$$M = aM + \frac{bM^2}{cM + dm}, \quad m = am + \frac{bm^2}{cm + dM}, \quad (3.2)$$

or

$$M(1 - a) = \frac{bM^2}{cM + dm}, \quad m(1 - a) = \frac{bm^2}{cm + dM}, \quad (3.3)$$

then

$$d(1 - a)Mm + c(1 - a)M^2 = bM^2, \quad d(1 - a)Mm + c(1 - a)m^2 = bm^2, \quad (3.4)$$

subtracting, we obtain

$$c(1 - a)(M^2 - m^2) = b(M^2 - m^2), \quad c(1 - a) \neq b. \quad (3.5)$$

Thus

$$M = m. \quad (3.6)$$

It follows from Theorem B that  $\bar{x}$  is a global attractor of (1.1), and then the proof is complete.  $\square$

#### 4. Boundedness of Solutions of (1.1)

In this section we study the boundedness of solutions of (1.1).

**Theorem 4.1.** *Every solution of (1.1) is bounded if  $(a + b/c) < 1$ .*

*Proof.* Let  $\{x_n\}_{n=-2}^{\infty}$  be a solution of (1.1). It follows from (1.1) that

$$x_{n+1} = ax_{n-1} + \frac{bx_n x_{n-1}}{cx_n + dx_{n-2}} \leq ax_{n-1} + \frac{bx_n x_{n-1}}{cx_n} = \left(a + \frac{b}{c}\right)x_{n-1}. \quad (4.1)$$

Then

$$x_{n+1} \leq x_{n-1} \quad \forall n \geq 0. \quad (4.2)$$

Then the subsequences  $\{x_{2n-1}\}_{n=0}^{\infty}$ ,  $\{x_{2n}\}_{n=0}^{\infty}$  are decreasing and so are bounded from above by  $M = \max\{x_{-2}, x_{-1}, x_0\}$ .  $\square$

## 5. Special Cases of (1.1)

Our goal in this section is to find a specific form of the solutions of some special cases of (1.1) when  $a, b, c$ , and  $d$  are integers and give numerical examples of each case and draw it by using MATLAB 6.5.

### 5.1. On the Difference Equation $x_{n+1} = x_{n-1} + x_n x_{n-1} / (x_n + x_{n-2})$

In this subsection we study the following special case of (1.1):

$$x_{n+1} = x_{n-1} + \frac{x_n x_{n-1}}{x_n + x_{n-2}}, \quad (5.1)$$

where the initial conditions  $x_{-2}, x_{-1}, x_0$  are arbitrary positive real numbers.

**Theorem 5.1.** *Let  $\{x_n\}_{n=-2}^{\infty}$  be a solution of (5.1). Then for  $n = 0, 1, 2, \dots$*

$$x_{2n-1} = k \prod_{i=0}^{n-1} \left( \frac{F_{4i+3}h + F_{4i+2}r}{F_{4i+2}h + F_{4i+1}r} \right), \quad x_{2n} = h \prod_{i=0}^{n-1} \left( \frac{F_{4i+5}h + F_{4i+4}r}{F_{4i+4}h + F_{4i+3}r} \right), \quad (5.2)$$

where  $x_{-2} = r, x_{-1} = k, x_0 = h, \{F_m\}_{m=0}^{\infty} = \{0, 1, 1, 2, 3, 5, 8, 13, \dots\}$ .

*Proof.* For  $n = 0$  the result holds. Now suppose that  $n > 0$  and that our assumption holds for  $n - 1, n - 2$ . That is,

$$\begin{aligned} x_{2n-3} &= k \prod_{i=0}^{n-2} \left( \frac{F_{4i+3}h + F_{4i+2}r}{F_{4i+2}h + F_{4i+1}r} \right), & x_{2n-2} &= h \prod_{i=0}^{n-2} \left( \frac{F_{4i+5}h + F_{4i+4}r}{F_{4i+4}h + F_{4i+3}r} \right), \\ x_{2n-4} &= h \prod_{i=0}^{n-3} \left( \frac{F_{4i+5}h + F_{4i+4}r}{F_{4i+4}h + F_{4i+3}r} \right). \end{aligned} \quad (5.3)$$

Now, it follows from (5.1) that

$$\begin{aligned} x_{2n-1} &= x_{2n-3} + \frac{x_{2n-2} x_{2n-3}}{x_{2n-2} + x_{2n-4}} \\ &= k \prod_{i=0}^{n-2} \left( \frac{F_{4i+3}h + F_{4i+2}r}{F_{4i+2}h + F_{4i+1}r} \right) \\ &\quad + \frac{h \prod_{i=0}^{n-2} ((F_{4i+5}h + F_{4i+4}r) / (F_{4i+4}h + F_{4i+3}r)) k \prod_{i=0}^{n-2} ((F_{4i+3}h + F_{4i+2}r) / (F_{4i+2}h + F_{4i+1}r))}{h \prod_{i=0}^{n-2} ((F_{4i+5}h + F_{4i+4}r) / (F_{4i+4}h + F_{4i+3}r)) + h \prod_{i=0}^{n-3} ((F_{4i+5}h + F_{4i+4}r) / (F_{4i+4}h + F_{4i+3}r))} \\ &= k \prod_{i=0}^{n-2} \left( \frac{F_{4i+3}h + F_{4i+2}r}{F_{4i+2}h + F_{4i+1}r} \right) \\ &\quad + \frac{((F_{4n-3}h + F_{4n-4}r) / (F_{4n-4}h + F_{4n-5}r)) k \prod_{i=0}^{n-2} ((F_{4i+3}h + F_{4i+2}r) / (F_{4i+2}h + F_{4i+1}r))}{((F_{4n-3}h + F_{4n-4}r) / (F_{4n-4}h + F_{4n-5}r)) + 1} \end{aligned}$$



$$\begin{aligned}
&= k \prod_{i=0}^{n-2} \left( \frac{F_{4i+3}h + F_{4i+2}r}{F_{4i+2}h + F_{4i+1}r} \right) + \frac{(F_{4n-3}h + F_{4n-4}r)k \prod_{i=0}^{n-2} ((F_{4i+3}h + F_{4i+2}r)/(F_{4i+2}h + F_{4i+1}r))}{F_{4n-3}h + F_{4n-4}r + F_{4n-4}h + F_{4n-5}r} \\
&= k \prod_{i=0}^{n-2} \left( \frac{F_{4i+3}h + F_{4i+2}r}{F_{4i+2}h + F_{4i+1}r} \right) + \frac{(F_{4n-3}h + F_{4n-4}r)k \prod_{i=0}^{n-2} ((F_{4i+3}h + F_{4i+2}r)/(F_{4i+2}h + F_{4i+1}r))}{F_{4n-2}h + F_{4n-3}r} \\
&= k \prod_{i=0}^{n-2} \left( \frac{F_{4i+3}h + F_{4i+2}r}{F_{4i+2}h + F_{4i+1}r} \right) \left( 1 + \frac{F_{4n-3}h + F_{4n-4}r}{F_{4n-2}h + F_{4n-3}r} \right) \\
&= k \prod_{i=0}^{n-2} \left( \frac{F_{4i+3}h + F_{4i+2}r}{F_{4i+2}h + F_{4i+1}r} \right) \left( \frac{F_{4n-1}h + F_{4n-2}r}{F_{4n-2}h + F_{4n-3}r} \right).
\end{aligned} \tag{5.4}$$

Therefore

$$x_{2n-1} = k \prod_{i=0}^{n-1} \left( \frac{F_{4i+3}h + F_{4i+2}r}{F_{4i+2}h + F_{4i+1}r} \right). \tag{5.5}$$

Also, we see from (5.1) that

$$\begin{aligned}
x_{2n} &= x_{2n-2} + \frac{x_{2n-1}x_{2n-2}}{x_{2n-1} + x_{2n-3}} \\
&= h \prod_{i=0}^{n-2} \left( \frac{F_{4i+5}h + F_{4i+4}r}{F_{4i+4}h + F_{4i+3}r} \right) \\
&\quad + \frac{k \prod_{i=0}^{n-1} ((F_{4i+3}h + F_{4i+2}r)/(F_{4i+2}h + F_{4i+1}r)) h \prod_{i=0}^{n-2} ((F_{4i+5}h + F_{4i+4}r)/(F_{4i+4}h + F_{4i+3}r))}{k \prod_{i=0}^{n-1} ((F_{4i+3}h + F_{4i+2}r)/(F_{4i+2}h + F_{4i+1}r)) + k \prod_{i=0}^{n-2} ((F_{4i+3}h + F_{4i+2}r)/(F_{4i+2}h + F_{4i+1}r))} \\
&= h \prod_{i=0}^{n-2} \left( \frac{F_{4i+5}h + F_{4i+4}r}{F_{4i+4}h + F_{4i+3}r} \right) \\
&\quad + \frac{((F_{4n-1}h + F_{4n-2}r)/(F_{4n-2}h + F_{4n-3}r)) h \prod_{i=0}^{n-2} ((F_{4i+5}h + F_{4i+4}r)/(F_{4i+4}h + F_{4i+3}r))}{(F_{4n-1}h + F_{4n-2}r/F_{4n-2}h + F_{4n-3}r) + 1} \\
&= h \prod_{i=0}^{n-2} \left( \frac{F_{4i+5}h + F_{4i+4}r}{F_{4i+4}h + F_{4i+3}r} \right) + \frac{(F_{4n-1}h + F_{4n-2}r) h \prod_{i=0}^{n-2} ((F_{4i+5}h + F_{4i+4}r)/(F_{4i+4}h + F_{4i+3}r))}{F_{4n-1}h + F_{4n-2}r + F_{4n-2}h + F_{4n-3}r} \\
&= h \prod_{i=0}^{n-2} \left( \frac{F_{4i+5}h + F_{4i+4}r}{F_{4i+4}h + F_{4i+3}r} \right) + \frac{(F_{4n-1}h + F_{4n-2}r) h \prod_{i=0}^{n-2} ((F_{4i+5}h + F_{4i+4}r)/(F_{4i+4}h + F_{4i+3}r))}{F_{4n}h + F_{4n-1}r} \\
&= h \prod_{i=0}^{n-2} \left( \frac{F_{4i+5}h + F_{4i+4}r}{F_{4i+4}h + F_{4i+3}r} \right) \left( 1 + \frac{F_{4n-1}h + F_{4n-2}r}{F_{4n}h + F_{4n-1}r} \right) \\
&= h \prod_{i=0}^{n-2} \left( \frac{F_{4i+5}h + F_{4i+4}r}{F_{4i+4}h + F_{4i+3}r} \right) \left( \frac{F_{4n+1}h + F_{4n}r}{F_{4n}h + F_{4n-1}r} \right).
\end{aligned} \tag{5.6}$$




Figure 1

Thus

$$x_{2n} = h \prod_{i=0}^{n-1} \left( \frac{F_{4i+5}h + F_{4i+4}r}{F_{4i+4}h + F_{4i+3}r} \right). \quad (5.7)$$

Hence, the proof is completed.  $\square$

For confirming the results of this section, we consider numerical example for  $x_{-2} = 7, x_{-1} = 6, x_0 = 3$ . (See Figure 1).

## 5.2. On the Difference Equation $x_{n+1} = x_{n-1} + \frac{x_n x_{n-1}}{x_n - x_{n-2}}$

In this subsection we give a specific form of the solutions of the difference equation

$$x_{n+1} = x_{n-1} + \frac{x_n x_{n-1}}{x_n - x_{n-2}}, \quad (5.8)$$

where the initial conditions  $x_{-2}, x_{-1}, x_0$  are arbitrary positive real numbers with  $x_{-2} \neq x_0$ .

**Theorem 5.2.** Let  $\{x_n\}_{n=-2}^{\infty}$  be a solution of (5.8). Then for  $n = 0, 1, 2, \dots$

$$x_{2n-1} = k \prod_{i=0}^{n-1} \left( \frac{F_{2i+3}h - F_{2i+1}r}{F_{2i+1}h - F_{2i-1}r} \right), \quad x_{2n} = h \prod_{i=0}^{n-1} \left( \frac{F_{2i+4}h - F_{2i+2}r}{F_{2i+2}h - F_{2i}r} \right), \quad (5.9)$$

where  $x_{-2} = r, x_{-1} = k, x_0 = h, \{F_m\}_{m=-1}^{\infty} = \{1, 0, 1, 1, 2, 3, 5, 8, 13, \dots\}$ .

*Proof.* For  $n = 0$  the result holds. Now suppose that  $n > 0$  and that our assumption holds for  $n - 1$ ,  $n - 2$ . That is,

$$\begin{aligned} x_{2n-3} &= k \prod_{i=0}^{n-2} \left( \frac{F_{2i+3}h - F_{2i+1}r}{F_{2i+1}h - F_{2i-1}r} \right), & x_{2n-2} &= h \prod_{i=0}^{n-2} \left( \frac{F_{2i+4}h - F_{2i+2}r}{F_{2i+2}h - F_{2i}r} \right), \\ x_{2n-4} &= h \prod_{i=0}^{n-3} \left( \frac{F_{2i+4}h - F_{2i+2}r}{F_{2i+2}h - F_{2i}r} \right). \end{aligned} \quad (5.10)$$

Now, it follows from (5.8) that

$$\begin{aligned} x_{2n-1} &= x_{2n-3} + \frac{x_{2n-2}x_{2n-3}}{x_{2n-2} - x_{2n-4}} \\ &= k \prod_{i=0}^{n-2} \left( \frac{F_{2i+3}h - F_{2i+1}r}{F_{2i+1}h - F_{2i-1}r} \right) \\ &\quad + \frac{h \prod_{i=0}^{n-2} ((F_{2i+4}h - F_{2i+2}r)/(F_{2i+2}h - F_{2i}r)) k \prod_{i=0}^{n-2} ((F_{2i+3}h - F_{2i+1}r)/(F_{2i+1}h - F_{2i-1}r))}{h \prod_{i=0}^{n-2} ((F_{2i+4}h - F_{2i+2}r)/(F_{2i+2}h - F_{2i}r)) - h \prod_{i=0}^{n-3} ((F_{2i+4}h - F_{2i+2}r)/(F_{2i+2}h - F_{2i}r))} \\ &= k \prod_{i=0}^{n-2} \left( \frac{F_{2i+3}h - F_{2i+1}r}{F_{2i+1}h - F_{2i-1}r} \right) \\ &\quad + \frac{((F_{2n}h - F_{2n-2}r)/(F_{2n-2}h - F_{2n-4}r)) k \prod_{i=0}^{n-2} ((F_{2i+3}h - F_{2i+1}r)/(F_{2i+1}h - F_{2i-1}r))}{(F_{2n}h - F_{2n-2}r)/F_{2n-2}h - F_{2n-4}r} - 1 \\ &= k \prod_{i=0}^{n-2} \left( \frac{F_{2i+3}h - F_{2i+1}r}{F_{2i+1}h - F_{2i-1}r} \right) + \frac{(F_{2n}h - F_{2n-2}r) k \prod_{i=0}^{n-2} ((F_{2i+3}h - F_{2i+1}r)/(F_{2i+1}h - F_{2i-1}r))}{F_{2n-1}h - F_{2n-3}r} \\ &= k \prod_{i=0}^{n-2} \left( \frac{F_{2i+3}h - F_{2i+1}r}{F_{2i+1}h - F_{2i-1}r} \right) \left( 1 + \frac{F_{2n}h - F_{2n-2}r}{F_{2n-1}h - F_{2n-3}r} \right) \\ &= k \prod_{i=0}^{n-2} \left( \frac{F_{2i+3}h - F_{2i+1}r}{F_{2i+1}h - F_{2i-1}r} \right) \left( \frac{F_{2n-1}h - F_{2n-3}r + F_{2n}h - F_{2n-2}r}{F_{2n-1}h - F_{2n-3}r} \right) \\ &= k \prod_{i=0}^{n-2} \left( \frac{F_{2i+3}h - F_{2i+1}r}{F_{2i+1}h - F_{2i-1}r} \right) \left( \frac{F_{2n+1}h - F_{2n-1}r}{F_{2n-1}h - F_{2n-3}r} \right). \end{aligned} \quad (5.11)$$

Therefore

$$x_{2n-1} = k \prod_{i=0}^{n-1} \left( \frac{F_{2i+3}h - F_{2i+1}r}{F_{2i+1}h - F_{2i-1}r} \right). \quad (5.12)$$

Also, we see from (5.8) that

$$\begin{aligned}
x_{2n} &= x_{2n-2} + \frac{x_{2n-1}x_{2n-2}}{x_{2n-1} - x_{2n-3}} \\
&= h \prod_{i=0}^{n-2} \left( \frac{F_{2i+4}h - F_{2i+2}r}{F_{2i+2}h - F_{2i}r} \right) \\
&\quad + \frac{k \prod_{i=0}^{n-1} ((F_{2i+3}h - F_{2i+1}r)/(F_{2i+1}h - F_{2i-1}r)) h \prod_{i=0}^{n-2} ((F_{2i+4}h - F_{2i+2}r)/(F_{2i+2}h - F_{2i}r))}{k \prod_{i=0}^{n-1} ((F_{2i+3}h - F_{2i+1}r)/(F_{2i+1}h - F_{2i-1}r)) - k \prod_{i=0}^{n-2} ((F_{2i+3}h - F_{2i+1}r)/(F_{2i+1}h - F_{2i-1}r))} \\
&= h \prod_{i=0}^{n-2} \left( \frac{F_{2i+4}h - F_{2i+2}r}{F_{2i+2}h - F_{2i}r} \right) \\
&\quad + \frac{((F_{2n+1}h - F_{2n-1}r)/(F_{2n-1}h - F_{2n-3}r)) h \prod_{i=0}^{n-2} ((F_{2i+4}h - F_{2i+2}r)/(F_{2i+2}h - F_{2i}r))}{((F_{2n+1}h - F_{2n-1}r)/(F_{2n-1}h - F_{2n-3}r)) - 1} \\
&= h \prod_{i=0}^{n-2} \left( \frac{F_{2i+4}h - F_{2i+2}r}{F_{2i+2}h - F_{2i}r} \right) + \frac{(F_{2n+1}h - F_{2n-1}r) h \prod_{i=0}^{n-2} ((F_{2i+4}h - F_{2i+2}r)/(F_{2i+2}h - F_{2i}r))}{F_{2n+1}h - F_{2n-1}r - F_{2n-1}h + F_{2n-3}r} \\
&= h \prod_{i=0}^{n-2} \left( \frac{F_{2i+4}h - F_{2i+2}r}{F_{2i+2}h - F_{2i}r} \right) + \frac{(F_{2n+1}h - F_{2n-1}r) h \prod_{i=0}^{n-2} ((F_{2i+4}h - F_{2i+2}r)/(F_{2i+2}h - F_{2i}r))}{F_{2n}h - F_{2n-2}r} \\
&= h \prod_{i=0}^{n-2} \left( \frac{F_{2i+4}h - F_{2i+2}r}{F_{2i+2}h - F_{2i}r} \right) \left( 1 + \frac{F_{2n+1}h - F_{2n-1}r}{F_{2n}h - F_{2n-2}r} \right) \\
&= h \prod_{i=0}^{n-2} \left( \frac{F_{2i+4}h - F_{2i+2}r}{F_{2i+2}h - F_{2i}r} \right) \left( \frac{F_{2n+2}h - F_{2n}r}{F_{2n}h - F_{2n-2}r} \right).
\end{aligned} \tag{5.13}$$

Thus

$$x_{2n} = h \prod_{i=0}^{n-1} \left( \frac{F_{2i+4}h - F_{2i+2}r}{F_{2i+2}h - F_{2i}r} \right). \tag{5.14}$$

Hence, the proof is completed.  $\square$

Assume that  $x_{-2} = 0.7, x_{-1} = 1.6, x_0 = 13$ . (See Figure 2), and for  $x_{-2} = 9, x_{-1} = 5, x_0 = 2$ . (See Figure 3).

The following cases can be treated similarly.

### 5.3. On the Difference Equation $x_{n+1} = x_{n-1} - x_n x_{n-1} / (x_n + x_{n-2})$

In this subsection we obtain the solution of the following difference equation

$$x_{n+1} = x_{n-1} - \frac{x_n x_{n-1}}{x_n + x_{n-2}}, \tag{5.15}$$

where the initial conditions  $x_{-2}, x_{-1}, x_0$  are arbitrary positive real numbers.




Figure 2




Figure 3

**Theorem 5.3.** Let  $\{x_n\}_{n=-2}^\infty$  be a solution of (5.15). Then for  $n = 0, 1, 2, \dots$

$$x_{2n-1} = k \prod_{i=0}^{n-1} \left( \frac{F_{2i}h + F_{2i+1}r}{F_{2i+1}h + F_{2i+2}r} \right), \quad x_{2n} = h \prod_{i=0}^{n-1} \left( \frac{F_{2i+1}h + F_{2i+2}r}{F_{2i+2}h + F_{2i+3}r} \right), \quad (5.16)$$

where  $x_{-2} = r$ ,  $x_{-1} = k$ ,  $x_0 = h$ ,  $\{F_m\}_{m=0}^\infty = \{0, 1, 1, 2, 3, 5, 8, 13, \dots\}$ .

Figure 4 shows the solution when  $x_{-2} = 3$ ,  $x_{-1} = 7$ ,  $x_0 = 12$ .




Figure 4




Figure 5

#### 5.4. On the Difference Equation $x_{n+1} = x_{n-1} - x_n x_{n-1} / (x_n - x_{n-2})$

In this subsection we give the solution of the following special case of (1.1)

$$x_{n+1} = x_{n-1} - \frac{x_n x_{n-1}}{x_n - x_{n-2}} \quad (5.17)$$

where the initial conditions  $x_{-2}, x_{-1}, x_0$  are arbitrary real numbers. with  $x_{-2} \neq x_0, x_{-2}, x_{-1}, x_0 \neq 0$ .

**Theorem 5.4.** Let  $\{x_n\}_{n=-2}^{\infty}$  be a solution of (5.17). Then every solution of (5.17) is periodic with period 12. Moreover  $\{x_n\}_{n=-2}^{\infty}$  takes the form

$$\left\{ r, k, h, \frac{kr}{r-h}, h-r, \frac{hk}{r-h}, -r, -k, -h, \frac{kr}{h-r}, -(h-r), \frac{hk}{h-r}, r, k, h, \dots \right\} \quad (5.18)$$

where  $x_{-2} = r, x_{-1} = k, x_0 = h$ .

Figure 5 shows the solution when  $x_{-2} = 3, x_{-1} = 7, x_0 = 2$ .

## 6. Conclusion

This paper discussed global stability, boundedness, and the solutions of some special cases of (1.1). In Section 2 we proved when  $b(c+3d) < (1-a)(c+d)^2$ , (1.1) local stability. In Section 3 we showed that the unique equilibrium of (1.1) is globally asymptotically stable if  $c(1-a) \neq b$ . In Section 4 we proved that the solution of (1.1) is bounded if  $(a+b/c) < 1$ . In Section 5 we gave the form of the solution of four special cases of (1.1) and gave numerical examples of each case and drew them by using Matlab 6.5.

## References

- [1] M. R. S. Kulenović and G. Ladas, *Dynamics of Second Order Rational Difference Equations*, Chapman & Hall/CRC, Boca Raton, Fla, USA, 2001.
- [2] V. L. Kocić and G. Ladas, *Global Behavior of Nonlinear Difference Equations of Higher Order with Applications*, vol. 256 of *Mathematics and Its Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1993.
- [3] N. Battaloglu, C. Cinar, and I. Yalçinkaya, *The Dynamics of the Difference Equation*, vol. 97, *Ars Combinatoria*, 2010.
- [4] C. Çinar, "On the positive solutions of the difference equation  $x_{n+1} = ax_{n-1}/(1+bx_nx_{n-1})$ ," *Applied Mathematics and Computation*, vol. 156, no. 2, pp. 587–590, 2004.
- [5] E. M. Elabbasy, H. El-Metwally, and E. M. Elsayed, "Global attractivity and periodic character of a fractional difference equation of order three," *Yokohama Mathematical Journal*, vol. 53, no. 2, pp. 89–100, 2007.
- [6] E. M. Elabbasy, H. El-Metwally, and E. M. Elsayed, "On the difference equation  $x_{n+1} = (a_0x_n + a_1x_{n-1} + \dots + a_kx_{n-k})/(b_0x_n + b_1x_{n-1} + \dots + b_kx_{n-k})$ ," *Mathematica Bohemica*, vol. 133, no. 2, pp. 133–147, 2008.
- [7] E. M. Elabbasy and E. M. Elsayed, "Dynamics of a rational difference equation," *Chinese Annals of Mathematics. Series B*, vol. 30, no. 2, pp. 187–198, 2009.
- [8] E. M. Elabbasy and E. M. Elsayed, "Global attractivity and periodic nature of a difference equation," *World Applied Sciences Journal*, vol. 12, no. 1, pp. 39–47, 2011.
- [9] H. El-Metwally, E. A. Grove, and G. Ladas, "A global convergence result with applications to periodic solutions," *Journal of Mathematical Analysis and Applications*, vol. 245, no. 1, pp. 161–170, 2000.
- [10] H. El-Metwally, E. A. Grove, G. Ladas, R. Levins, and M. Radin, "On the difference equation  $x_{n+1} = \alpha + x_{n-1}e^{-x_n}$ ," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 47, no. 7, pp. 4623–4634, 2001.
- [11] H. El-Metwally, "Global behavior of an economic model," *Chaos, Solitons and Fractals*, vol. 33, no. 3, pp. 994–1005, 2007.
- [12] H. El-Metwally, E. A. Grove, G. Ladas, and H. D. Voulov, "On the global attractivity and the periodic character of some difference equations," *Journal of Difference Equations and Applications*, vol. 7, no. 6, pp. 837–850, 2001.
- [13] E. M. Elsayed, "On the solution of recursive sequence of order two," *Fasciculi Mathematici*, no. 40, pp. 5–13, 2008.

- [14] E. M. Elsayed, "Dynamics of a recursive sequence of higher order," *Communications on Applied Nonlinear Analysis*, vol. 16, no. 2, pp. 37–50, 2009.
- [15] E. M. Elsayed, "Dynamics of recursive sequence of order two," *Kyungpook Mathematical Journal*, vol. 50, no. 4, pp. 483–497, 2010.
- [16] E. M. Elsayed, "On the difference equation  $x_{n+1} = x_{n-5}/(-1 + x_{n-2}x_{n-5})$ ," *International Journal of Contemporary Mathematical Sciences*, vol. 3, no. 33–36, pp. 1657–1664, 2008.
- [17] E. M. Elsayed, "Qualitative behavior of a difference equation of order three," *Acta Scientiarum Mathematicarum*, vol. 75, no. 1-2, pp. 113–129, 2009.
- [18] E. M. Elsayed, "Qualitative behavior of a rational recursive sequence," *Indagationes Mathematicae. New Series*, vol. 19, no. 2, pp. 189–201, 2008.
- [19] E. Elsayed, "On the global attractivity and the solution of recursive sequence," *Studia Scientiarum Mathematicarum Hungarica*, vol. 47, no. 3, pp. 401–418, 2010.
- [20] E. M. Elsayed, "Qualitative properties for a fourth order rational difference equation," *Acta Applicandae Mathematicae*, vol. 110, no. 2, pp. 589–604, 2010.
- [21] E. M. Elsayed, "On the global attractivity and the periodic character of a recursive sequence," *Opuscula Mathematica*, vol. 30, no. 4, pp. 431–446, 2010.
- [22] E. M. Elsayed, B. Iričanin, and S. Stević, "On the max-type equation," *Ars Combinatoria*, vol. 95, pp. 187–192, 2010.
- [23] A. E. Hamza and R. Khalaf-Allah, "On the recursive sequence  $x_{n+1} = A \prod_{i=1}^k x_{n-2i-1} / (B + C \prod_{i=1}^{k-1} x_{n-2i})$ ," *Computers & Mathematics with Applications*, vol. 56, no. 7, pp. 1726–1731, 2008.
- [24] M. Aloqeili, "Dynamics of a rational difference equation," *Applied Mathematics and Computation*, vol. 176, no. 2, pp. 768–774, 2006.
- [25] A. M. Amleh, V. Kirk, and G. Ladas, "On the dynamics of  $x_{n+1} = (a + bx_{n-1}) / (A + Bx_{n-2})$ ," *Mathematical Sciences Research Hot-Line*, vol. 5, no. 7, pp. 1–15, 2001.
- [26] C. Çinar, "On the positive solutions of the difference equation  $x_{n+1} = x_{n-1} / (1 + ax_n x_{n-1})$ ," *Applied Mathematics and Computation*, vol. 158, no. 3, pp. 809–812, 2004.
- [27] C. Çinar, "On the positive solutions of the difference equation  $x_{n+1} = x_{n-1} / (-1 + ax_n x_{n-1})$ ," *Applied Mathematics and Computation*, vol. 158, no. 3, pp. 809–812, 2004.
- [28] E. M. Elabbasy, H. El-Metwally, and E. M. Elsayed, "On the difference equation  $x_{n+1} = ax_n - bx_n / (cx_n - dx_{n-1})$ ," *Advances in Difference Equations*, vol. 2006, Article ID 82579, 10 pages, 2006.
- [29] E. M. Elabbasy, H. El-Metwally, and E. M. Elsayed, "On the difference equation  $x_{n+1} = \alpha x_{n-k} / (\beta + \gamma \prod_{i=0}^k x_{n-i})$ ," *Journal of Concrete and Applicable Mathematics*, vol. 5, no. 2, pp. 101–113, 2007.
- [30] T. F. Ibrahim, "On the third order rational difference equation  $x_{n+1} = x_n x_{n-2} / (x_{n-1}(a + bx_n x_{n-2}))$ ," *International Journal of Contemporary Mathematical Sciences*, vol. 4, no. 25–28, pp. 1321–1334, 2009.
- [31] R. Karatas, C. Cinar, and D. Simsek, "On positive solutions of the difference equation  $x_{n+1} = x_{n-5} / (1 + x_{n-2}x_{n-5})$ ," *International Journal of Contemporary Mathematical Sciences*, vol. 1, no. 9–12, pp. 495–500, 2006.
- [32] I. Yalçinkaya and C. Çinar, "On the dynamics of the difference equation  $x_{n+1} = ax_{n-k} / (b + cx_n^p)$ ," *Fasciculi Mathematici*, no. 42, pp. 141–148, 2009.
- [33] X. Yang, "On the global asymptotic stability of the difference equation  $x_{n+1} = (x_{n-1}x_{n-2} + x_{n-3} + a) / (x_{n-1} + x_{n-2}x_{n-3} + a)$ ," *Applied Mathematics and Computation*, vol. 171, no. 2, pp. 857–861, 2005.
- [34] M. R. S. Kulenović and Z. Nurkanović, "Global behavior of a three-dimensional linear fractional system of difference equations," *Journal of Mathematical Analysis and Applications*, vol. 310, no. 2, pp. 673–689, 2005.
- [35] W.-T. Li and H.-R. Sun, "Dynamics of a rational difference equation," *Applied Mathematics and Computation*, vol. 163, no. 2, pp. 577–591, 2005.
- [36] A. Rafiq, "Convergence of an iterative scheme due to Agarwal et al," *Rostocker Mathematisches Kolloquium*, no. 61, pp. 95–105, 2006.
- [37] M. Saleh and M. Aloqeili, "On the difference equation  $y_{n+1} = A + y_n / y_{n-k}$  with  $A < 0$ ," *Applied Mathematics and Computation*, vol. 176, no. 1, pp. 359–363, 2006.
- [38] D. Simsek, C. Cinar, and I. Yalçinkaya, "On the recursive sequence  $x_{n+1} = x_{n-3} / (1 + x_{n-1})$ ," *International Journal of Contemporary Mathematical Sciences*, vol. 1, no. 9–12, pp. 475–480, 2006.
- [39] C. Wang and S. Wang, "Global behavior of equilibrium point for A class of fractional difference equation," in *Proceedings of 7th Asian Control Conference (ASCC '09)*, pp. 288–291, Hong Kong, August 2009.



- [40] C.-Y. Wang, S. Wang, and X.-P. Yan, "Global asymptotic stability of 3-species mutualism models with diffusion and delay effects," *Discrete Dynamics in Nature and Society*, vol. 2009, Article ID 317298, 20 pages, 2009.
- [41] C.-Y. Wang, F. Gong, S. Wang, L.-R. Li, and Q.-H. Shi, "Asymptotic behavior of equilibrium point for a class of nonlinear difference equation," *Advances in Difference Equations*, vol. 2009, Article ID 214309, 8 pages, 2009.
- [42] I. Yalçinkaya, "Global asymptotic stability in a rational difference equation," *Selçuk Journal of Applied Mathematics*, vol. 6, no. 2, pp. 59–68, 2005.
- [43] I. Yalçinkaya, "On the difference equation  $x_{n+1} = \alpha + x_{n-2}/x_n^k$ ," *Polytechnica Posnaniensis. Institutum Mathematicum. Fasciculi Mathematici*, no. 42, pp. 133–139, 2009.
- [44] I. Yalçinkaya, "On the difference equation  $x_{n+1} = \alpha + x_{n-m}/x_n^k$ ," *Discrete Dynamics in Nature and Society*, vol. 2008, Article ID 805460, 8 pages, 2008.
- [45] I. Yalçinkaya, "On the global asymptotic stability of a second-order system of difference equations," *Discrete Dynamics in Nature and Society*, vol. 2008, Article ID 860152, 12 pages, 2008.
- [46] I. Yalçinkaya, C. Çinar, and M. Atalay, "On the solutions of systems of difference equations," *Advances in Difference Equations*, vol. 2008, Article ID 143943, 9 pages, 2008.
- [47] E. M. E. Zayed and M. A. El-Moneam, "On the rational recursive sequence  $x_{n+1} = (\alpha + \beta x_n + \gamma x_{n-1}) / (A + \beta x_n + C x_{n-1})$ ," *Communications on Applied Nonlinear Analysis*, vol. 12, no. 4, pp. 15–28, 2005.
- [48] E. M. E. Zayed and M. A. El-Moneam, "On the rational recursive sequence  $x_{n+1} = (\alpha x_n + \beta x_{n-1} + \gamma x_{n-2} + \delta x_{n-3}) / (A x_n + B x_{n-1} + C x_{n-2} + D x_{n-3})$ ," *Communications on Applied Nonlinear Analysis*, vol. 12, pp. 15–28, 2005.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

