

Research Article

Strong Convergence of Modified Halpern Iterations in CAT(0) Spaces

A. Cuntavepanit¹ and B. Panyanak^{1,2}

¹ Department of Mathematics, Faculty of Science, Chiang Mai University, Chiang Mai 50200, Thailand

² Materials Science Research Center, Faculty of Science, Chiang Mai University, Chiang Mai 50200, Thailand

Correspondence should be addressed to B. Panyanak, banchap@chiangmai.ac.th

Received 28 November 2010; Accepted 10 January 2011

Academic Editor: Qamrul Hasan Ansari

Copyright © 2011 A. Cuntavepanit and B. Panyanak. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Strong convergence theorems are established for the modified Halpern iterations of nonexpansive mappings in CAT(0) spaces. Our results extend and improve the recent ones announced by Kim and Xu (2005), Hu (2008), Song and Chen (2008), Saejung (2010), and many others.

1. Introduction

Let C be a nonempty subset of a metric space (X, d) . A mapping $T : C \rightarrow C$ is said to be *nonexpansive* if

$$d(Tx, Ty) \leq d(x, y), \quad \forall x, y \in C. \quad (1.1)$$

A point $x \in C$ is called a fixed point of T if $x = Tx$. We will denote by $F(T)$ the set of fixed points of T . In 1967, Halpern [1] introduced an explicit iterative scheme for a nonexpansive mapping T on a subset C of a Hilbert space by taking any points $u, x_1 \in C$ and defined the iterative sequence $\{x_n\}$ by

$$x_{n+1} = \alpha_n u + (1 - \alpha_n)Tx_n, \quad \text{for } n \geq 1, \quad (1.2)$$

where $\alpha_n \in [0, 1]$. He pointed out that the control conditions: (C1) $\lim_n \alpha_n = 0$ and (C2) $\sum_{n=1}^{\infty} \alpha_n = \infty$ are necessary for the convergence of $\{x_n\}$ to a fixed point of T . Subsequently, many mathematicians worked on the Halpern iterations both in Hilbert and Banach spaces

(see, e.g., [2–11] and the references therein). Among other things, Wittmann [7] proved strong convergence of the Halpern iteration under the control conditions (C1), (C2), and (C4) $\sum_{n=1}^{\infty} |\alpha_{n+1} - \alpha_n| < \infty$ in a Hilbert space. In 2005, Kim and Xu [12] generalized Wittmann's result by introducing a modified Halpern iteration in a Banach space as follows. Let C be a closed convex subset of a uniformly smooth Banach space X , and let $T : C \rightarrow C$ be a nonexpansive mapping. For any points $u, x_1 \in C$, the sequence $\{x_n\}$ is defined by

$$x_{n+1} = \beta_n u + (1 - \beta_n)T(\alpha_n x_n + (1 - \alpha_n)Tx_n), \quad \text{for } n \geq 1, \quad (1.3)$$

where $\{\alpha_n\}$ and $\{\beta_n\}$ are sequences in $[0, 1]$. They proved under the following control conditions:

$$\begin{aligned} (D1) \quad & \lim_n \alpha_n = 0, & \lim_n \beta_n = 0, \\ (D2) \quad & \sum_{n=1}^{\infty} \alpha_n = \infty, & \sum_{n=1}^{\infty} \beta_n = \infty, \\ (D3) \quad & \sum_{n=1}^{\infty} |\alpha_{n+1} - \alpha_n| < \infty, & \sum_{n=1}^{\infty} |\beta_{n+1} - \beta_n| < \infty, \end{aligned} \quad (1.4)$$

that the sequence $\{x_n\}$ converges strongly to a fixed point of T .

The purpose of this paper is to extend Kim-Xu's result to a special kind of metric spaces, namely, CAT(0) spaces. We also prove a strong convergence theorem for another kind of modified Halpern iteration defined by Hu [13] in this setting.

2. CAT(0) Spaces

A metric space X is a CAT(0) space if it is geodesically connected and if every geodesic triangle in X is at least as "thin" as its comparison triangle in the Euclidean plane. The precise definition is given below. It is well known that any complete, simply connected Riemannian manifold having nonpositive sectional curvature is a CAT(0) space. Other examples include Pre-Hilbert spaces (see [14]), \mathbb{R} -trees (see [15]), Euclidean buildings (see [16]), the complex Hilbert ball with a hyperbolic metric (see [17]), and many others. For a thorough discussion of these spaces and of the fundamental role they play in geometry, we refer the reader to Bridson and Haefliger [14].

Fixed point theory in CAT(0) spaces was first studied by Kirk (see [18, 19]). He showed that every nonexpansive (single-valued) mapping defined on a bounded closed convex subset of a complete CAT(0) space always has a fixed point. Since then, the fixed point theory for single-valued and multivalued mappings in CAT(0) spaces has been rapidly developed, and many papers have appeared (see, e.g., [20–31] and the references therein). It is worth mentioning that fixed point theorems in CAT(0) spaces (specially in \mathbb{R} -trees) can be applied to graph theory, biology, and computer science (see, e.g., [15, 32–35]).

Let (X, d) be a metric space. A *geodesic path* joining $x \in X$ to $y \in X$ (or, more briefly, a *geodesic* from x to y) is a map c from a closed interval $[0, l] \subset \mathbb{R}$ to X such that $c(0) = x$, $c(l) = y$ and $d(c(t), c(t')) = |t - t'|$ for all $t, t' \in [0, l]$. In particular, c is an isometry and $d(x, y) = l$. The image α of c is called a *geodesic* (or *metric*) *segment* joining x and y . When it is unique, this geodesic segment is denoted by $[x, y]$. The space (X, d) is said to be a *geodesic space* if every

two points of X are joined by a geodesic, and X is said to be *uniquely geodesic* if there is exactly one geodesic joining x and y for each $x, y \in X$. A subset $Y \subseteq X$ is said to be *convex* if Y includes every geodesic segment joining any two of its points.

A *geodesic triangle* $\Delta(x_1, x_2, x_3)$ in a geodesic metric space (X, d) consists of three points x_1, x_2 , and x_3 in X (the *vertices* of Δ) and a geodesic segment between each pair of vertices (the *edges* of Δ). A *comparison triangle* for the geodesic triangle $\Delta(x_1, x_2, x_3)$ in (X, d) is a triangle $\bar{\Delta}(x_1, x_2, x_3) := \Delta(\bar{x}_1, \bar{x}_2, \bar{x}_3)$ in the Euclidean plane \mathbb{E}^2 such that $d_{\mathbb{E}^2}(\bar{x}_i, \bar{x}_j) = d(x_i, x_j)$ for $i, j \in \{1, 2, 3\}$.

A geodesic space is said to be a CAT(0) space if all geodesic triangles satisfy the following comparison axiom.

CAT(0): let Δ be a geodesic triangle in X , and let $\bar{\Delta}$ be a comparison triangle for Δ . Then, $\bar{\Delta}$ is said to satisfy the CAT(0) *inequality* if for all $x, y \in \Delta$ and all comparison points $\bar{x}, \bar{y} \in \bar{\Delta}$,

$$d(x, y) \leq d_{\mathbb{E}^2}(\bar{x}, \bar{y}). \quad (2.1)$$

Let $x, y \in X$, and by Lemma 2.1 (iv) of [23] for each $t \in [0, 1]$, there exists a unique point $z \in [x, y]$ such that

$$d(x, z) = td(x, y), \quad d(y, z) = (1 - t)d(x, y). \quad (2.2)$$

From now on, we will use the notation $(1 - t)x \oplus ty$ for the unique point z satisfying (2.2). We now collect some elementary facts about CAT(0) spaces which will be used in the proofs of our main results.

Lemma 2.1. *Let X be a CAT(0) space. Then,*

(i) (see [23, Lemma 2.4]) *for each $x, y, z \in X$ and $t \in [0, 1]$, one has*

$$d((1 - t)x \oplus ty, z) \leq (1 - t)d(x, z) + td(y, z), \quad (2.3)$$

(ii) (see [21]) *for each $x, y \in X$ and $t, s \in [0, 1]$, one has*

$$d((1 - t)x \oplus ty, (1 - s)x \oplus sy) = |t - s|d(x, y), \quad (2.4)$$

(iii) (see [19, Lemma 3]) *for each $x, y, z \in X$ and $t \in [0, 1]$, one has*

$$d((1 - t)z \oplus tx, (1 - t)z \oplus ty) \leq td(x, y), \quad (2.5)$$

(iv) (see [23, Lemma 2.5]) *for each $x, y, z \in X$ and $t \in [0, 1]$, one has*

$$d((1 - t)x \oplus ty, z)^2 \leq (1 - t)d(x, z)^2 + td(y, z)^2 - t(1 - t)d(x, y)^2. \quad (2.6)$$

Recall that a continuous linear functional μ on ℓ_∞ , the Banach space of bounded real sequences, is called a *Banach limit* if $\|\mu\| = \mu(1, 1, \dots) = 1$ and $\mu_n(a_n) = \mu_n(a_{n+1})$ for all $\{a_n\} \in \ell_\infty$.

Lemma 2.2 (see [8, Proposition 2]). Let $\{a_1, a_2, \dots\} \in \ell_\infty$ be such that $\mu_n(a_n) \leq 0$ for all Banach limits μ and $\limsup_n (a_{n+1} - a_n) \leq 0$. Then, $\limsup_n a_n \leq 0$.

Lemma 2.3 (see [28, Lemma 2.1]). Let C be a closed convex subset of a complete CAT(0) space X , and let $T : C \rightarrow C$ be a nonexpansive mapping. Let $u \in C$ be fixed. For each $t \in (0, 1)$, the mapping $S_t : C \rightarrow C$ defined by

$$S_t z = tu \oplus (1-t)Tz, \quad \text{for } z \in C \quad (2.7)$$

has a unique fixed point $z_t \in C$, that is,

$$z_t = S_t(z_t) = tu \oplus (1-t)T(z_t). \quad (2.8)$$

Lemma 2.4 (see [28, Lemma 2.2]). Let C and T be as the preceding lemma. Then, $F(T) \neq \emptyset$ if and only if $\{z_t\}$ given by (2.8) remains bounded as $t \rightarrow 0$. In this case, the following statements hold:

- (1) $\{z_t\}$ converges to the unique fixed point z of T which is nearest u ,
- (2) $d^2(u, z) \leq \mu_n d^2(u, x_n)$ for all Banach limits μ and all bounded sequences $\{x_n\}$ with $\lim_n d(x_n, Tx_n) = 0$.

Lemma 2.5 (see [10, Lemma 2.1]). Let $\{\alpha_n\}_{n=1}^\infty$ be a sequence of nonnegative real numbers satisfying the condition

$$\alpha_{n+1} \leq (1 - \gamma_n)\alpha_n + \gamma_n\sigma_n, \quad n \geq 1, \quad (2.9)$$

where $\{\gamma_n\}$ and $\{\sigma_n\}$ are sequences of real numbers such that

- (1) $\{\gamma_n\} \subset [0, 1]$ and $\sum_{n=1}^\infty \gamma_n = \infty$,
- (2) either $\limsup_{n \rightarrow \infty} \sigma_n \leq 0$ or $\sum_{n=1}^\infty |\gamma_n\sigma_n| < \infty$.

Then, $\lim_{n \rightarrow \infty} \alpha_n = 0$.

Lemma 2.6 (see [27, 36]). Let $\{x_n\}$ and $\{y_n\}$ be bounded sequences in a CAT(0) space X , and let $\{\alpha_n\}$ be a sequence in $[0, 1]$ with $0 < \liminf_n \alpha_n \leq \limsup_n \alpha_n < 1$. Suppose that $x_{n+1} = \alpha_n y_n \oplus (1 - \alpha_n)x_n$ for all $n \in \mathbb{N}$ and

$$\limsup_{n \rightarrow \infty} (d(y_{n+1}, y_n) - d(x_{n+1}, x_n)) \leq 0. \quad (2.10)$$

Then, $\lim_n d(x_n, y_n) = 0$.

3. Main Results

The following result is an analog of Theorem 1 of Kim and Xu [12]. They prove the theorem by using the concept of duality mapping, while we use the concept of Banach limit. We also observe that the condition $\sum_{n=1}^\infty \alpha_n = \infty$ in [12, Theorem 1] is superfluous.

Theorem 3.1. *Let C be a nonempty closed convex subset of a complete CAT(0) space X , and let $T : C \rightarrow C$ be a nonexpansive mapping such that $F(T) \neq \emptyset$. Given a point $u \in C$ and sequences $\{\alpha_n\}$ and $\{\beta_n\}$ in $[0, 1]$, the following conditions are satisfied:*

- (A1) $\lim_n \alpha_n = 0$ and $\sum_{n=1}^{\infty} |\alpha_{n+1} - \alpha_n| < \infty$,
 (A2) $\lim_n \beta_n = 0$, $\sum_{n=1}^{\infty} \beta_n = \infty$ and $\sum_{n=1}^{\infty} |\beta_{n+1} - \beta_n| < \infty$.

Define a sequence $\{x_n\}$ in C by $x_1 = x \in C$ arbitrarily, and

$$x_{n+1} = \beta_n u \oplus (1 - \beta_n)(\alpha_n x_n \oplus (1 - \alpha_n)Tx_n), \quad \forall n \geq 1. \quad (3.1)$$

Then, $\{x_n\}$ converges to a fixed point $z \in F(T)$ which is nearest u .

Proof. For each $n \geq 1$, we let $y_n := \alpha_n x_n \oplus (1 - \alpha_n)Tx_n$. We divide the proof into 3 steps. (i) We will show that $\{x_n\}$, $\{y_n\}$, and $\{Tx_n\}$ are bounded sequences. (ii) We show that $\lim_n d(x_n, Tx_n) = 0$. Finally, we show that (iii) $\{x_n\}$ converges to a fixed point $z \in F(T)$ which is nearest u .

(i) As in the first part of the proof of [12, Theorem 1], we can show that $\{x_n\}$ is bounded and so is $\{y_n\}$ and $\{Tx_n\}$. Notice also that

$$d(y_n, p) \leq d(x_n, p), \quad \forall p \in F(T). \quad (3.2)$$

(ii) It suffices to show that

$$\lim_{n \rightarrow \infty} d(x_n, x_{n+1}) = 0. \quad (3.3)$$

Indeed, if (3.3) holds, we obtain

$$\begin{aligned} d(x_n, Tx_n) &\leq d(x_n, x_{n+1}) + d(x_{n+1}, y_n) + d(y_n, Tx_n) \\ &= d(x_n, x_{n+1}) + d(\beta_n u \oplus (1 - \beta_n)y_n, y_n) + d(\alpha_n x_n \oplus (1 - \alpha_n)Tx_n, Tx_n) \\ &\leq d(x_n, x_{n+1}) + \beta_n d(u, y_n) + \alpha_n d(x_n, Tx_n) \rightarrow 0, \quad \text{as } n \rightarrow \infty. \end{aligned} \quad (3.4)$$

By using Lemma 2.1, we get

$$\begin{aligned} d(x_{n+1}, x_n) &= d(\beta_n u \oplus (1 - \beta_n)y_n, \beta_{n-1} u \oplus (1 - \beta_{n-1})y_{n-1}) \\ &\leq d(\beta_n u \oplus (1 - \beta_n)y_n, \beta_n u \oplus (1 - \beta_n)y_{n-1}) \\ &\quad + d(\beta_n u \oplus (1 - \beta_n)y_{n-1}, \beta_{n-1} u \oplus (1 - \beta_{n-1})y_{n-1}) \end{aligned}$$

$$\begin{aligned}
&\leq (1 - \beta_n)d(y_n, y_{n-1}) + |\beta_n - \beta_{n-1}|d(u, y_{n-1}) \\
&= (1 - \beta_n)d(\alpha_n x_n \oplus (1 - \alpha_n)Tx_n, \alpha_{n-1}x_{n-1} \oplus (1 - \alpha_{n-1})Tx_{n-1}) \\
&\quad + |\beta_n - \beta_{n-1}|d(u, \alpha_{n-1}x_{n-1} \oplus (1 - \alpha_{n-1})Tx_{n-1}) \\
&\leq (1 - \beta_n)[d(\alpha_n x_n \oplus (1 - \alpha_n)Tx_n, \alpha_n x_{n-1} \oplus (1 - \alpha_n)Tx_n) \\
&\quad + d(\alpha_n x_{n-1} \oplus (1 - \alpha_n)Tx_n, \alpha_n x_{n-1} \oplus (1 - \alpha_n)Tx_{n-1}) \\
&\quad + d(\alpha_n x_{n-1} \oplus (1 - \alpha_n)Tx_{n-1}, \alpha_{n-1}x_{n-1} \oplus (1 - \alpha_{n-1})Tx_{n-1})] \\
&\quad + |\beta_n - \beta_{n-1}|[\alpha_{n-1}d(u, x_{n-1}) + (1 - \alpha_{n-1})d(u, Tx_{n-1})] \\
&\leq (1 - \beta_n)[\alpha_n d(x_n, x_{n-1}) + (1 - \alpha_n)d(Tx_n, Tx_{n-1}) + |\alpha_n - \alpha_{n-1}|d(x_{n-1}, Tx_{n-1})] \\
&\quad + |\beta_n - \beta_{n-1}|[\alpha_{n-1}d(u, x_{n-1}) + (1 - \alpha_{n-1})d(u, Tx_{n-1})] \\
&= (1 - \beta_n)d(x_n, x_{n-1}) + (1 - \beta_n)|\alpha_n - \alpha_{n-1}|d(x_{n-1}, Tx_{n-1}) \\
&\quad + |\beta_n - \beta_{n-1}|\alpha_{n-1}d(u, x_{n-1}) + |\beta_n - \beta_{n-1}|(1 - \alpha_{n-1})d(u, Tx_{n-1}) \\
&\leq (1 - \beta_n)d(x_n, x_{n-1}) + (1 - \beta_n)|\alpha_n - \alpha_{n-1}|d(x_{n-1}, Tx_{n-1}) \\
&\quad + |\beta_n - \beta_{n-1}|\alpha_{n-1}[d(u, Tx_{n-1}) + d(Tx_{n-1}, x_{n-1})] \\
&\quad + |\beta_n - \beta_{n-1}|d(u, Tx_{n-1}) - |\beta_n - \beta_{n-1}|\alpha_{n-1}d(u, Tx_{n-1}) \\
&= (1 - \beta_n)d(x_n, x_{n-1}) + (1 - \beta_n)|\alpha_n - \alpha_{n-1}|d(x_{n-1}, Tx_{n-1}) \\
&\quad + |\beta_n - \beta_{n-1}|\alpha_{n-1}d(x_{n-1}, Tx_{n-1}) + |\beta_n - \beta_{n-1}|d(u, Tx_{n-1}).
\end{aligned} \tag{3.5}$$

Hence,

$$d(x_{n+1}, x_n) \leq (1 - \beta_n)d(x_n, x_{n-1}) + \gamma(|\alpha_n - \alpha_{n-1}| + 2|\beta_n - \beta_{n-1}|), \tag{3.6}$$

where $\gamma > 0$ is a constant such that $\gamma \geq \max\{d(u, Tx_{n-1}), d(x_{n-1}, Tx_{n-1})\}$ for all $n \in \mathbb{N}$. By assumptions, we have

$$\lim_{n \rightarrow \infty} \beta_n = 0, \quad \sum_{n=1}^{\infty} \beta_n = \infty, \quad \sum_{n=1}^{\infty} (|\alpha_n - \alpha_{n-1}| + 2|\beta_n - \beta_{n-1}|) < \infty. \tag{3.7}$$

Hence, Lemma 2.5 is applicable to (3.6), and we obtain $\lim_n d(x_{n+1}, x_n) = 0$.

(iii) From Lemma 2.3, let $z = \lim_{t \rightarrow 0} z_t$, where z_t is given by (2.8). Then, z is the point of $F(T)$ which is nearest u . We observe that

$$\begin{aligned}
 d^2(x_{n+1}, z) &= d^2(\beta_n u \oplus (1 - \beta_n)y_n, z) \\
 &\leq \beta_n d^2(u, z) + (1 - \beta_n)d^2(y_n, z) - \beta_n(1 - \beta_n)d^2(u, y_n) \\
 &\leq \beta_n d^2(u, z) + (1 - \beta_n)d^2(x_n, z) - \beta_n(1 - \beta_n)d^2(u, y_n) \\
 &= (1 - \beta_n)d^2(x_n, z) + \beta_n [d^2(u, z) - (1 - \beta_n)d^2(u, y_n)].
 \end{aligned} \tag{3.8}$$

By Lemma 2.4, we have $\mu_n(d^2(u, z) - d^2(u, x_n)) \leq 0$ for all Banach limit μ . Moreover, since $\lim_n d(x_{n+1}, x_n) = 0$,

$$\limsup_{n \rightarrow \infty} \left[\left(d^2(u, z) - d^2(u, x_{n+1}) \right) - \left(d^2(u, z) - d^2(u, x_n) \right) \right] = 0. \tag{3.9}$$

It follows from $\lim_n d(y_n, x_n) = 0$ and Lemma 2.2 that

$$\limsup_{n \rightarrow \infty} \left(d^2(u, z) - (1 - \beta_n)d^2(u, y_n) \right) = \limsup_{n \rightarrow \infty} \left(d^2(u, z) - d^2(u, x_n) \right) \leq 0. \tag{3.10}$$

Hence, the conclusion follows from Lemma 2.5. \square

By using the similar technique as in the proof of Theorem 3.1, we can obtain a strong convergence theorem which is an analog of [13, Theorem 3.1] (see also [37, 38] for subsequence comments).

Theorem 3.2. *Let C be a nonempty closed and convex subset of a complete CAT(0) space X , and let $T : C \rightarrow C$ be a nonexpansive mapping such that $F(T) \neq \emptyset$. Given a point $u \in C$ and an initial value $x_1 \in C$. The sequence $\{x_n\}$ is defined iteratively by*

$$x_{n+1} = \beta_n x_n \oplus (1 - \beta_n)(\alpha_n u \oplus (1 - \alpha_n)Tx_n), \quad n \geq 1. \tag{3.11}$$

Suppose that both $\{\alpha_n\}$ and $\{\beta_n\}$ are sequences in $[0, 1]$ satisfying

- (B1) $\lim_{n \rightarrow \infty} \alpha_n = 0$,
- (B2) $\sum_{n=1}^{\infty} \alpha_n = \infty$,
- (B3) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$.

Then, $\{x_n\}$ converges to a fixed point $z \in F(T)$ which is nearest u .

Proof. Let $y_n := \alpha_n u \oplus (1 - \alpha_n)Tx_n$. We divide the proof into 3 steps.

Step 1. We show that $\{x_n\}$, $\{y_n\}$, and $\{Tx_n\}$ are bounded sequences. Let $p \in F(T)$, then we have

$$\begin{aligned}
d(x_{n+1}, p) &= d(\beta_n x_n \oplus (1 - \beta_n)(\alpha_n u \oplus (1 - \alpha_n)Tx_n), p) \\
&\leq \beta_n d(x_n, p) + (1 - \beta_n)d(\alpha_n u \oplus (1 - \alpha_n)Tx_n, p) \\
&\leq \beta_n d(x_n, p) + (1 - \beta_n)\alpha_n d(u, p) + (1 - \beta_n)(1 - \alpha_n)d(Tx_n, p) \\
&\leq (\beta_n + (1 - \beta_n)(1 - \alpha_n))d(x_n, p) + (1 - \beta_n)\alpha_n d(u, p) \\
&= [1 - (1 - \beta_n)\alpha_n]d(x_n, p) + (1 - \beta_n)\alpha_n d(u, p) \\
&\leq \max\{d(x_n, p), d(u, p)\}.
\end{aligned} \tag{3.12}$$

Now, an induction yields

$$d(x_{n+1}, p) \leq \max\{d(x_1, p), d(u, p)\}, \quad n \geq 1. \tag{3.13}$$

Hence, $\{x_n\}$ is bounded and so are $\{y_n\}$ and $\{Tx_n\}$.

Step 2. We show that $\lim_n d(x_n, Tx_n) = 0$. By using Lemma 2.1, we get

$$\begin{aligned}
d(y_{n+1}, y_n) &= d(\alpha_{n+1}u \oplus (1 - \alpha_{n+1})Tx_{n+1}, \alpha_n u \oplus (1 - \alpha_n)Tx_n) \\
&\leq \alpha_n d(\alpha_{n+1}u \oplus (1 - \alpha_{n+1})Tx_{n+1}, u) \\
&\quad + (1 - \alpha_n)d(\alpha_{n+1}u \oplus (1 - \alpha_{n+1})Tx_{n+1}, Tx_n) \\
&\leq \alpha_n(1 - \alpha_{n+1})d(Tx_{n+1}, u) + (1 - \alpha_n)\alpha_{n+1}d(u, Tx_n) \\
&\quad + (1 - \alpha_n)(1 - \alpha_{n+1})d(Tx_{n+1}, Tx_n) \\
&\leq \alpha_n(1 - \alpha_{n+1})d(Tx_{n+1}, u) + (1 - \alpha_n)\alpha_{n+1}d(u, Tx_n) \\
&\quad + (1 - \alpha_n)(1 - \alpha_{n+1})d(x_{n+1}, x_n).
\end{aligned} \tag{3.14}$$

This implies that

$$\begin{aligned}
d(y_{n+1}, y_n) - d(x_{n+1}, x_n) &\leq \alpha_n(1 - \alpha_{n+1})d(Tx_{n+1}, u) + (1 - \alpha_n)\alpha_{n+1}d(u, Tx_n) \\
&\quad + [\alpha_n\alpha_{n+1} - \alpha_n - \alpha_{n+1}]d(x_{n+1}, x_n).
\end{aligned} \tag{3.15}$$

Since $\{x_n\}$ and $\{Tx_n\}$ are bounded and $\lim_{n \rightarrow \infty} \alpha_n = 0$, it follows that

$$\limsup_{n \rightarrow \infty} (d(y_{n+1}, y_n) - d(x_{n+1}, x_n)) \leq 0. \tag{3.16}$$

Hence, by Lemma 2.6, we get

$$\lim_{n \rightarrow \infty} d(x_n, y_n) = 0. \quad (3.17)$$

On the other hand,

$$d(y_n, Tx_n) = d(\alpha_n u \oplus (1 - \alpha_n)Tx_n, Tx_n) \leq \alpha_n d(u, Tx_n) \rightarrow 0, \quad \text{as } n \rightarrow \infty. \quad (3.18)$$

Using (3.17) and (3.18), we get

$$d(x_n, Tx_n) \leq d(x_n, y_n) + d(y_n, Tx_n) \rightarrow 0, \quad \text{as } n \rightarrow \infty. \quad (3.19)$$

Step 3. We show that $\{x_n\}$ converges to a fixed point of T . Let $z = \lim_{t \rightarrow 0} z_t$, where z_t is given by (2.8), then $z \in F(T)$. Finally, we show that $\lim_n x_n = z$

$$\begin{aligned} d^2(x_{n+1}, z) &= d^2(\beta_n x_n \oplus (1 - \beta_n)y_n, z) \\ &\leq \beta_n d^2(x_n, z) + (1 - \beta_n)d^2(y_n, z) - \beta_n(1 - \beta_n)d^2(x_n, y_n) \\ &\leq \beta_n d^2(x_n, z) + (1 - \beta_n)d^2(\alpha_n u \oplus (1 - \alpha_n)Tx_n, z) - \beta_n(1 - \beta_n)d^2(x_n, y_n) \\ &\leq (1 - \beta_n) \left[\alpha_n d^2(u, z) + (1 - \alpha_n)d^2(Tx_n, z) - \alpha_n(1 - \alpha_n)d^2(u, Tx_n) \right] \\ &\quad - \beta_n(1 - \beta_n)d^2(x_n, y_n) + \beta_n d^2(x_n, z) \\ &\leq [\beta_n + (1 - \beta_n)(1 - \alpha_n)]d^2(x_n, z) + (1 - \beta_n)\alpha_n \left[d^2(u, z) - (1 - \alpha_n)d^2(u, Tx_n) \right] \\ &= [1 - (1 - \beta_n)\alpha_n]d^2(x_n, z) + (1 - \beta_n)\alpha_n \left[d^2(u, z) - (1 - \alpha_n)d^2(u, Tx_n) \right]. \end{aligned} \quad (3.20)$$

By Lemma 2.4, we have $\mu_n(d^2(u, z) - d^2(u, x_n)) \leq 0$ for all Banach limit μ . Moreover, since

$$\begin{aligned} d(x_{n+1}, x_n) &= d(\beta_n x_n \oplus (1 - \beta_n)y_n, x_n) \\ &\leq (1 - \beta_n)d(y_n, x_n) \rightarrow 0, \quad \text{as } n \rightarrow \infty, \\ \limsup_{n \rightarrow \infty} \left(d^2(u, z) + d^2(u, x_{n+1}) - d^2(u, z) - d^2(u, x_n) \right) &= 0, \end{aligned} \quad (3.21)$$

it follows from condition (B1), $\lim_n d(x_n, Tx_n) = 0$ and Lemma 2.2 that

$$\limsup_{n \rightarrow \infty} \left(d^2(u, z) - (1 - \alpha_n)d^2(u, Tx_n) \right) = \limsup_{n \rightarrow \infty} \left(d^2(u, z) - d^2(u, x_n) \right) \leq 0. \quad (3.22)$$

Hence, the conclusion follows by Lemma 2.5. \square

Acknowledgments

The authors are grateful to Professor Sompong Dhompongsa for his suggestions and advices during the preparation of the paper. This research was supported by the National Research University Project under Thailand's Office of the Higher Education Commission.

References

- [1] B. Halpern, "Fixed points of nonexpanding maps," *Bulletin of the American Mathematical Society*, vol. 73, pp. 957–961, 1967.
- [2] S. Reich, "Some fixed point problems," *Atti della Accademia Nazionale dei Lincei*, vol. 57, no. 3-4, pp. 194–198, 1974.
- [3] P.-L. Lions, "Approximation de points fixes de contractions," *Comptes Rendus de l'Académie des Sciences de Paris A-B*, vol. 284, no. 21, pp. A1357–A1359, 1977 (French).
- [4] S. Reich, "Strong convergence theorems for resolvents of accretive operators in Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 75, no. 1, pp. 287–292, 1980.
- [5] S. Reich, "Some problems and results in fixed point theory," in *Topological Methods in Nonlinear Functional Analysis (Toronto, Ont., 1982)*, vol. 21 of *Contemporary Mathematics*, pp. 179–187, American Mathematical Society, Providence, RI, USA, 1983.
- [6] W. Takahashi and Y. Ueda, "On Reich's strong convergence theorems for resolvents of accretive operators," *Journal of Mathematical Analysis and Applications*, vol. 104, no. 2, pp. 546–553, 1984.
- [7] R. Wittmann, "Approximation of fixed points of nonexpansive mappings," *Archiv der Mathematik*, vol. 58, no. 5, pp. 486–491, 1992.
- [8] N. Shioji and W. Takahashi, "Strong convergence of approximated sequences for nonexpansive mappings in Banach spaces," *Proceedings of the American Mathematical Society*, vol. 125, no. 12, pp. 3641–3645, 1997.
- [9] H.-K. Xu, "Iterative algorithms for nonlinear operators," *Journal of the London Mathematical Society. Second Series*, vol. 66, no. 1, pp. 240–256, 2002.
- [10] H.-K. Xu, "An iterative approach to quadratic optimization," *Journal of Optimization Theory and Applications*, vol. 116, no. 3, pp. 659–678, 2003.
- [11] H.-K. Xu, "A strong convergence theorem for contraction semigroups in Banach spaces," *Bulletin of the Australian Mathematical Society*, vol. 72, no. 3, pp. 371–379, 2005.
- [12] T.-H. Kim and H.-K. Xu, "Strong convergence of modified Mann iterations," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 61, no. 1-2, pp. 51–60, 2005.
- [13] L.-G. Hu, "Strong convergence of a modified Halpern's iteration for nonexpansive mappings," *Fixed Point Theory and Applications*, vol. 2008, Article ID 649162, 9 pages, 2008.
- [14] M. R. Bridson and A. Haefliger, *Metric Spaces of Non-Positive Curvature*, vol. 319 of *Grundlehren der Mathematischen Wissenschaften*, Springer, Berlin, Germany, 1999.
- [15] W. A. Kirk, "Fixed point theorems in CAT(0) spaces and \mathbb{R} -trees," *Fixed Point Theory and Applications*, vol. 2004, no. 4, pp. 309–316, 2004.
- [16] K. S. Brown, *Buildings*, Springer, New York, NY, USA, 1989.
- [17] K. Goebel and S. Reich, *Uniform Convexity, Hyperbolic Geometry, and Nonexpansive Mappings*, vol. 83 of *Monographs and Textbooks in Pure and Applied Mathematics*, Marcel Dekker Inc., New York, NY, USA, 1984.
- [18] W. A. Kirk, "Geodesic geometry and fixed point theory," in *Seminar of Mathematical Analysis (Malaga/Seville, 2002/2003)*, vol. 64 of *Colección Abierta*, pp. 195–225, University of Seville, Secretary of Publications, Seville, Spain, 2003.
- [19] W. A. Kirk, "Geodesic geometry and fixed point theory. II," in *International Conference on Fixed Point Theory and Applications*, pp. 113–142, Yokohama Publishers, Yokohama, Japan, 2004.
- [20] S. Dhompongsa, A. Kaewkhao, and B. Panyanak, "Lim's theorems for multivalued mappings in CAT(0) spaces," *Journal of Mathematical Analysis and Applications*, vol. 312, no. 2, pp. 478–487, 2005.
- [21] P. Chaoha and A. Phon-on, "A note on fixed point sets in CAT(0) spaces," *Journal of Mathematical Analysis and Applications*, vol. 320, no. 2, pp. 983–987, 2006.
- [22] L. Leustean, "A quadratic rate of asymptotic regularity for CAT(0)-spaces," *Journal of Mathematical Analysis and Applications*, vol. 325, no. 1, pp. 386–399, 2007.
- [23] S. Dhompongsa and B. Panyanak, "On Δ -convergence theorems in CAT(0) spaces," *Computers & Mathematics with Applications*, vol. 56, no. 10, pp. 2572–2579, 2008.

- [24] N. Shahzad, "Fixed point results for multimaps in CAT(0) spaces," *Topology and Its Applications*, vol. 156, no. 5, pp. 997–1001, 2009.
- [25] R. Espinola and A. Fernandez-Leon, "CAT(k)-spaces, weak convergence and fixed points," *Journal of Mathematical Analysis and Applications*, vol. 353, no. 1, pp. 410–427, 2009.
- [26] N. Hussain and M. A. Khamsi, "On asymptotic pointwise contractions in metric spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 71, no. 10, pp. 4423–4429, 2009.
- [27] W. Laowang and B. Panyanak, "Strong and Δ convergence theorems for multivalued mappings in CAT(0) spaces," *Journal of Inequalities and Applications*, vol. 2009, Article ID 730132, 16 pages, 2009.
- [28] S. Saejung, "Halpern's iteration in CAT(0) spaces," *Fixed Point Theory and Applications*, vol. 2010, Article ID 471781, 13 pages, 2010.
- [29] A. R. Khan, M. A. Khamsi, and H. Fukhar-Ud-Din, "Strong convergence of a general iteration scheme in CAT(0) spaces," *Nonlinear Analysis: Theory, Methods and Applications*, vol. 74, no. 3, pp. 783–791, 2011.
- [30] S. H. Khan and M. Abbas, "Strong and Δ -convergence of some iterative schemes in CAT(0) spaces," *Computers and Mathematics with Applications*, vol. 61, no. 1, pp. 109–116, 2011.
- [31] A. Abkar and M. Eslamian, "Common fixed point results in CAT(0) spaces," *Nonlinear Analysis: Theory, Methods and Applications*, vol. 74, no. 5, pp. 1835–1840, 2011.
- [32] M. Bestvina, " \mathbb{R} -trees in topology, geometry, and group theory," in *Handbook of Geometric Topology*, pp. 55–91, North-Holland, Amsterdam, The Netherlands, 2002.
- [33] C. Semple and M. Steel, *Phylogenetics*, vol. 24 of *Oxford Lecture Series in Mathematics and Its Applications*, Oxford University Press, Oxford, UK, 2003.
- [34] R. Espinola and W. A. Kirk, "Fixed point theorems in \mathbb{R} -trees with applications to graph theory," *Topology and Its Applications*, vol. 153, no. 7, pp. 1046–1055, 2006.
- [35] W. A. Kirk, "Some recent results in metric fixed point theory," *Journal of Fixed Point Theory and Applications*, vol. 2, no. 2, pp. 195–207, 2007.
- [36] T. Suzuki, "Strong convergence theorems for infinite families of nonexpansive mappings in general Banach spaces," *Fixed Point Theory and Applications*, vol. 2005, no. 1, pp. 103–123, 2005.
- [37] Y. Song and R. Chen, "Strong convergence of an iterative method for non-expansive mappings," *Mathematische Nachrichten*, vol. 281, no. 8, pp. 1196–1204, 2008.
- [38] S. Wang, "A note on strong convergence of a modified Halpern's iteration for nonexpansive mappings," *Fixed Point Theory and Applications*, vol. 2010, Article ID 805326, 2 pages, 2010.