

ON A SUBGROUP OF THE AFFINE WEYL GROUP \tilde{C}_4

MUHAMMAD A. ALBAR

(Received 16 March 1999)

ABSTRACT. We study a subgroup of the affine Weyl group \tilde{C}_4 and show that this subgroup is a homomorphic image of the triangle group $\Delta(3, 4, 4)$.

Keywords and phrases. Group presentation, Reidemeister-Schreier process, Coxeter group, SQ -universality.

2000 Mathematics Subject Classification. Primary 20F05.

1. Introduction. In the algebraic structures of the Coxeter groups $\tilde{A}_{n-1}, B_n, D_n$, we observe the following. \tilde{A}_{n-1} is the subgroup of the wreath product $Z_2 S_n$ such that $\tilde{A}_{n-1} \cong Z^{n-1} \rtimes S_n$, where Z^{n-1} is the subgroup of Z^n consisting of all elements of exponent sum zero [2]; D_n is a subgroup of $B_n \cong Z_2 S_n$ such that $D_n \cong Z_2^{n-1} \rtimes S_n$ and Z_2^{n-1} is the subgroup of Z_2^n containing all elements of exponent sum zero [4]. We have the following natural question about $\tilde{C}_n \cong D_\infty^{n-1} \rtimes S_{n-1}$. What is the subgroup K of \tilde{C}_n , where $K \cong H \rtimes S_{n-1}$ and H is the subgroup of D_∞^{n-1} consisting of all elements of exponent sum zero [3]. In this paper we answer the question for $n = 4$ and find that the subgroup $H \rtimes S_3$ is a factor group of the triangle group $\Delta(3, 4, 4)$.

We begin by giving a presentation for the direct product of three copies of the infinite dihedral group

$$\begin{aligned}
 D_\infty^3 = \langle a_1, a_2, a_3, b_1, b_2, b_3 \mid & a_i^2 = b_i^2 = e, 1 \leq i \leq 3; \\
 & a_i a_j = a_j a_i, 1 \leq i < j \leq 3; \\
 & b_i b_j = b_j b_i, 1 \leq i < j \leq 3; \\
 & a_i b_j = b_j b_i \text{ if } i \neq j, 1 \leq i, j \leq 3 \rangle.
 \end{aligned}
 \tag{1.1}$$

A presentation for the symmetric group of degree 3 is

$$S_3 = \langle x_1, x_2 \mid x_1^2 = x_2^2 = (x_1 x_2)^3 = e \rangle.
 \tag{1.2}$$

In [3], it is shown that \tilde{C}_4 is the semi-direct product $\tilde{C}_4 \cong D_\infty^3 \rtimes S_3$ with the natural action

$$(a_1, a_2, a_3)^{x_1} = (a_2, a_1, a_3), (a_1, a_2, a_3)^{x_2} = (a_1, a_3, a_2),
 \tag{1.3}$$

$$(b_1, b_2, b_3)^{x_1} = (b_2, b_1, b_3), (b_1, b_2, b_3)^{x_2} = (b_1, b_3, b_2).
 \tag{1.4}$$

We consider the subgroup H of D_∞^3 containing all elements of exponent sum zero. H is a normal subgroup of D_∞^3 and $D_\infty^3/H \cong \langle a_1 \mid a_1^2 = e \rangle$. Using the Reidemeister-Schreier

process we find the following presentation for H :

$$\begin{aligned}
 H &= \langle \mathcal{Y}_1, \mathcal{Y}_2, \mathcal{Y}_3, \mathcal{Y}_4, \mathcal{Y}_5 \mid \mathcal{Y}_1^2 = \mathcal{Y}_2^2 = \mathcal{Y}_3^2 = \mathcal{Y}_4^2 = (\mathcal{Y}_1\mathcal{Y}_2)^2 = (\mathcal{Y}_2\mathcal{Y}_3)^2 = (\mathcal{Y}_3\mathcal{Y}_4)^2 \\
 &= (\mathcal{Y}_4\mathcal{Y}_5)^2 = (\mathcal{Y}_5\mathcal{Y}_1)^2 = (\mathcal{Y}_2\mathcal{Y}_4)^2 = (\mathcal{Y}_3\mathcal{Y}_5)^2 = (\mathcal{Y}_1\mathcal{Y}_4)^2 = e \rangle,
 \end{aligned}
 \tag{1.5}$$

where $\mathcal{Y}_1 = a_1b_3, \mathcal{Y}_2 = a_2a_1, \mathcal{Y}_3 = a_1a_3, \mathcal{Y}_4 = a_1b_1, \mathcal{Y}_5 = a_1b_2$. From the action of S_3 on D_∞^3 we easily compute the following action of S_3 on H :

$$(\mathcal{Y}_1, \mathcal{Y}_2, \mathcal{Y}_3, \mathcal{Y}_4, \mathcal{Y}_5)^{x_1} = (\mathcal{Y}_2\mathcal{Y}_1, \mathcal{Y}_2, \mathcal{Y}_2\mathcal{Y}_3, \mathcal{Y}_2\mathcal{Y}_5, \mathcal{Y}_2\mathcal{Y}_4),
 \tag{1.6}$$

$$(\mathcal{Y}_1, \mathcal{Y}_2, \mathcal{Y}_3, \mathcal{Y}_4, \mathcal{Y}_5)^{x_2} = (\mathcal{Y}_5, \mathcal{Y}_3, \mathcal{Y}_2, \mathcal{Y}_4, \mathcal{Y}_1).
 \tag{1.7}$$

2. The group $H \rtimes S_3$. We use the method of presentation of group extensions described in [1] to find a presentation for $H \rtimes S_3$ with the action computed in Section 1. A presentation for $H \rtimes S_3$ is

$$H \rtimes S_3 = \langle x_1, x_2, \mathcal{Y}_1, \mathcal{Y}_2, \mathcal{Y}_3, \mathcal{Y}_4, \mathcal{Y}_5 \mid RH, RS_3, H^{S_3} \rangle,
 \tag{2.1}$$

where RH are the relations of H , RS_3 are the relations of S_3 , the relations H^{S_3} are the action of S_3 on H . Lengthy computations using Tietze transformations give the following presentation for $H \rtimes S_3$,

$$H \rtimes S_3 = \langle a, b, c \mid a^2 = b^2 = c^2 = (ab)^3 = (bc)^4 = (ca)^4 = (bacac)^3 = e \rangle.
 \tag{2.2}$$

We observe that if $\Delta(3,4,4)$ is the hyperbolic triangle group generated by a, b , and c and N is the normal closure of $(bacac)^3$ in $\Delta(3,4,4)$, then $H \rtimes S_3$ is the factor group $(\Delta(3,4,4))/N$.

3. The triangle group $\Delta(3,4,4)$. The triangle group $\Delta(3,4,4)$ is given by the presentations

$$\Delta(3,4,4) = \langle a, b, c \mid a^2 = b^2 = c^2 = (ab)^3 = (bc)^4 = (ca)^4 = e \rangle.
 \tag{3.1}$$

It is one of the hyperbolic triangle groups. $\Delta(3,4,4)$ is SQ -universal [6]. We find the derived subgroup of $\Delta(3,4,4)$ and show that it is SQ -universal using a method different from that in [7]. We also compute the growth series (word growth in the sense of Milnor and Gromov) of $\Delta(3,4,4)$. Using the Reidemeister-Schreier process we find that $\Delta'(3,4,4)$ is

$$\Delta'(3,4,4) = \langle x, y, z \mid x^2 = y^4 = (xy)^3 = (yz^{-1})^2 = e \rangle.
 \tag{3.2}$$

We consider the map $\theta : \Delta(3,4,4) \rightarrow Z_2 = \langle v \mid v^2 = e \rangle$ defined by $\theta(x) = \theta(y) = \theta(z) = v$. It is easy to see that

$$\ker \theta = \langle a, b, c, d \mid (ab)^2 = c^3 = d^3 = (ab^{-1})^2 = (bd^{-1})^2 = e \rangle.
 \tag{3.3}$$

We define another map $\phi : \ker \theta \rightarrow Z_2 = \langle u \mid u^2 = e \rangle$ by $\phi(a) = \phi(b) = u$ and $\phi(c) = \phi(d) = e$. Then $\ker \phi$ has the presentation

$$\begin{aligned}
 \ker \phi &= \langle x_1, x_2, x_3, x_4, x_5, x_6 \mid x_3^2 = x_4^3 = x_5^3 = x_6^3 = (x_1x_2)^2 \\
 &= (x_1x_4)^3 = x_2x_6^{-1}x_3x_5^{-1} = x_3x_5^{-1}x_2x_6^{-1} = e \rangle.
 \end{aligned}
 \tag{3.4}$$

Letting $x_1 = x_5 = x_6 = e$ and $x_2 = x_3$ in $\ker \phi$ we get $\langle x_2, x_4 | x_2^2 = x_4^3 = e \rangle = Z_2 * Z_3$. Since the free product $Z_2 * Z_3$ is *SQU* [7], therefore $\ker \theta$ is *SQU*. But $\ker \theta$ is of finite index in $\Delta(3, 4, 4)$. Hence $\Delta(3, 4, 4)$ is *SQU* [7]. The growth series of $\Delta(3, 4, 4)$ is computed using exercise 26 in Section 1 of Chapter 4 in Bourbaki [5] as

$$y(t) = \frac{(1+t)(1+t+t^2)(1+t+t^2+t^3)}{1-t^2-2t^3-t^4+t^6}. \quad (3.5)$$

We observe that zeros of the denominator of $y(t)$ are not in the unit circle which implies that $\Delta(3, 4, 4)$ does not have a nilpotent subgroup of finite index. This is also known since $\Delta(3, 4, 4)$ is *SQU*.

REMARK 3.1. It is interesting to know what subgroup of \tilde{C}_n we get for $n > 4$. We did not find that yet.

ACKNOWLEDGEMENT. The author thanks King Fahd University of Petroleum and Minerals for support.

REFERENCES

- [1] M. A. Albar, *On presentation of group extensions*, Comm. Algebra **12** (1984), no. 23-24, 2967-2975. MR 86g:20040. Zbl 551.20017.
- [2] ———, *On the affine Weyl group of type \tilde{A}_{n-1}* , Int. J. Math. Math. Sci. **10** (1987), no. 1, 147-154. MR 88b:20051. Zbl 634.20014.
- [3] M. A. Albar and M. Al-Hamed, *The structure of the affine Weyl group \tilde{C}_n* , To appear in the Royal Irish Academy.
- [4] M. A. Albar and N. A. Al-Saleh, *On the affine Weyl group of type B_n* , Math. Japon. **35** (1990), no. 4, 599-602. MR 91d:20030. Zbl 790.20048.
- [5] N. Bourbaki, *Éléments de Mathématique. Fasc. XXXIV. Groupes et Algèbres de Lie. Chapitre IV: Groupes de Coxeter et Systèmes de Tits. Chapitre V: Groupes engendrés par des Réflexions. Chapitre VI: Systèmes de Racines*, Actualités Scientifiques et Industrielles, No. 1337, Hermann, Paris, 1968. MR 39#1590. Zbl 186.33001.
- [6] J. E. Humphreys, *Reflection Groups and Coxeter Groups*, Cambridge Studies in Advanced Mathematics, vol. 29, Cambridge University Press, Cambridge, 1990. MR 92h:20002. Zbl 725.20028.
- [7] P. M. Neumann, *The SQ-universality of some finitely presented groups*, J. Austral. Math. Soc. **16** (1973), 1-6, Collection of articles dedicated to the memory of Hanna Neumann, I. MR 48#11342. Zbl 267.20026.

MUHAMMAD A. ALBAR: DEPARTMENT OF MATHEMATICAL SCIENCES, KING FAHD UNIVERSITY OF PETROLEUM AND MINERALS, DHAHRAN 31261, SAUDI ARABIA