

Research Article

Asymptotic Stability Results for Nonlinear Fractional Difference Equations

Fulai Chen and Zhigang Liu

Department of Mathematics, Xiangnan University, Chenzhou 423000, China

Correspondence should be addressed to Fulai Chen, cflmath@163.com

Received 1 August 2011; Revised 27 December 2011; Accepted 2 January 2012

Academic Editor: Michela Redivo-Zaglia

Copyright © 2012 F. Chen and Z. Liu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We present some results for the asymptotic stability of solutions for nonlinear fractional difference equations involving Riemann-Liouville-like difference operator. The results are obtained by using Krasnoselskii's fixed point theorem and discrete Arzela-Ascoli's theorem. Three examples are also provided to illustrate our main results.

1. Introduction

In this paper we consider the asymptotic stability of solutions for nonlinear fractional difference equations:

$$\begin{aligned}\Delta^\alpha x(t) &= f(t + \alpha, x(t + \alpha)), \quad t \in N_0, \quad 0 < \alpha \leq 1, \\ \Delta^{\alpha-1} x(t)|_{t=0} &= x_0,\end{aligned}\tag{1.1}$$

where Δ^α is a Riemann-Liouville-like discrete fractional difference, $f : [0, +\infty) \times R \rightarrow R$ is continuous with respect to t and x , $N_a = \{a, a + 1, a + 2, \dots\}$.

Fractional differential equations have received increasing attention during recent years since these equations have been proved to be valuable tools in the modeling of many phenomena in various fields of science and engineering. Most of the present works were focused on fractional differential equations, see [1–12] and the references therein. However, very little progress has been made to develop the theory of the analogous fractional finite difference equation [13–19].

Due to the lack of geometry interpretation of the fractional derivatives, it is difficult to find a valid tool to analyze the stability of fractional difference equations. In the case that it

is difficult to employ Liapunov's direct method, fixed point theorems are usually considered in stability [20–25]. Motivated by this idea, in this paper, we discuss asymptotic stability of nonlinear fractional difference equations by using Krasnoselskii's fixed point theorem and discrete Arzela-Ascoli's theorem. Different from our previous work [18], in this paper, the sufficient conditions of attractivity are irrelevant to the initial value x_0 .

2. Preliminaries

In this section, we introduce preliminary facts of discrete fractional calculus. For more details, see [14].

Definition 2.1 (see [14]). Let $\nu > 0$. The ν -th fractional sum x is defined by

$$\Delta^{-\nu} f(t) = \frac{1}{\Gamma(\nu)} \sum_{s=a}^{t-\nu} (t-s-1)^{(\nu-1)} f(s), \quad (2.1)$$

where f is defined for $s = a \bmod(1)$ and $\Delta^{-\nu} f$ is defined for $t = (a + \nu) \bmod(1)$, and $t^{(\nu)} = \Gamma(t+1)/\Gamma(t-\nu+1)$. The fractional sum $\Delta^{-\nu}$ maps functions defined on N_a to functions defined on $N_{a+\nu}$.

Definition 2.2 (see [14]). Let $\mu > 0$ and $m-1 < \mu < m$, where m denotes a positive integer, $m = [\mu]$, $[\cdot]$ ceiling of number. Set $\nu = m - \mu$. The μ -th fractional difference is defined as

$$\Delta^{\mu} f(t) = \Delta^{m-\nu} f(t) = \Delta^m (\Delta^{-\nu} f(t)). \quad (2.2)$$

Theorem 2.3 (see [15]). *Let f be a real-value function defined on N_a and $\mu, \nu > 0$, then the following equalities hold:*

- (i) $\Delta^{-\nu} [\Delta^{-\mu} f(t)] = \Delta^{-(\mu+\nu)} f(t) = \Delta^{-\mu} [\Delta^{-\nu} f(t)];$
- (ii) $\Delta^{-\nu} \Delta f(t) = \Delta \Delta^{-\nu} f(t) - \frac{(t-a)^{(\nu-1)}}{\Gamma(\nu)} f(a).$

Lemma 2.4 (see [15]). *Let $\mu \neq 1$ and assume $\mu + \nu + 1$ is not a nonpositive integer, then*

$$\Delta^{-\nu} t^{(\mu)} = \frac{\Gamma(\mu+1)}{\Gamma(\mu+\nu+1)} t^{(\mu+\nu)}. \quad (2.3)$$

Lemma 2.5 (see [15]). *Assume that the following factorial functions are well defined:*

- (i) *If $0 < \alpha < 1$, then $t^{(\alpha\gamma)} \geq (t^{(\gamma)})^{\alpha};$*
- (ii) $t^{(\beta+\gamma)} = (t-\gamma)^{(\beta)} t^{(\gamma)}.$

Lemma 2.6 (see [13]). *Let $\mu > 0$ be noninteger, $m = [\mu]$, $[\cdot]$, $\nu = m - \mu$, thus one has*

$$\sum_{s=a+\nu}^{t-\mu} (t-s-1)^{(\mu-1)} = \frac{(t-a-\nu)^{(\mu)}}{\mu}. \quad (2.4)$$

Lemma 2.7. *The equivalent fractional Taylor's difference formula of (1.1) is*

$$x(t) = \frac{x_0}{\Gamma(\alpha)} t^{(\alpha-1)} + \frac{1}{\Gamma(\alpha)} \sum_{s=0}^{t-\alpha} (t-s-1)^{(\alpha-1)} f(s+\alpha, x(s+\alpha)), \quad t \in N_\alpha. \quad (2.5)$$

Proof. Apply the $\Delta^{-\alpha}$ operator to each side of the first formula of (1.1) to obtain

$$\Delta^{-\alpha} \Delta^\alpha x(t) = \Delta^{-\alpha} f(t+\alpha, x(t+\alpha)), \quad t \in N_\alpha. \quad (2.6)$$

Apply Theorem 2.3 to the left-hand side of (2.6) to obtain

$$\begin{aligned} \Delta^{-\alpha} \Delta^\alpha x(t) &= \Delta^{-\alpha} \Delta \Delta^{-(1-\alpha)} x(t) = \Delta \Delta^{-\alpha} \Delta^{-(1-\alpha)} x(t) - \frac{t^{(\alpha-1)}}{\Gamma(\alpha)} x(\alpha-1) \\ &= x(t) - \frac{x_0}{\Gamma(\alpha)} t^{(\alpha-1)}. \end{aligned} \quad (2.7)$$

So, applying Definition 2.1 to the right-hand side of (2.6), for $t \in N_\alpha$ we obtain (2.5). The recursive iteration to this Taylor's difference formula implies that (2.5) represents the unique solution of the IVP (1.1). This completes the proof. \square

Lemma 2.8 (see [4, (1.5.15)]). *The quotient expansion of two gamma functions at infinity is*

$$\frac{\Gamma(z+a)}{\Gamma(z+b)} = z^{a-b} \left[1 + O\left(\frac{1}{z}\right) \right], \quad (|\arg(z+a)| < \pi, |z| \rightarrow \infty). \quad (2.8)$$

Corollary 2.9. *One has*

$$t^{(-\beta)} > (t+\alpha)^{(-\beta)} \quad \text{for } \alpha, \beta, t > 0. \quad (2.9)$$

Proof. According to Lemma 2.8,

$$\begin{aligned} \frac{t^{(-\beta)}}{(t+\alpha)^{(-\beta)}} &= \frac{\Gamma(t+1)}{\Gamma(t+\beta+1)} \cdot \frac{\Gamma(t+\alpha+\beta+1)}{\Gamma(t+\alpha+1)} \\ &= \frac{\Gamma(t+1)}{\Gamma(t+\alpha+1)} \cdot \frac{\Gamma(t+\alpha+\beta+1)}{\Gamma(t+\beta+1)} \\ &= t^{-\alpha} \left[1 + O\left(\frac{1}{t}\right) \right] \cdot (t+\beta)^\alpha \left[1 + O\left(\frac{1}{t+\beta}\right) \right] \\ &= \left(1 + \frac{\beta}{t} \right)^\alpha \left[1 + O\left(\frac{1}{t}\right) \right] \left[1 + O\left(\frac{1}{t+\beta}\right) \right] \\ &> 1. \end{aligned} \quad (2.10)$$

Then, $t^{(-\beta)} > (t+\alpha)^{(-\beta)}$ for $\alpha, \beta, t > 0$. This completes the proof. \square

Definition 2.10. The solution $x = \varphi(t)$ of the IVP (1.1) is said to be

- (i) stable if for any $\varepsilon > 0$ and $t_0 \in \mathbb{R}^+$, there exists a $\delta = \delta(t_0, \varepsilon) > 0$ such that

$$|x(t, x_0, t_0) - \varphi(t)| < \varepsilon \quad (2.11)$$

for $|x_0 - \varphi(t_0)| \leq \delta(t_0, \varepsilon)$ and all $t \geq t_0$;

- (ii) attractive if there exists $\eta(t_0) > 0$ such that $\|x_0\| \leq \eta$ implies

$$\lim_{t \rightarrow \infty} x(t, x_0, t_0) = 0; \quad (2.12)$$

- (iii) asymptotically stable if it is stable and attractive.

The space $l_{n_0}^\infty$ is the set of real sequences defined on the set of positive integers where any individual sequence is bounded with respect to the usual supremum norm. It is well known that under the supremum norm $l_{n_0}^\infty$ is a Banach space [26].

Definition 2.11 (see [27]). A set Ω of sequences in $l_{n_0}^\infty$ is uniformly Cauchy (or equi-Cauchy), if for every $\varepsilon > 0$, there exists an integer N such that $|x(i) - x(j)| < \varepsilon$, whenever $i, j > N$ for any $x = \{x(n)\}$ in Ω .

Theorem 2.12 (see [27, (discrete Arzela-Ascoli's theorem)]). *A bounded, uniformly Cauchy subset Ω of $l_{n_0}^\infty$ is relatively compact.*

Theorem 2.13 (see [20, (Krasnoselskii's fixed point theorem)]). *Let S be a nonempty, closed, convex, and bounded subset of the Banach space X and let $A : X \rightarrow X$ and $B : S \rightarrow X$ be two operators such that*

- (a) A is a contraction with constant $L < 1$,
- (b) B is continuous, BS resides in a compact subset of X ,
- (c) $[x = Ax + By, y \in S] \Rightarrow x \in S$.

Then the operator equation $Ax + Bx = x$ has a solution in S .

3. Main Results

Let l_α^∞ be the set of all real sequences $x = \{x(t)\}_{t=\alpha}^\infty$ with norm $\|x\| = \sup_{t \in N_\alpha} |x(t)|$, then l_α^∞ is a Banach space.

Define the operator

$$\begin{aligned} Px(t) &= \frac{x_0}{\Gamma(\alpha)} t^{(\alpha-1)} + \frac{1}{\Gamma(\alpha)} \sum_{s=0}^{t-\alpha} (t-s-1)^{(\alpha-1)} f(s+\alpha, x(s+\alpha)), \\ Ax(t) &= \frac{x_0}{\Gamma(\alpha)} t^{(\alpha-1)}, \\ Bx(t) &= \frac{1}{\Gamma(\alpha)} \sum_{s=0}^{t-\alpha} (t-s-1)^{(\alpha-1)} f(s+\alpha, x(s+\alpha)), \quad t \in N_\alpha. \end{aligned} \quad (3.1)$$

Obviously, $Px = Ax + Bx$, the operator A is a contraction with the constant 0, which implies that condition (a) of Theorem 2.13 holds, and $x(t)$ is a solution of (1.1) if it is a fixed point of the operator P .

Lemma 3.1. Assume that the following condition is satisfied:

(H_1) there exist constants $\beta_1 \in (\alpha, 1)$ and $L_1 \geq 0$ such that

$$|f(t, x(t))| \leq L_1 t^{(-\beta_1)} \quad \text{for } t \in N_\alpha. \quad (3.2)$$

Then the operator B is continuous and BS_1 is a compact subset of R for $t \in N_{\alpha+n_1}$, where

$$S_1 = \left\{ x(t) : |x(t)| \leq t^{(-\gamma_1)} \text{ for } t \in N_{\alpha+n_1} \right\}, \quad (3.3)$$

$\gamma_1 = (-1/2)(\alpha - \beta_1)$, and $n_1 \in N$ satisfies that

$$\frac{|x_0|}{\Gamma(\alpha)} (\alpha + n_1 + \gamma_1)^{((1/2)(\alpha + \beta_1) - 1)} + \frac{L_1 \Gamma(1 - \beta_1)}{\Gamma(1 + \alpha - \beta_1)} (\alpha + n_1 + \gamma_1)^{(-\gamma_1)} \leq 1. \quad (3.4)$$

Proof. For $t \in N_\alpha$, apply Lemma 2.8 and $\gamma_1 > 0$,

$$t^{(-\gamma_1)} = \frac{\Gamma(t+1)}{\Gamma(t+\gamma_1+1)} = t^{-\gamma_1} \left[1 + O\left(\frac{1}{t}\right) \right], \quad (3.5)$$

and we have that $t^{(-\gamma_1)} \rightarrow 0$ as $t \rightarrow \infty$, then there exists a $n_1 \in N$ such that inequality (3.4) holds, which implies that the set S_1 exists.

We firstly show that B maps S_1 in S_1 .

It is easy to know that S_1 is a closed, bounded, and convex subset of R .

Apply condition (H_1) , Lemma 2.5, Corollary 2.9 and (3.4), for $t \in N_{\alpha+n_1}$, we have

$$\begin{aligned} |Bx(t)| &\leq \frac{1}{\Gamma(\alpha)} \sum_{s=0}^{t-\alpha} (t-s-1)^{(\alpha-1)} |f(s+\alpha, x(s+\alpha))| \\ &\leq \frac{1}{\Gamma(\alpha)} \sum_{s=0}^{t-\alpha} (t-s-1)^{(\alpha-1)} L_1 (s+\alpha)^{(-\beta_1)} \\ &= L_1 \Delta^{-\alpha} (t+\alpha)^{(-\beta_1)} \\ &= \frac{L_1 \Gamma(1-\beta_1)}{\Gamma(1+\alpha-\beta_1)} (t+\alpha)^{(\alpha-\beta_1)} \\ &< \frac{L_1 \Gamma(1-\beta_1)}{\Gamma(1+\alpha-\beta_1)} t^{(\alpha-\beta_1)} \\ &= \frac{L_1 \Gamma(1-\beta_1)}{\Gamma(1+\alpha-\beta_1)} (t+\gamma_1)^{(-\gamma_1)} t^{(-\gamma_1)} \end{aligned}$$

$$\begin{aligned}
&\leq \frac{L_1\Gamma(1-\beta_1)}{\Gamma(1+\alpha-\beta_1)}(\alpha+n_1+\gamma_1)^{(-\gamma_1)}t^{(-\gamma_1)} \\
&\leq t^{(-\gamma_1)},
\end{aligned} \tag{3.6}$$

which implies that $BS_1 \subset S_1$ for $t \in N_{\alpha+n_1}$.

Next, we show that B is continuous on S_1 .

Let $\varepsilon > 0$ be given then there exist $T_1 \in \mathbb{N}$ and $T_1 \geq n_1$ such that $t \in N_{\alpha+T_1}$ implies that

$$\frac{L_1\Gamma(1-\beta_1)}{\Gamma(1+\alpha-\beta_1)}t^{(\alpha-\beta_1)} < \frac{\varepsilon}{2}. \tag{3.7}$$

Let $\{x_n\}$ be a sequence such that $x_n \rightarrow x$. For $t \in \{\alpha+n_1, \alpha+n_1+1, \dots, \alpha+T_1-1\}$, applying the continuity of f and Lemma 2.6, we have

$$\begin{aligned}
|Bx_n(t) - Bx(t)| &\leq \frac{1}{\Gamma(\alpha)} \sum_{s=0}^{t-\alpha} (t-s-1)^{(\alpha-1)} |f(s+\alpha, x_n(s+\alpha)) - f(s+\alpha, x(s+\alpha))| \\
&\leq \max_{s \in \{0, 1, \dots, T_1-1\}} |f(s+\alpha, x_n(s+\alpha)) - f(s+\alpha, x(s+\alpha))| \\
&\quad \times \frac{1}{\Gamma(\alpha)} \sum_{s=0}^{t-\alpha} (t-s-1)^{(\alpha-1)} \\
&= \frac{t^{(\alpha)}}{\Gamma(\alpha+1)} \max_{s \in \{0, 1, \dots, T_1-1\}} |f(s+\alpha, x_n(s+\alpha)) - f(s+\alpha, x(s+\alpha))| \\
&\leq \frac{(\alpha+T_1-1)^{(\alpha)}}{\Gamma(\alpha+1)} \max_{s \in \{0, 1, \dots, T_1-1\}} |f(s+\alpha, x_n(s+\alpha)) - f(s+\alpha, x(s+\alpha))| \\
&\rightarrow 0 \quad \text{as } n \rightarrow \infty.
\end{aligned} \tag{3.8}$$

For $t \in N_{\alpha+T_1}$,

$$\begin{aligned}
|Bx_n(t) - Bx(t)| &\leq \frac{1}{\Gamma(\alpha)} \sum_{s=0}^{t-\alpha} (t-s-1)^{(\alpha-1)} [|f(s+\alpha, x_n(s+\alpha))| + |f(s+\alpha, x(s+\alpha))|] \\
&\leq \frac{2L_1}{\Gamma(\alpha)} \sum_{s=0}^{t-\alpha} (t-s-1)^{(\alpha-1)} (s+\alpha)^{(-\beta_1)} \\
&= 2L_1 \Delta^{-\alpha} (t+\alpha)^{(-\beta_1)} \\
&= \frac{2L_1\Gamma(1-\beta_1)}{\Gamma(1+\alpha-\beta_1)} (t+\alpha)^{(\alpha-\beta_1)} \\
&< \frac{2L_1\Gamma(1-\beta_1)}{\Gamma(1+\alpha-\beta_1)} t^{(\alpha-\beta_1)} \\
&< \varepsilon.
\end{aligned} \tag{3.9}$$

Thus, for all $t \in N_{\alpha+n_1}$, we have

$$|Bx_n(t) - Bx(t)| \longrightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (3.10)$$

which implies that B is continuous.

Lastly, we show that BS_1 is relatively compact.

Let $t_1, t_2 \in N_{\alpha+T_1}$ and $t_2 > t_1$, thus we have

$$\begin{aligned} |Bx(t_2) - Bx(t_1)| &= \left| \frac{1}{\Gamma(\alpha)} \sum_{s=0}^{t_2-\alpha} (t_2 - s - 1)^{(\alpha-1)} f(s + \alpha, x(s + \alpha)) \right. \\ &\quad \left. - \frac{1}{\Gamma(\alpha)} \sum_{s=0}^{t_1-\alpha} (t_1 - s - 1)^{(\alpha-1)} f(s + \alpha, x(s + \alpha)) \right| \\ &\leq \frac{1}{\Gamma(\alpha)} \sum_{s=0}^{t_2-\alpha} (t_2 - s - 1)^{(\alpha-1)} |f(s + \alpha, x(s + \alpha))| \\ &\quad + \frac{1}{\Gamma(\alpha)} \sum_{s=0}^{t_1-\alpha} (t_1 - s - 1)^{(\alpha-1)} |f(s + \alpha, x(s + \alpha))| \\ &\leq \frac{L_1 \Gamma(1 - \beta_1)}{\Gamma(1 + \alpha - \beta_1)} (t_2 + \alpha)^{(\alpha-\beta_1)} + \frac{L_1 \Gamma(1 - \beta_1)}{\Gamma(1 + \alpha - \beta_1)} (t_1 + \alpha)^{(\alpha-\beta_1)} \\ &< \frac{L_1 \Gamma(1 - \beta_1)}{\Gamma(1 + \alpha - \beta_1)} t_2^{(\alpha-\beta_1)} + \frac{L_1 \Gamma(1 - \beta_1)}{\Gamma(1 + \alpha - \beta_1)} t_1^{(\alpha-\beta_1)} \\ &< \varepsilon. \end{aligned} \quad (3.11)$$

Thus, $\{Bx : x \in S_1\}$ is a bounded and uniformly Cauchy subset by Definition 2.11, and BS_1 is relatively compact by means of Theorem 2.12. This completes the proof. \square

Lemma 3.2. Assume that condition (H_1) holds, then a solution of (1.1) is in S_1 for $t \in N_{\alpha+n_1}$.

Proof. Notice if that $x(t)$ is a fixed point of P , then it is a solution of (1.1). To prove this, it remains to show that, for fixed $y \in S_1$, $x = Ax + By \Rightarrow x \in S_1$ holds.

If $x = Ax + By$, applying condition (H_1) and (3.4), for $t \in N_{\alpha+n_1}$, we have

$$\begin{aligned} |x(t)| &\leq |Ax(t)| + |By(t)| \\ &\leq \frac{|x_0|}{\Gamma(\alpha)} t^{(\alpha-1)} + \frac{1}{\Gamma(\alpha)} \sum_{s=0}^{t-\alpha} (t - s - 1)^{(\alpha-1)} |f(s + \alpha, y(s + \alpha))| \\ &\leq \frac{|x_0|}{\Gamma(\alpha)} t^{(\alpha-1)} + \frac{L_1 \Gamma(1 - \beta_1)}{\Gamma(1 + \alpha - \beta_1)} (t + \alpha)^{(\alpha-\beta_1)} \\ &< \frac{|x_0|}{\Gamma(\alpha)} t^{(\alpha-1)} + \frac{L_1 \Gamma(1 - \beta_1)}{\Gamma(1 + \alpha - \beta_1)} t^{(\alpha-\beta_1)} \end{aligned}$$

$$\begin{aligned}
&= \left[\frac{|x_0|}{\Gamma(\alpha)} (t + \gamma_1)^{((1/2)(\alpha+\beta_1)-1)} + \frac{L_1 \Gamma(1 - \beta_1)}{\Gamma(1 + \alpha - \beta_1)} (t + \gamma_1)^{(-\gamma_1)} \right] t^{(-\gamma_1)} \\
&\leq \left[\frac{|x_0|}{\Gamma(\alpha)} (\alpha + n_1 + \gamma_1)^{((1/2)(\alpha+\beta_1)-1)} + \frac{L_1 \Gamma(1 - \beta_1)}{\Gamma(1 + \alpha - \beta_1)} (\alpha + n_1 + \gamma_1)^{(-\gamma_1)} \right] t^{(-\gamma_1)} \\
&\leq t^{(-\gamma_1)}.
\end{aligned} \tag{3.12}$$

Thus, $x(t) \in S_1$ for $t \in N_{\alpha+n_1}$. According to Theorem 2.13 and Lemma 3.1, there exists a $x \in S_1$ such that $x = Ax+Bx$, that is, P has a fixed point in S_1 which is a solution of (1.1) for $t \in N_{\alpha+n_1}$. This completes the proof. \square

Theorem 3.3. Assume that condition (H_1) holds, then the solutions of (1.1) is attractive.

Proof. By Lemma 3.2, the solutions of (1.1) exist and are in S_1 . All functions $x(t)$ in S_1 tend to 0 as $t \rightarrow \infty$. Then the solutions of (1.1) tend to zero as $t \rightarrow \infty$. This completes the proof. \square

Theorem 3.4. Assume that the following condition is satisfied:

(H_2) there exist constants $\beta_2 \in (\alpha, 1)$ and $L_2 \geq 0$ such that

$$|f(t, x(t)) - f(t, y(t))| \leq L_2 t^{(-\beta_2)} \|x - y\| \quad \text{for } t \in N_\alpha. \tag{3.13}$$

Then the solutions of (1.1) are stable provided that

$$c := \frac{L_2 \Gamma(1 + \alpha) \Gamma(1 - \beta_2)}{\Gamma(1 + \alpha - \beta_2) \Gamma(1 + \beta_2)} < 1. \tag{3.14}$$

Proof. Let $x(t)$ be a solution of (1.1), and let $\tilde{x}(t)$ be a solution of (1.1) satisfying the initial value condition $\tilde{x}(0) = \tilde{x}_0$. For $t \in N_\alpha$, applying condition (H_2) , we have

$$\begin{aligned}
|x(t) - \tilde{x}(t)| &\leq \frac{t^{(\alpha-1)}}{\Gamma(\alpha)} |x_0 - \tilde{x}_0| + \frac{1}{\Gamma(\alpha)} \sum_{s=0}^{t-\alpha} (t-s-1)^{(\alpha-1)} \\
&\quad \times |f(s+\alpha, x(s+\alpha)) - f(s+\alpha, \tilde{x}(s+\alpha))| \\
&\leq \frac{t^{(\alpha-1)}}{\Gamma(\alpha)} |x_0 - \tilde{x}_0| + \frac{L_2}{\Gamma(\alpha)} \sum_{s=0}^{t-\alpha} (t-s-1)^{(\alpha-1)} (s+\alpha)^{(-\beta_2)} \|x - \tilde{x}\| \\
&= \frac{t^{(\alpha-1)}}{\Gamma(\alpha)} |x_0 - \tilde{x}_0| + L_2 \Delta^{-\alpha} (t+\alpha)^{(-\beta_2)} \|x - \tilde{x}\| \\
&= \frac{t^{(\alpha-1)}}{\Gamma(\alpha)} |x_0 - \tilde{x}_0| + \frac{L_2 \Gamma(1 - \beta_2)}{\Gamma(1 + \alpha - \beta_2)} (t+\alpha)^{(\alpha-\beta_2)} \|x - \tilde{x}\| \\
&\leq \frac{\alpha^{(\alpha-1)}}{\Gamma(\alpha)} |x_0 - \tilde{x}_0| + \frac{L_2 \Gamma(1 - \beta_2)}{\Gamma(1 + \alpha - \beta_2)} \alpha^{(\alpha-\beta_2)} \|x - \tilde{x}\|
\end{aligned}$$

$$\begin{aligned}
&= \alpha|x_0 - \tilde{x}_0| + \frac{L_2\Gamma(1+\alpha)\Gamma(1-\beta_2)}{\Gamma(1+\alpha-\beta_2)\Gamma(1+\beta_2)}\|x - \tilde{x}\| \\
&= \alpha|x_0 - \tilde{x}_0| + c\|x - \tilde{x}\|,
\end{aligned} \tag{3.15}$$

which implies that

$$\|x - \tilde{x}\| \leq \frac{\alpha}{1-c}|x_0 - \tilde{x}_0|. \tag{3.16}$$

For any given $\varepsilon > 0$, let $\delta = ((1-c)/\alpha)\varepsilon$, $|x_0 - \tilde{x}_0| < \delta$ follows that $\|x - \tilde{x}\| < \varepsilon$, which yields that the solutions of (1.1) are stable. This completes the proof. \square

Theorem 3.5. Assume that conditions (H_1) and (H_2) hold, then the solutions of (1.1) are asymptotically stable provided that (3.14) holds.

Theorem 3.5 is the simple consequence of Theorems 3.3 and 3.4.

Theorem 3.6. Assume that the following condition is satisfied:

(H_3) there exist constants $\beta_3 \in (\alpha, (1/2)(1+\alpha))$, $\gamma_2 = (1/2)(1-\alpha)$, and $L_3 \geq 0$ such that

$$|f(t, x(t))| \leq L_3(t + \gamma_2)^{(-\beta_3)}|x(t)| \quad \text{for } t \in N_\alpha. \tag{3.17}$$

Then the solutions of (1.1) is attractive.

Proof. Set

$$S_2 = \left\{ x(t) : |x(t)| \leq t^{(-\gamma_2)} \text{ for } t \in N_{\alpha+n_2} \right\}, \tag{3.18}$$

where $n_2 \in N$ satisfies that

$$\frac{|x_0|}{\Gamma(\alpha)}(\alpha + n_2 + \gamma_2)^{(-\gamma_2)} + \frac{L_3\Gamma(1-\beta_3-\gamma_2)}{\Gamma(1+\alpha-\beta_3-\gamma_2)}(\alpha + n_2 + \gamma_2)^{(\alpha-\beta_3)} \leq 1. \tag{3.19}$$

We first prove condition (c) of Theorem 2.13, that is, for fixed $y \in S_2$ and for all $x \in R$, $x = Ax + By \Rightarrow x \in S_2$ holds.

If $x = Ax + By$, applying condition (H_3) and (3.19), for $t \in N_{\alpha+n_2}$, we have

$$\begin{aligned}
|x(t)| &\leq |Ax(t)| + |By(t)| \\
&\leq \frac{|x_0|}{\Gamma(\alpha)} t^{(\alpha-1)} + \frac{1}{\Gamma(\alpha)} \sum_{s=0}^{t-\alpha} (t-s-1)^{(\alpha-1)} |f(s+\alpha, y(s+\alpha))| \\
&\leq \frac{|x_0|}{\Gamma(\alpha)} t^{(\alpha-1)} + \frac{1}{\Gamma(\alpha)} \sum_{s=0}^{t-\alpha} (t-s-1)^{(\alpha-1)} L_3 (s+\alpha+\gamma_2)^{(-\beta_3)} |y(s+\alpha)| \\
&\leq \frac{|x_0|}{\Gamma(\alpha)} t^{(\alpha-1)} + \frac{L_3}{\Gamma(\alpha)} \sum_{s=0}^{t-\alpha} (t-s-1)^{(\alpha-1)} (s+\alpha+\gamma_2)^{(-\beta_3)} (s+\alpha)^{(-\gamma_2)} \\
&\leq \frac{|x_0|}{\Gamma(\alpha)} t^{(\alpha-1)} + \frac{L_3}{\Gamma(\alpha)} \sum_{s=0}^{t-\alpha} (t-s-1)^{(\alpha-1)} (s+\alpha)^{(-\beta_3-\gamma_2)} \\
&\leq \frac{|x_0|}{\Gamma(\alpha)} t^{(\alpha-1)} + \frac{L_3 \Gamma(1-\beta_3-\gamma_2)}{\Gamma(1+\alpha-\beta_3-\gamma_2)} (t+\alpha)^{(\alpha-\beta_3-\gamma_2)} \\
&< \frac{|x_0|}{\Gamma(\alpha)} t^{(\alpha-1)} + \frac{L_3 \Gamma(1-\beta_3-\gamma_2)}{\Gamma(1+\alpha-\beta_3-\gamma_2)} t^{(\alpha-\beta_3-\gamma_2)} \\
&= \left[\frac{|x_0|}{\Gamma(\alpha)} (t+\gamma_2)^{(-\gamma_2)} + \frac{L_3 \Gamma(1-\beta_3-\gamma_2)}{\Gamma(1+\alpha-\beta_3-\gamma_2)} (t+\gamma_2)^{(\alpha-\beta_3)} \right] t^{(-\gamma_2)} \\
&\leq \left[\frac{|x_0|}{\Gamma(\alpha)} (\alpha+n_2+\gamma_2)^{(-\gamma_2)} + \frac{L_3 \Gamma(1-\beta_3-\gamma_2)}{\Gamma(1+\alpha-\beta_3-\gamma_2)} (\alpha+n_2+\gamma_2)^{(\alpha-\beta_3)} \right] t^{(-\gamma_2)} \\
&\leq t^{(-\gamma_2)}.
\end{aligned} \tag{3.20}$$

Thus, condition (c) of Theorem 2.13 holds.

The proof of condition (b) of Theorem 2.13 is similar to that of Lemma 3.1, and we omit it. Therefore, P has a fixed point in S_2 by using Theorem 2.13, that is, the IVP (1.1) has a solution in S_2 . Moreover, all functions in S_2 tend to 0 as $t \rightarrow \infty$, then the solution of (1.1) tends to zero as $t \rightarrow \infty$, which shows that the zero solution of (1.1) is attractive. This completes the proof. \square

Theorem 3.7. Assume that conditions (H_2) and (H_3) hold, then the solutions of (1.1) are asymptotically stable provided that (3.14) holds.

Theorem 3.8. Assume that the following condition is satisfied:

(H_4) there exist constants $\eta \in (0, 1)$, $\beta_4 \in (\alpha, (2+\alpha\eta)/(2+\eta))$, and $L_4 \geq 0$ such that

$$|f(t, x(t))| \leq L_4 (t+1)^{(-\beta_4)} |x(t)|^\eta \quad \text{for } t \in N_\alpha. \tag{3.21}$$

Then the solutions of (1.1) is attractive.

Proof. Set

$$S_3 = \left\{ x(t) : |x(t)| \leq t^{(-\gamma_3)} \text{ for } t \in N_{\alpha+n_3} \right\}, \tag{3.22}$$

where $\gamma_3 = (1/2)(\beta_4 - \alpha)$, and $n_3 \in N$ satisfies that

$$\frac{|x_0|}{\Gamma(\alpha)} (\alpha + n_3 + \gamma_3)^{(\alpha-1+\gamma_3)} + \frac{L_4 \Gamma(1 - \beta_4 - \gamma_3 \eta)}{\Gamma(1 + \alpha - \beta_4 - \gamma_3 \eta)} (\alpha + n_3 + \gamma_3)^{-\gamma_3} \leq 1. \quad (3.23)$$

Here we only prove that condition (c) of Theorem 2.13 holds, and the remaining part of the proof is similar to that of Theorem 3.6.

Since $\eta \in (0, 1)$, $\beta_4 \in (\alpha, (2 + \alpha\eta)/(2 + \eta))$, and $\gamma_3 = (1/2)(\beta_4 - \alpha)$, then $\gamma_3, \gamma_3\eta, \alpha + \gamma_3 \in (0, 1)$, $\beta_4 + \gamma_3\eta \in (\alpha, 1)$.

If $x = Ax + By$, applying condition (H_4) , Lemma 2.5 and (3.23), for $t \in N_{\alpha+n_3}$, we have

$$\begin{aligned} |x(t)| &\leq |Ax(t)| + |By(t)| \leq \frac{|x_0|}{\Gamma(\alpha)} t^{(\alpha-1)} + \frac{1}{\Gamma(\alpha)} \sum_{s=0}^{t-\alpha} (t-s-1)^{(\alpha-1)} |f(s+\alpha, y(s+\alpha))| \\ &\leq \frac{|x_0|}{\Gamma(\alpha)} t^{(\alpha-1)} + \frac{1}{\Gamma(\alpha)} \sum_{s=0}^{t-\alpha} (t-s-1)^{(\alpha-1)} L_4 (s+\alpha+1)^{(-\beta_4)} |y(s+\alpha)|^\eta \\ &\leq \frac{|x_0|}{\Gamma(\alpha)} t^{(\alpha-1)} + \frac{L_4}{\Gamma(\alpha)} \sum_{s=0}^{t-\alpha} (t-s-1)^{(\alpha-1)} (s+\alpha+\gamma_3\eta)^{(-\beta_4)} \left[(s+\alpha)^{(-\gamma_3)} \right]^\eta \\ &\leq \frac{|x_0|}{\Gamma(\alpha)} t^{(\alpha-1)} + \frac{L_4}{\Gamma(\alpha)} \sum_{s=0}^{t-\alpha} (t-s-1)^{(\alpha-1)} (s+\alpha+\gamma_3\eta)^{(-\beta_4)} (s+\alpha)^{(-\gamma_3\eta)} \\ &= \frac{|x_0|}{\Gamma(\alpha)} t^{(\alpha-1)} + \frac{L_4}{\Gamma(\alpha)} \sum_{s=0}^{t-\alpha} (t-s-1)^{(\alpha-1)} (s+\alpha)^{(-\beta_4-\gamma_3\eta)} \\ &\leq \frac{|x_0|}{\Gamma(\alpha)} t^{(\alpha-1)} + \frac{L_4 \Gamma(1 - \beta_4 - \gamma_3 \eta)}{\Gamma(1 + \alpha - \beta_4 - \gamma_3 \eta)} (t + \alpha)^{(\alpha - \beta_4 - \gamma_3 \eta)} \\ &< \frac{|x_0|}{\Gamma(\alpha)} t^{(\alpha-1)} + \frac{L_4 \Gamma(1 - \beta_4 - \gamma_3 \eta)}{\Gamma(1 + \alpha - \beta_4 - \gamma_3 \eta)} t^{(\alpha - \beta_4 - \gamma_3 \eta)} \\ &\leq \frac{|x_0|}{\Gamma(\alpha)} t^{(\alpha-1)} + \frac{L_4 \Gamma(1 - \beta_4 - \gamma_3 \eta)}{\Gamma(1 + \alpha - \beta_4 - \gamma_3 \eta)} t^{(\alpha - \beta_4)} \\ &= \left[\frac{|x_0|}{\Gamma(\alpha)} (t + \gamma_3)^{(\alpha-1+\gamma_3)} + \frac{L_4 \Gamma(1 - \beta_4 - \gamma_3 \eta)}{\Gamma(1 + \alpha - \beta_4 - \gamma_3 \eta)} (t + \gamma_3)^{(-\gamma_3)} \right] t^{(-\gamma_3)} \\ &\leq \left[\frac{|x_0|}{\Gamma(\alpha)} (\alpha + n_3 + \gamma_3)^{(\alpha-1+\gamma_3)} \right. \\ &\quad \left. + \frac{L_4 \Gamma(1 - \beta_4 - \gamma_3 \eta)}{\Gamma(1 + \alpha - \beta_4 - \gamma_3 \eta)} (\alpha + n_3 + \gamma_3)^{(-\gamma_3)} \right] t^{(-\gamma_3)} \\ &\leq t^{(-\gamma_3)}. \end{aligned} \quad (3.24)$$

Thus, condition (c) of Theorem 2.13 holds. This completes the proof. \square

4. Examples

Example 4.1. Consider

$$\begin{aligned} \Delta^{0.5}x(t) &= 0.2(t+0.5)^{(-0.75)} \sin(x(t+0.5)), \quad t \in N_0, \\ \Delta^{-0.5}x(t)|_{t=0} &= x_0, \end{aligned} \quad (4.1)$$

where $f(t, x(t)) = 0.2t^{(-0.75)} \sin(x(t))$, $t \in N_{0.5}$.

Since

$$|f(t, x(t))| = \left| 0.2t^{(-0.75)} \sin(x(t)) \right| \leq 0.2t^{(-0.75)}, \quad (4.2)$$

this implies that condition (H_1) holds.

In addition,

$$|f(t, x(t)) - f(t, y(t))| \leq 0.2t^{(-0.75)} \|x - y\|. \quad (4.3)$$

Thus, condition (H_2) is satisfied.

Moreover, from $L_2 = 0.2$, $\alpha = 0.5$, and $\beta_2 = 0.75$, we have

$$c = \frac{L_2 \Gamma(1 + \alpha) \Gamma(1 - \beta_2)}{\Gamma(1 + \alpha - \beta_2) \Gamma(1 + \beta_2)} = \frac{0.2 \Gamma(1.5) \Gamma(0.25)}{\Gamma(1.25) \Gamma(1.75)} \approx 0.7716 < 1, \quad (4.4)$$

which implies that inequality (3.14) holds.

Thus the solutions of (4.1) are asymptotically stable by Theorem 3.5.

Example 4.2. Consider

$$\begin{aligned} \Delta^{0.5}x(t) &= 0.2(t+1.5)^{(-0.6)} x(t+0.5), \quad t \in N_0, \\ \Delta^{-0.5}x(t)|_{t=0} &= x_0, \end{aligned} \quad (4.5)$$

where $f(t, x(t)) = 0.2(t+1)^{(-0.6)} x(t)$, $t \in N_{0.5}$.

Since $\beta_3 = 0.6$, $\alpha = 0.5$, we have that $\beta_3 \in (\alpha, (1/2)(1 + \alpha))$, $\gamma_2 = 0.25$ and

$$|f(t, x(t))| = \left| 0.2(t+1)^{(-0.6)} x(t) \right| \leq 0.2(t+0.25)^{(-0.6)} |x(t)|, \quad (4.6)$$

which implies that condition (H_3) is satisfied.

Meanwhile,

$$|f(t, x(t)) - f(t, y(t))| \leq 0.2(t+1)^{(-0.6)} \|x - y\| \leq 0.2t^{(-0.6)} \|x - y\|, \quad (4.7)$$

which implies that condition (H_2) is satisfied.

From $L_2 = 0.2$, $\alpha = 0.5$, and $\beta_2 = 0.6$, we have

$$c = \frac{L_2 \Gamma(1 + \alpha) \Gamma(1 - \beta_2)}{\Gamma(1 + \alpha - \beta_2) \Gamma(1 + \beta_2)} = \frac{0.2 \Gamma(1.5) \Gamma(0.4)}{\Gamma(0.9) \Gamma(1.6)} \approx 0.4120 < 1, \quad (4.8)$$

which implies that inequality (3.14) holds.

Thus the solutions of (4.5) are asymptotically stable by Theorem 3.7.

Example 4.3. Consider

$$\begin{aligned} \Delta^{0.5} x(t) &= (t + 1.5)^{(-0.6)} x^{1/3}(t + 0.5), \quad t \in N_0, \\ \Delta^{-0.5} x(t)|_{t=0} &= x_0, \end{aligned} \quad (4.9)$$

where $f(t, x(t)) = (t + 1)^{(-0.6)} x^{1/3}(t)$, $t \in N_{0.5}$.

Since $\alpha = 0.5$, $\beta_4 = 0.6$, $\eta = 1/3$, we have that $\eta \in (0, 1)$, $\beta_4 \in (\alpha, (2 + \alpha\eta)/(2 + \eta))$ and

$$|f(t, x(t))| \leq (t + 1)^{(-0.6)} |x(t)|^{1/3}, \quad (4.10)$$

then condition (H_4) is satisfied.

The solutions of (4.9) are attractive by Theorem 3.8.

Acknowledgments

This research was supported by the NSF of Hunan Province (10JJ6007, 2011FJ3013), the Scientific Research Foundation of Hunan Provincial Education Department, and the Construct Program of the Key Discipline in Hunan Province.

References

- [1] R. P. Agarwal, V. Lakshmikantham, and J. J. Nieto, "On the concept of solution for fractional differential equations with uncertainty," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 72, no. 6, pp. 2859–2862, 2010.
- [2] M. Benchohra, J. Henderson, S. K. Ntouyas, and A. Ouahab, "Existence results for fractional order functional differential equations with infinite delay," *Journal of Mathematical Analysis and Applications*, vol. 338, no. 2, pp. 1340–1350, 2008.
- [3] F. Chen, A. Chen, and X. Wang, "On the solutions for impulsive fractional functional differential equations," *Differential Equations and Dynamical Systems*, vol. 17, no. 4, pp. 379–391, 2009.
- [4] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, North-Holland Mathematics Studies, Elsevier Science B.V., Amsterdam, the Netherlands, 2006.
- [5] V. Lakshmikantham, "Theory of fractional functional differential equations," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 69, no. 10, pp. 3337–3343, 2008.
- [6] V. Lakshmikantham, S. Leela, and J. V. Devi, *Theory of Fractional Dynamic Systems*, Cambridge Scientific Publishers, Cambridge, UK, 2009.
- [7] K. S. Miller and B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*, John Wiley & Sons, New York, NY, USA, 1993.
- [8] J. J. Nieto, "Maximum principles for fractional differential equations derived from Mittag-Leffler functions," *Applied Mathematics Letters*, vol. 23, no. 10, pp. 1248–1251, 2010.

- [9] I. Podlubny, *Fractional Differential Equations*, Academic Press, San Diego, Calif, USA, 1999.
- [10] J. Wang and Y. Zhou, "A class of fractional evolution equations and optimal controls," *Nonlinear Analysis. Real World Applications*, vol. 12, no. 1, pp. 262–272, 2011.
- [11] Y. Zhou, F. Jiao, and J. Li, "Existence and uniqueness for P -type fractional neutral differential equations," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 71, no. 7-8, pp. 2724–2733, 2009.
- [12] Y. Zhou, F. Jiao, and J. Li, "Existence and uniqueness for fractional neutral differential equations with infinite delay," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 71, no. 7-8, pp. 3249–3256, 2009.
- [13] G. A. Anastassiou, "Discrete fractional calculus and inequalities," *Classical Analysis and ODEs*. In press.
- [14] F. M. Atici and P. W. Eloe, "Initial value problems in discrete fractional calculus," *Proceedings of the American Mathematical Society*, vol. 137, no. 3, pp. 981–989, 2009.
- [15] F. M. Atici and P. W. Eloe, "A transform method in discrete fractional calculus," *International Journal of Difference Equations*, vol. 2, no. 2, pp. 165–176, 2007.
- [16] F. M. Atici and P. W. Eloe, "Discrete fractional calculus with the nabla operator," *Electronic Journal of Qualitative Theory of Differential Equations I*, no. 3, pp. 1–12, 2009.
- [17] F. M. Atici and S. Sengül, "Modeling with fractional difference equations," *Journal of Mathematical Analysis and Applications*, vol. 369, no. 1, pp. 1–9, 2010.
- [18] F. Chen, "Fixed points and asymptotic stability of nonlinear fractional difference equations," *Electronic Journal of Qualitative Theory of Differential Equations*, vol. 39, pp. 1–18, 2011.
- [19] F. Chen and Y. Zhou, "Attractivity of fractional functional differential equations," *Computers & Mathematics with Applications*, vol. 62, no. 3, pp. 1359–1369, 2011.
- [20] T. A. Burton and T. Furumochi, "Krasnoselskii's fixed point theorem and stability," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 49, no. 4, pp. 445–454, 2002.
- [21] T. A. Burton, "Fixed points, stability, and exact linearization," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 61, no. 5, pp. 857–870, 2005.
- [22] T. A. Burton, *Stability by Fixed Point Theory for Functional Differential Equations*, Dover Publications, Mineola, NY, USA, 2006.
- [23] B. C. Dhage, "Global attractivity results for nonlinear functional integral equations via a Krasnoselskii type fixed point theorem," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 70, no. 7, pp. 2485–2493, 2009.
- [24] C. Jin and J. Luo, "Stability in functional differential equations established using fixed point theory," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 68, no. 11, pp. 3307–3315, 2008.
- [25] Y. N. Raffoul, "Stability in neutral nonlinear differential equations with functional delays using fixed-point theory," *Mathematical and Computer Modelling*, vol. 40, no. 7-8, pp. 691–700, 2004.
- [26] Y. Zhou, *Oscillatory Behavior of Delay Differential Equations*, Science Press, Beijing, China, 2007.
- [27] S. S. Cheng and W. T. Patula, "An existence theorem for a nonlinear difference equation," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 20, no. 3, pp. 193–203, 1993.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

