

Research Article

Parallel Rank of Two Sandpile Models of Signed Integer Partitions

G. Chiaselotti, T. Gentile, G. Marino, and P. A. Oliverio

Dipartimento di Matematica, Università della Calabria, Via Pietro Bucci, 87036 Arcavacata di Rende, Italy

Correspondence should be addressed to G. Marino; gmarino@unical.it

Received 4 December 2012; Accepted 24 January 2013

Academic Editor: Filomena Cianciaruso

Copyright © 2013 G. Chiaselotti et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We introduce the concept of fundamental sequence for a finite graded poset X which is also a discrete dynamical model. The concept of fundamental sequence is a refinement of the concept of parallel convergence time for these models. We compute the parallel convergence time and the fundamental sequence when X is the finite lattice $P(n, r)$ of all the signed integer partitions $a_r, \dots, a_1, b_1, \dots, b_{n-r}$ such that $r \geq a_r \geq \dots \geq a_1 \geq 0 \geq b_1 \geq \dots \geq b_{n-r} \geq -(n-r)$, where $n \geq r \geq 0$, and when X is the sublattice $P(n, d, r)$ of all the signed integer partitions of $P(n, r)$ having exactly d nonzero parts.

1. Introduction

The study of the order relation in particular subsets of integer partitions has been recently addressed from the point of view of the discrete dynamical models. The excellent survey [1] describes several subsets of integer partitions which have a lattice structure in terms of discrete dynamical models. Before going further, we borrow from [1] the following description of *discrete dynamical model* (briefly DDM): “At each (discrete) time step, such a model is in some state, which we call a *configuration*. Configurations are described by combinatorial objects, like graphs, integer partitions, and others, and we will not distinguish a configuration and its combinatorial description. A discrete dynamical model is then defined by an initial configuration and (some) evolution (rules) which (say) under which conditions the (configurations) may be changed, and which (describe) the new configurations one may obtain. These (rules) can generally be applied under a local condition, and (they imply) a local modification of the current configuration. Note that in the general case the evolution (rules) can be applied in several places in a configuration, leading to several configurations. If a configuration c' can be obtained from a configuration c after one application of (some) evolution (rule), we say that c' is a successor of c , or c is a predecessor of c' , which is denoted by $c \rightarrow c'$.” In this paper, we consider integer

partitions; therefore, we use the term *configuration* as an equivalent to *integer partition*. There are two ways in which we can apply the evolution rules on a given configuration: a *sequential* way (in this case, we refer also to *sequential dynamic* of the DDM) and a *parallel* way (in this case, we refer also to *parallel dynamic* of the DDM). In the sequential way, we apply the evolution rules separately on each summand of the configuration. In the parallel way, we apply the evolution rules concurrently on all the summands of the configuration. When we start from the initial configuration and apply the evolution rules in a sequential way, generally we obtain an oriented graph which represents the Hasse diagram of a poset of integer partitions with an order relation \sqsubseteq . In this case, one usually proves that the relation $c \rightarrow c'$ is equivalent to saying that c is covered by c' with respect to the partial order \sqsubseteq . Therefore, several concepts of the classical order theory find their interesting formulation in dynamic terms. The first implicit formulation of the covering rules of an integer partitions lattice as evolution rules of a DDM was given in the famous Brylawski paper [2]. In this paper, Brylawski proposed a dynamical approach in order to study the lattice $L_B(n)$ of all the partitions of a fixed positive integer n with the dominance order. However, the explicit identification of a specific set of integer partitions with a DDM begins in [3, 4]. In the DDM introduced by Goles and Kiwi in [4], denoted by $SPM(n)$, a configuration is an integer partition

$a = (a_1, \dots, a_n)$ of the positive integer n , and each summand of a is identified with a pile of sand grains; hence, the name is *sandpile model* (briefly SPM). In this model, there is only one evolution rule.

Rule 1 (vertical rule). One grain can move from a column to the next one if the difference of height of these two columns is greater than or equal to 2.

In the model $L_B(n)$ (introduced by Brylawski [2]), the movement of a sand grain respects Rule 1 and also the following.

Rule 2 (horizontal rule). If a column containing $p + 1$ grains is followed by a sequence of columns containing p grains and then one column containing $p - 1$ grains, one grain of the first column can slip to the last one.

There are a lot of specializations and extensions of this model which have been introduced and studied under different names, different aspects, and different approaches. The SPM(n) can be also related to the self-organized criticality (SOC) property, introduced by Bak et al. in [5]. The study of such systems has been developed in an algebraic context [6], in a combinatorial game theory setting [4, 7, 8], in the theory of cellular automata [9–11], and in the setting of discrete dynamical systems [12–14].

There exists a wide literature concerning dynamical models related in several ways to the original model SPM(n), [8, 15–29].

When we study the parallel dynamic of an SPM which also has a lattice structure, starting from its initial configuration (denoted by $\hat{0}$), we always arrive after s unit time steps to a unique final configuration (denoted by $\hat{1}$), that is, the maximum of the correspondent lattice. This nonnegative integer s , denoted by $T_{\text{par}}(\hat{0})$, was introduced explicitly in [3, 4]. The number $T_{\text{par}}(\hat{0})$ is relevant to the study of a DDM of integer partitions which is also a graded lattice because it represents a type of “parallel rank” of the graded lattice. The classical concept of rank can be identified with the number of sequential unit steps needed to reach $\hat{1}$ starting from $\hat{0}$, and in [4] this number is denoted by $T_{\text{seq}}(\hat{0})$. Our question is then as follows: is it possible to refine the concept of “parallel rank” $T_{\text{par}}(\hat{0})$ in such a way that this refinement conducts us to have further information concerning the graded lattice? For example, can the parallel dynamic of the model, in some sense, characterize the symmetry of the corresponding lattice? It is very difficult, in general, to establish, for example, when a graded lattice has the Sperner property, or it is rank unimodal or rank symmetric. Can a refinement of the study of the parallel dynamic of an SPM that is also a graded lattice be helpful in addressing these problems? In this paper, we introduce and study some concepts related to the parallel dynamic of SPMs, and we study these concepts in the wider environment of the signed integer partitions, that is, integer partitions whose summands can be also negative. The signed integer partitions have been recently introduced by Andrews in [30] and further studied in [31] from an arithmetical

point of view. The reason why we work in the signed integer partitions environment is not only formal. In the context of the signed integer partitions, it is easier to see the symmetry of the examined models. Almost all these models of the signed integer partitions are equipped with an involution order-reversing map that makes their self-duality immediately clear. Moreover, a signed integer partition is a generalization of a usual integer partition; therefore, all the classical results can be considered as particular cases of their analogues obtained with signed integer partitions. In the sequel, we assume that (X, \sqsubseteq) is a finite graded lattice of signed integer partitions with minimum $\hat{0}$ and maximum $\hat{1}$. We denote by $\text{rank}(X)$ the rank of X . Moreover, we also assume that X is a generic *deterministic* DDM having k distinct evolution rules R_1, \dots, R_k . In our case, this means that at each position at most one rule is applied (however the same rule may be applied at several different positions). In our model X , given two configurations $w, w' \in X$, it results that w is covered by w' (with respect to the partial order \sqsubseteq) if and only if w' is obtained from w with the application of some rule R_i . Therefore, in the sequel, we can always identify $T_{\text{seq}}(\hat{0})$ with $\text{rank}(X)$.

If w and w' are two different configurations of X , we say that w' is a *parallel successor* of w , and we write $w \Rightarrow w'$ or $w' = w \Rightarrow$ if w' is the configuration which is obtained with all the possible parallel applications of the rules R_1, \dots, R_k on the parts of w . If we apply in parallel m_i times the rule R_i on w , for $i = 1, \dots, k$, in order to obtain w' , we write $w \Rightarrow \langle (m_1, \dots, m_k) \rangle w'$, and we also set $M(w) := (m_1, \dots, m_k)$ and $|M(w)| := m_1 + \dots + m_k$. We write $w \Rightarrow \langle M(w), M(u), \dots, M(z) \rangle w'$ if there exists u, v, \dots, z such that $w \Rightarrow \langle M(w) \rangle u \Rightarrow \langle M(u) \rangle v \Rightarrow \dots \Rightarrow \langle M(z) \rangle w'$. Let us note that there is a unique finite sequence (w_0, w_1, \dots, w_s) of configurations in X such that

$$w_0 = \hat{0} \Rightarrow w_1 \Rightarrow \dots \Rightarrow w_{s-1} \Rightarrow w_s = \hat{1}. \quad (1)$$

The sequence in (1) is a chain of length s in X that we call *fundamental chain* of X . It is clear that

$$\sum_{i=1}^{s-1} |M(w_i)| = T_{\text{seq}}(\hat{0}) \quad (2)$$

and $T_{\text{par}}(\hat{0}) = s$. We call *parallel rank* of X the integer $T_{\text{par}}(\hat{0})$ and *sequential rank* of X the integer $T_{\text{seq}}(\hat{0})$. We also call *fundamental sequence* of X the integer sequence

$$(|M(w_0)|, |M(w_1)|, \dots, |M(w_{s-1})|). \quad (3)$$

We say that X is *parallel unimodal* if the fundamental sequence of X is a unimodal sequence; we also say that X is *parallel symmetric* if the fundamental sequence of X is a symmetric sequence. In this paper, we compute the parallel rank and the fundamental sequence in two cases. In the first case, when X is the lattice $P(n, r)$ of all the signed integer partitions of the form $a_r, \dots, a_1, b_1, \dots, b_{n-r}$, where $n \geq r \geq 0$ are two fixed integers and $r \geq a_r \geq \dots \geq a_1 \geq 0 \geq b_1 \geq \dots \geq b_{n-r} \geq -(n-r)$. In the second (more difficult) case, when X is the sublattice $P(n, d, r)$ of all the signed integer partitions

TABLE 1: Parallel rank in $P(n, d, r)$.

Case	Relation between n, d , and r	Parallel rank $T_{\text{par}}(\bar{0})$
I	$d \leq \min\{r, n-r\}$	$n+d-2$
II	$r \geq d \geq 2(n-r)$	$d+r-2$
III	$r \geq d \geq n-r$ and $2(n-r) \geq d \geq n-r$	$2n-r-2$
IV	$n-r \geq d \geq 2r$	$d+n-r-2$
V	$n-r \geq d \geq r$ and $2r \geq d \geq r$	$n+r-2$
VI	$\min\{2r, 2(n-r)\} \geq d \geq \max\{r, n-r\}$	$2n-d-2$
VII	$2r > d \geq 2(n-r)$ and $d \geq \max\{r, n-r\}$	$2r-2$
VIII	$2(n-r) > d \geq 2r$ and $d \geq \max\{r, n-r\}$	$2(n-r)-2$

TABLE 2: Fundamental sequence in $P(n, d, r)$.

Case	Fundamental sequence
Case I	$(1, 2, \dots, d-1, d, d, \dots, d, d-1, \dots, 1)$
Case II	$(2, 4, \dots, 2(n-r-1), 2(n-r), \dots, 2(n-r), 2(n-r)+1, \dots, d-1, d, \dots, d, d-1, \dots, 1)$
Case III	$(2, 4, \dots, 2(d-n+r-1), 2(d-n+r), 2(d-n+r)+1, \dots, d-1, d, \dots, d, d-1, \dots, 1)$
Case IV	$(1, 2, \dots, d-1, d, \dots, d, d-2, \dots, d-2r, d-2r-1, \dots, 1)$
Case V	$(1, 2, \dots, d-1, d, \dots, d, d-2, \dots, 2r-d, 2r-d-1, \dots, 1)$
Case VI.1	$(2, 4, \dots, 2(d-n+r), 2(d-n+r)+1, \dots, d, \dots, d, d-1, \dots, n-r, \dots, n-r, n-r-1, \dots, n-d, \dots, n-d, n-d-1, \dots, 1)$
Case VI.2	$(2, 4, \dots, 2(d-n+r), 2(d-n+r)+1, \dots, d, \dots, d, d-1, \dots, 2(d-n+r), 2(d-n+r-1), \dots, 4n-4r-2d, 4n-4r-2d-1, \dots, n-d, \dots, n-d, n-d-1, \dots, 1)$
Case VI.3	$(2, 4, \dots, 2(d-n+r), 2(d-n+r)+1, \dots, d, \dots, d, d-1, \dots, 2(d-n+r), 2(d-n+r)-1, 2(d-n+r)-3, \dots, 4r-2n-1, 4r-2n-2, \dots, n-d, \dots, n-d, n-d-1, \dots, 1)$
Case VI.4	$(2, 4, \dots, 2(d-n+r), 2(d-n+r)+1, \dots, d-n+2r-1, \dots, d-n+2r-1, d-n+2r, \dots, n-r, \dots, n-r, n-r-1, \dots, 2n-d-2r, 2n-d-2r-2, \dots, 2r-d, 2r-d-1, \dots, 1)$
Case VI.5	$(2, 4, \dots, 2(d-n+r), 2(d-n+r)+1, \dots, d-n+2r-1, \dots, d-n+2r-1, d-n+2r-2, \dots, n-r, \dots, n-r, n-r-1, \dots, 2n-d-2r, 2n-d-2r-2, \dots, 2r-d, 2r-d-1, \dots, 1)$
Case VI.6	$(2, 4, \dots, 2(d-n+r), 2(d-n+r)+1, \dots, d-n+2r-1, \dots, d-n+2r-1, d-n+2r-2, \dots, n-r, \dots, n-r, n-r-1, \dots, 2n-d-2r, 2n-d-2r-2, \dots, 2r-d, 2r-d-1, \dots, 1)$
Case VII.1	$(2, 4, \dots, 2(n-r), \dots, 2(n-r), 2(n-r)-1, \dots, 2(n-r)-1, 2(n-r)-2, 2(n-r)-4, \dots, 4r-2n, 4r-2n-1, \dots, 1)$
Case VII.2	$(2, 4, \dots, 2(n-r), \dots, 2(n-r), 2(n-r)-1, \dots, 2(n-r)-1, 2(n-r), 2(n-r)+1, \dots, r-1, r-1, \dots, 1)$
Case VIII.1	$(2, 4, \dots, 2(d-n+r), 2(d-n+r)+1, \dots, d-n+2r-1, \dots, d-n+2r-1, d-n+2r, \dots, n-r, \dots, n-r, n-r-1, \dots, 2n-2r-d, 2n-2r-d-2, \dots, d-2r, d-2r-1, \dots, 1)$
Case VIII.2	$(2, 4, \dots, 2(d-n+r), 2(d-n+r)+1, \dots, d-n+2r-1, \dots, d-n+2r-1, d-n+2r-2, \dots, 2(d-n+r), 2(d-n+r-1), \dots, 2(2n-2r-d), 2(2n-2r-d)-1, \dots, 2n-2r-d, 2n-2r-d-2, \dots, d-2r, d-2r-1, \dots, 1)$
Case VIII.3	$(2, 4, \dots, 2(d-n+r), 2(d-n+r)+1, \dots, d-n+2r-1, \dots, d-n+2r-1, d-n+2r-2, \dots, n-r+1, n-r, \dots, n-r, n-r-1, \dots, 2n-2r-d, 2n-2r-d-2, \dots, d-2r, d-2r-1, \dots, 1)$

of $P(n, r)$ having exactly d nonzero parts. The paper is structured as follows. In Section 2, we define the lattice $P(n, r)$ and establish the evolution rules which determine its covering relations, and we compute the usual rank, the parallel rank, and the fundamental sequence of $P(n, r)$. In Section 3, we introduce a new integer parameter $1 \leq d \leq n$ and define the sublattice $P(n, d, r)$. We establish three evolution rules which determine the covering relations in $P(n, d, r)$ and determine the usual rank of this lattice. In Section 4, we compute the parallel rank and the fundamental sequence of $P(n, d, r)$ for all the values of the integers parameters n, d , and r . The

complete results of our computations are listed in Tables 1 and 2.

2. The Sandpile Model $P(n, r)$

In this section, we introduce the lattice of signed integer partitions $P(n, r)$, and we examine some of its basic properties. Let n, r be two fixed nonnegative integers such that $n \geq r$. We call (n, r) -signed partition an n -tuple of integers


$$a_r \cdots a_1 \mid b_1 \cdots b_{n-r}, \tag{4}$$

such that

- (i) $a_1, \dots, a_r \in \{1, \dots, r, 0\}$,
- (ii) $b_1, \dots, b_{n-r} \in \{-1, \dots, -(n-r), 0\}$,
- (iii) $a_r \geq \dots \geq a_1 \geq 0 \geq b_1 \geq \dots \geq b_{n-r}$.

In some cases, when n and r are clear from the context, we say simply *signed partition* instead of (n, r) -signed partition. We denote by $P(n, r)$ the set of all the (n, r) -signed partitions. If w is an (n, r) -signed partition, we call *parts* of w the integers $a_r, \dots, a_1, b_1 \cdots b_{n-r}$, *nonnegative parts* of w the integers a_r, \dots, a_1 , and *non-positive parts* of w the integers b_1, \dots, b_{n-r} . We set $\sum(w) := \sum_{i=1}^r a_i + \sum_{j=1}^{n-r} b_j$, and if $m \in \mathbb{Z}$ is such that $\sum(w) = m$, we say that w is an (n, r) -signed partition of m ; in this case, we write $w \vdash m$. We set $w_+ = a_r \cdots a_1$ and $w_- = |b_1 \cdots b_{n-r}|$. In order to simplify the notations, in the numerical examples we write the integers b_1, \dots, b_{n-r} of a generic (n, r) -signed partition without the minus sign. For example, we write $4410 \mid 01125$ instead of $4410 \mid 0(-1)(-1)(-2)(-5)$. Sometimes, when we do not distinguish between nonnegative and nonpositive parts, we write an (n, r) -signed partition without vertical bar $|$. Also, we denote by $|w|_>$ the number of parts of w that are strictly positive, with $|w|_<$ the number of parts of w that are strictly negative, and we set $\|w\| = |w|_> + |w|_<$. If $w = a_r \cdots a_1 \mid b_1 \cdots b_{n-r}$ and $w' = a'_r \cdots a'_1 \mid b'_1 \cdots b'_{n-r}$ are two (n, r) -signed partitions, we set $w_+ = w'_+$ if $a'_i = a_i$ for all $i = r, \dots, 1$, $w_- = w'_-$ if $b'_j = b_j$ for all $j = 1, \dots, n-r$, and $w = w'$ if $w_+ = w'_+$ and $w_- = w'_-$. On $P(n, r)$, we consider the partial order on the components that we denote by \sqsubseteq . As usual, we write $w \sqsubset w'$ if $w \sqsubseteq w'$ and $w \neq w'$. Moreover, we write $w < w'$ (or $w' > w$) if w' covers w . Since $(P(n, r), \sqsubseteq)$ is a finite distributive lattice (because its partial order is on the components), it is also graded with minimum element $0 \cdots 0 \mid (n-r) \cdots (n-r)$ and maximum element $r \cdots r \mid 0 \cdots 0$.

Now let us describe the evolution rules which describe $P(n, r)$ as a discrete dynamical model. In the sequel, if $w \in P(n, r)$, we represent the sequence of the positive parts of w as a not-increasing sequence of columns of stacked squares and the sequence of the negative parts of w as a not-decreasing sequence of columns of stacked squares. We call *pile* a column of stacked squares and *grain* each square of a pile. For example, if $n = 10$, $r = 6$, the configuration




is identified with the signed partition $433100 \mid 0113 \in P(10, 6)$.

Let $w = a_r \cdots a_1 \mid b_1 \cdots b_{n-r} \in P(n, r)$. We formally set $a_0 = 0$, $a_{r+1} = r$, and $b_0 = 0$. If $0 \leq i \leq r+1$, we call a_i the i th-plus pile of w , and if $0 \leq j \leq n-r$, we call b_j the j th-minus pile of w . We call a_i *plus singleton pile* if $a_i = 1$ and b_j *minus singleton pile* if $b_j = -1$. If $1 \leq i \leq r+1$, we say that w has a *plus step* at i if $a_i - a_{i-1} \geq 1$. If $1 \leq j \leq n-r$, we say that w has a *minus step* at j if $|b_j| - |b_{j-1}| \geq 1$.


Remark 1. The choice to set $a_0 = 0$, $a_{r+1} = r$, and $b_0 = 0$ is a formal trick to decrease the number of rules necessary for our model. This means that when we apply the next rules to one element $w \in P(n, r)$, we think that there is an “invisible” extra pile in the imaginary place $r+1$ having exactly r grains, an “invisible” extra pile with 0 grains in the imaginary place to the right of a_1 and to the left of $|$, and another “invisible” extra pile with 0 grains in the imaginary place to the left of b_1 and to the right of $|$. However, the piles corresponding, respectively, to $a_0 = 0$, $a_{r+1} = r$, and $b_0 = 0$ must not be considered as parts of w .

2.1. Evolution Rules.

Rule 1. If w has a plus step at $i+1$, then one grain must be added on the i th-plus pile



Rule 2. One grain must be deleted from the j th-minus pile if w has a minus step at j



We write $w \mapsto^k w'$ (or $w' = w \mapsto^k$) to denote that w' is an n -tuple of integers obtained from w applying the Rule k , for $k = 1, 2$. We also set

$$\nabla(w) = \{w' : w \mapsto^k w', k = 1, 2\}. \tag{8}$$

Theorem 2. *If $w \in P(n, r)$, then $\nabla(w) = \{w' \in P(n, r) : w' > w\}$.*

Proof. We start to show that $\nabla(w) \subseteq \{w' \in P(n, r) : w' > w\}$. Let $w = a_r \cdots a_1 \mid b_1 \cdots b_{n-r} \in P(n, r)$ and $a_{r+1} = r$ be the invisible pile in the place $r+1$. We distinguish the two possible cases related to the previous rules.

Case 1. Let us assume that $r \geq i \geq 1$, $a_i \neq 0$, and that w has a plus step at $i+1$. If $w' = w \mapsto^1$, then $w' = a_r \cdots a_{i+1} (a_i + 1) a_{i-1} \cdots a_1 \mid b_1 \cdots b_{n-r}$. Since there is a plus step at $i+1$, we have $a_{i+1} - a_i \geq 1$; hence, $a_{i+1} \geq a_i + 1 > a_i \geq a_{i-1}$, and this implies that $w' \in P(n, r)$. We must show now that w' covers w in $P(n, r)$. Since w and w' differ between them only in the place i for a_i and $a_i + 1$, respectively, it is clear that there does not exist an element $z \in P(n, r)$ such that $w \sqsubset z \sqsubset w'$. Hence, $w' > w$.

Case 2. If $1 \leq j \leq n-r$ and w has a minus step at j , we apply Rule 2 to w on the minus pile b_j , and we obtain $w' = w \mapsto^2$, where $w' = a_r \cdots a_1 \mid b_1 \cdots b_{j-1} (b_j + 1) b_{j+1} \cdots b_{n-r}$. Since w has a minus step at j , we have $-b_j + b_{j-1} = |b_j| - |b_{j-1}| \geq 1$, and therefore, $w' \in P(n, r)$ because $0 \geq b_{j-1} \geq b_j + 1 > b_j \geq b_{j+1}$.

As in the Case 1, we note that w' covers w in $P(n, r)$ because they differ between them only for a grain in the place j .

We must show now that $\{w' \in P(n, r) : w' \succ w\} \subseteq \nabla(w)$. Let $w = a_n \cdots a_1$ and $w' = a'_n \cdots a'_1$ be two (n, r) -signed partitions in $P(n, r)$. We prove at first that w' covers w if and only if there exists exactly a place k where w and w' are different and that $a'_k - a_k = 1$. In fact, if $a'_k - a_k = 1$ and $a'_i = a_i$ if $i \neq k$, it is immediate to see that w' covers w . In order to prove the other implication, we assume that w' covers w , and we proceed by contradiction. We must distinguish three cases.

Case A. w and w' are equal in all their parts. In this case, $w = w'$, against the hypothesis.

Case B. There exist at least two places k and h , with $h > k$, where w and w' differ, with $l'_k > l_k$ and $l'_h > l_h$. We consider the signed partition

$$u = a'_n \cdots a'_{h+1} a'_h a_{h-1} \cdots a_{k+1} a_k a_{k-1} \cdots a_1. \quad (9)$$

Then, $r \geq a'_n \geq \cdots \geq a'_{h+1} \geq a'_h > a_h \geq a_{h-1} \geq \cdots \geq a_{k+1} \geq a_k \geq a_{k-1} \geq \cdots \geq a_1 \geq -(n-r)$; therefore, $u \in P(n, r)$, and since $w \sqsubset u \sqsubset w'$, this is against the hypothesis that w' covers w .

Case C. There exists exactly one place k where w and w' are different, but $a'_k - a_k > 1$. This implies that $a_k < a_k + 1 < l'_k$. We consider now the signed partition

$$u = a_n \cdots a_{k+1} (a_k + 1) a_{k-1} \cdots a_1. \quad (10)$$

Let us observe then that $w \sqsubset u \sqsubset w'$ and $u \in P(n, r)$, and this is absurd because w' covers w .

Therefore, if $w = a_r \cdots a_1 \mid b_1 \cdots b_{n-r}$ and $w' = a'_r \cdots a'_1 \mid b'_1 \cdots b'_{n-r}$ are two (n, r) -signed partitions such that $w \prec w'$, the condition that w' covers w can occur just by one of the following possibilities:

- (i) there exists exactly an index $i \in \{1, \dots, r\}$, with $a'_i = a_i + 1$, and so $w' = w \mapsto^1$,
- (ii) there exists exactly an index $j \in \{1, \dots, n-r\}$, with $b'_j = b_j + 1$, and so $w' = w \mapsto^2$.

According to the definition of $\nabla(w)$, the thesis follows. \square

Now let us study the sequential and the parallel dynamic of the model $P(n, r)$. In such a context, we call *configuration* an element of $P(n, r)$. The initial configuration is $\hat{0} = 0 \cdots 0 \mid (r-n) \cdots (r-n)$. Each configuration converges, in sequential and in parallel, toward the unique fixed point $\hat{1} = r \cdots r \mid 0 \cdots 0$ because of the lattice structure of the model. Let us note that if w is a configuration, when we use the evolution rules in parallel, on each column of w , we can apply (due to the nature of Rules 1 and 2) at most one evolution rule; hence, our model is deterministic. Obviously, $T_{\text{seq}}(w)$ is independent of the order in which the sites are updated because $P(n, r)$ is a finite distributive and hence also a graded lattice. From the previous theorem, we can easily obtain the rank function of $P(n, r)$.

Proposition 3. *Let one denote by ϱ the rank function of the graded lattice $P(n, r)$. Then, $\varrho(w) = \sum(w) - \sum(\hat{0})$ for each $w \in P(n, r)$ and $T_{\text{seq}}(\hat{0}) = r^2 + (n-r)^2$.*

Proof. Let $w, w' \in P(n, r)$ such that $w' \succ w$. From the previous theorem, w' is obtained from w with an application of Rules 1 or 2. If we take then the function $\varrho(w) := \sum(w) - \sum(\hat{0})$, we obtain $\varrho(w') = \varrho(w) + 1$ and $\varrho(\hat{0}) = 0$. Hence, ϱ is exactly the rank function of $P(n, r)$. The second part of the thesis is followed by $T_{\text{seq}}(\hat{0}) = \varrho(\hat{1}) = \sum(\hat{1}) - \sum(\hat{0}) = r^2 + (n-r)^2$. \square

For the sake of completeness, we recall that in [36] we have computed the rank of the sublattice $S(n, r)$ of all the signed partitions of $P(n, r)$ having positive and negative parts all distinct between them.

We recall now the concept of involution poset. An *involution poset* (IP) is a poset (X, \leq, c) with a unary operation $c : x \in X \mapsto x^c \in X$, such that

$$(I1) \quad (x^c)^c = x, \text{ for all } x \in X,$$

$$(I2) \quad \text{if } x, y \in X \text{ and if } x \leq y, \text{ then } y^c \leq x^c.$$

The map c is called *complementation* of X and x^c the *complement* of x . Let us observe that if X is an involution poset, by (I1) it follows that c is bijective, and by (I1) and (I2) it holds that if $x, y \in X$ are such that $x < y$, then $y^c < x^c$. If (X, \leq, c) is an involution poset and if $Z \subseteq X$, we will set $Z^c = \{z^c : z \in Z\}$. We note that if X is an involution poset, then X is a self-dual poset because from (I1) and (I2) it follows that if $x, y \in X$ we have that $x \leq y$, if and only if $y^c \leq x^c$, and this is equivalent to say that the complementation is an isomorphism between X and its dual poset X^* .

Now, if $w = a_r \cdots a_1 \mid b_1 \cdots b_{n-r} \in P(n, r)$, we set $w^c = (r-a_1) \cdots (r-a_r) \mid (|b_{n-r}| - (n-r)) \cdots (|b_1| - (n-r))$, and let us note that w^c is still a signed partition in $P(n, r)$; therefore, we can define a unary operation $c : P(n, r) \rightarrow P(n, r)$ such that $w \mapsto w^c$. Then, it is immediate to verify that $(P(n, r), \sqsubseteq, c)$ is an involution poset. If $w = a_r \cdots a_1 \mid b_1 \cdots b_{n-r} \in P(n, r)$, we also set $w^t = (-b_{n-r}) \cdots (-b_1) \mid (-a_1) \cdots (-a_r)$, and we call w^t the *transposed* of w . Let us note that $w^t \in P(n, n-r)$.

Proposition 4. *We have:*

$$(i) \quad |P(n, r)| = \binom{2r}{r} \binom{2(n-r)}{n-r}$$

$$(ii) \quad P(n, r) \cong P(n, n-r).$$

Proof. (i) It is sufficient to observe that $P(n, r)$ can be identified with the set of all the ordered pairs (z_1, z_2) , where z_1 is a decreasing string of length r on the ordered alphabet $r > \cdots > 1 > 0$ and z_2 is a decreasing string of length $n-r$ on the ordered alphabet $0 > -1 > \cdots > -(n-r)$; therefore, $|P(n, r)| = \binom{(r+1)+r-1}{r} \binom{(n-r+1)+(n-r)-1}{n-r} = \binom{2r}{r} \binom{2(n-r)}{n-r}$.

(ii) We define the map $\phi : P(n, r) \rightarrow P(n, n-r)$ such that $\phi(w) = (w^t)^c$, for all $w \in P(n, r)$. We prove then that ϕ is an isomorphism of posets. By part (i), it follows that $|P(n, r)| = |P(n, n-r)|$; therefore, ϕ is bijective since it is easily seen that it is injective. If $w, w' \in P(n, r)$, the condition $w \sqsubseteq w' \Leftrightarrow \phi(w) \sqsubseteq \phi(w')$ follows at once from the conditions (I1), (I2) and because the transposed map $w \mapsto w^t$ is order-reversing and such that $(w^t)^t = w$. \square

Theorem 5. The parallel rank $T_{par}(\widehat{0})$ in $P(n, r)$ is given by

$$T_{par}(\widehat{0}) = \begin{cases} 2r - 1 & \text{if } r \geq n - r \\ 2(n - r) - 1 & \text{if } r < n - r. \end{cases} \quad (11)$$

The fundamental sequence of $P(n, r)$ is equal to

$$\begin{aligned} & (2, 4, \dots, 2(n - r), 2(n - r), \dots, 2(n - r), 2(n - r) + 1, \dots, r, r - 1, \dots, 1) && \text{if } r > 2(n - r) \\ & (2, 4, \dots, 2(n - r), 2(n - r), \dots, 2(n - r), 2(n - r) - 2, \dots, 4r - 2n, 4r - 2n - 1, \dots, 1) && \text{if } n - r \leq r \leq 2(n - r) \\ & (2, 4, \dots, 2r, 2r, \dots, 2r, 2r + 1, \dots, n - r, n - r - 1, \dots, 1) && \text{if } n - r > 2r \\ & (2, 4, \dots, 2r, 2r, \dots, 2r, 2r - 2, \dots, 2n - 4r, 2n - 4r - 1, \dots, 1) && \text{if } r \leq n - r \leq 2r. \end{aligned} \quad (12)$$

Proof. By Proposition 4 (ii), we can consider $r \geq n - r$. The initial configuration is $\widehat{0} = 0 \cdots 0 \mid (n - r) \cdots (n - r)$. Since our model is deterministic, we can consider separately the action of the Rules 1 and 2. On the nonnegative part, Rule 1 is applied as in the sequence

$$(1, 2, \dots, r, (r - 1), \dots, 1) \quad (13)$$

of length $2r - 1$ and similarly Rule 2 on the nonpositive part as in the sequence

$$(1, 2, \dots, (n - r), (n - r - 1), \dots, 1), \quad (14)$$


of length $2(n - r) - 1$. It follows that the parallel rank of $P(n, r)$ is equal to the maximum of the length of the two sequences (13) and (14) and that the fundamental sequence is the sum of the two sequences. The other parts follow easily by replacing r with $n - r$ and conversely. \square

3. The Submodel $P(n, d, r)$ with Three Evolution Rules


In this section, we consider a submodel of $P(n, r)$ which presents an evolution rule which permits us to move one single grain from the right part of a configuration into its left part. Let d be a positive integer such that $d \leq n$. We define $P(n, d, r)$ as the set of all the (n, r) -signed partitions $w \in P(n, r)$ such that $\|w\| = d$. It is immediate to see that $P(n, d, r)$ is a sublattice of $P(n, r)$. The lattice $P(n, d, r)$ is related to some extremal combinatorial sum problems studied in [32–35]. The interesting problem is to determine some evolution rules which describe such a lattice as a discrete dynamical model. We consider then the following rules.

3.1. Evolution Rules.


Rule R_1 . If the i th-plus pile has at least one grain, and if w has a plus step at $i + 1$, then one grain must be added on the i th-plus pile



Rule R_2 . If there are some minus singleton piles, then the first from the left of them must be shifted to the side of the lowest not empty plus pile



Rule R_3 . One grain must be deleted from the j th-minus pile if w has a minus step at j and $|b_j| > 1$



Remark 6. (i) In the previous rule, the lowest not empty plus pile can also be the invisible column in the place $r + 1$. In this case, all the plus piles are empty, and an eventual minus singleton pile must be shifted in the place r .

(ii) We take implicitly for intended that the shift of one minus singleton pile into a plus singleton pile can be made if the number of plus piles (excluding $a_{r+1} = r$) with at least a grain is less than r (otherwise we obtain a configuration that does not belong to $P(n, d, r)$).

We write $w \rightarrow^k w'$ (or $w' = w \rightarrow^k$) to denote that w' is an n -tuple of integers obtained from w applying Rule R_k , for $k = 1, 2, 3$. We also set

$$\nabla(w) = \{w' : w \rightarrow^k w', k = 1, 2, 3\}. \quad (18)$$

Theorem 7. If $w \in P(n, d, r)$, then $\nabla(w) = \{w' \in P(n, d, r) : w' > w\}$.

Proof. We start to show that $\nabla(w) \subseteq \{w' \in P(n, d, r) : w' > w\}$. Let $w = a_r \cdots a_1 \mid b_1 \cdots b_{n-r} \in P(n, d, r)$ and $a_{r+1} = r$ be the invisible pile in the place $r + 1$. We distinguish the three possible cases related to the previous rules.

Case 1. Let us assume that $r \geq i \geq 1$, $a_i \neq 0$ and that w has a plus step at $i + 1$. If $w' = w \rightarrow^1$, then $w' = a_r \cdots a_{i+1}(a_i + 1)a_{i-1} \cdots a_1 \mid b_1 \cdots b_{n-r}$. It is clear that $\|w'\| = d$ because $a_i \neq 0$. Since there is a plus step at $i + 1$, we have $a_{i+1} - a_i \geq 1$;

hence, $a_{i+1} \geq a_i + 1 > a_i \geq a_{i-1}$, and this implies that $w' \in P(n, d, r)$. We must show now that w' covers w in $P(n, d, r)$. Since w and w' differ between them only in the place i for a_i and $a_i + 1$, respectively, it is clear that there does not exist an element $z \in P(n, d, r)$ such that $w \sqsubset z \sqsubset w'$. Hence, $w' \succ w$.

Case 2. Let us assume that in w there is a minus singleton pile b_j , for some $1 \leq j \leq n - r$. Since $a_{r+1} = r > 0$, we can assume that $a_{i+1} > 0$, $a_i = 0$, for some $1 \leq i \leq r$. This means that w has the following form: $w = a_r \cdots a_{i+1} 00 \cdots 0 \mid 0 \cdots 0(-1)b_{j+1} \cdots b_{n-r}$. Applying Rule R_2 to w , we obtain $w' = w \rightarrow^2$, where $w' = a_r \cdots a_{i+1} 10 \cdots 0 \mid 0 \cdots 00b_{j+1} \cdots b_{n-r}$. It is clear then that $w' \in P(n, r)$ and $\|w'\| = d$ since w' is obtained from w with only a shift of the pile -1 to the left in the place i . Let us note that the only elements $z_1, z_2 \in P(n, r)$ such that $w \sqsubset z_1 \sqsubset w'$, and $w \sqsubset z_2 \sqsubset w'$ are $z_1 = a_r \cdots a_{i+1} 10 \cdots 0 \mid 0 \cdots 0(-1)b_{j+1} \cdots b_{n-r}$ and $z_2 = a_r \cdots a_{i+1} 00 \cdots 0 \mid 0 \cdots 00b_{j+1} \cdots b_{n-r}$, but $\|z_1\| = d + 1$ and $\|z_2\| = d - 1$, hence z_1, z_2 are not elements of $P(n, d, r)$. This implies that w' covers w in $P(n, d, r)$.

Case 3. If $1 \leq j \leq n - r$ and w has a minus step at j , we apply Rule R_3 to w on the minus pile b_j , and we obtain $w' = w \rightarrow^3$, where $w' = a_r \cdots a_1 \mid b_1 \cdots b_{j-1}(b_j + 1)b_{j+1} \cdots b_{n-r}$. Since w has a minus step at j , we have $-b_j + b_{j-1} = |b_j| - |b_{j-1}| \geq 1$, therefore $w' \in P(n, r)$ because $0 \geq b_{j-1} \geq b_j + 1 > b_j \geq b_{j+1}$ and $\|w'\| = d$ since $b_j \leq -2$ implies $b_j + 1 < 0$. As in Case 1, we note that w' covers w in $P(n, d, r)$ because they differ between them only for a grain in the place j . We now must show that $\{w' \in P(n, d, r) : w' \succ w\} \subseteq \nabla(w)$. Let $w'' = a_r'' \cdots a_1'' \mid b_1'' \cdots b_{n-r}''$ be a generic element of $P(n, d, r)$ such that $w'' \sqsupset w$. If we show that there exists an element $w' = a_r' \cdots a_1' \mid b_1' \cdots b_{n-r}'$ of $P(n, d, r)$ such that $w' \in \nabla(w)$ and $w'' \sqsupseteq w'$, we complete the proof of the theorem. Since $w'' \sqsupset w$, there is a place where the corresponding component of w'' is an integer strictly bigger than the integer component of w corresponding to the same place. We distinguish several cases.

Case A. We assume that $a_i'' > a_i$ and $a_{i+1} \geq a_i + 1$ for some $i \in \{r - 1, \dots, 1\}$. In this case, we can apply Rule R_1 in the place i to obtain $w' = w \rightarrow^1$ such that $w'' \sqsupseteq w'$.

Case B. We assume that $a_i'' > a_i$ and $a_{i+1} = a_i$ for some $i \in \{r - 1, \dots, 1\}$. In this case, we have $a_{i+1}'' \geq a_i'' > a_i = a_{i+1}$, that is, $a_{i+1}'' \geq a_{i+1} + 1$; therefore if $a_{i+2} \geq a_{i+1} + 1$, we can apply Rule R_1 in the place $i + 1$ to obtain w' , otherwise we have $a_{i+2} = a_{i+1} = a_i$. Iterating this procedure, to each step $k \geq 1$ we can apply Rule R_1 in the place $i + k$ to obtain w' or it necessarily results that $a_{i+k} = \cdots = a_{i+1} = a_i$. Hence, if for no one k we can apply Rule R_1 in the place $i + k$, we necessarily arrive to the condition $a_r = \cdots = a_{i+1} = a_i$. Since $r \geq a_i'' > a_i$, it must be $a_r < r$; therefore, we can apply Rule R_1 in the place r to obtain $a_r' = a_r + 1$, with $a_r'' \geq a_r'$ because $a_r'' \geq \cdots \geq a_i'' > a_i = \cdots = a_r$.

Case C. We assume that $a_r'' > a_r$. In this case, we can apply Rule R_1 in the place r .

Case D. We assume that $0 > b_j'' > b_j$, for some $j \in \{1, \dots, n - r\}$. In this case, we can apply Rule R_3 in the place j .


Case E. We assume that $0 = b_j'' > b_j$ and $b_j \leq -2$, for some $j \in \{1, \dots, n - r\}$. Also in this case, we can apply Rule R_3 in the place j .

Case F. We assume that $0 = b_j'' > b_j = -1$, for some $j \in \{1, \dots, n - r\}$. In this case, the number of negative parts of w'' is strictly lower than the number of negative parts of w , and since $\|w\| = \|w''\| = d$, it follows that there exists at least one index $i \in \{1, \dots, r\}$ such that $a_i'' > 0$ and $a_i = 0$. We choose then such index i maximal, so that we have $i = r$ or $i \leq r - 1$ and $a_{i+1} > 0$.

We suppose at first that $i = r$. In this case, we have $a_r'' \geq 1$ and $0 = a_r = \cdots = a_1$; therefore, we can apply Rule R_2 to move the “negative” grain from the place j into the place r , so that the (n, r) -partition $w' = 100 \cdots 0 \mid 0 \cdots 0b_{j+1} \cdots b_{n-r}$ is such that $\|w'\| = d$ and $w'' \sqsupseteq w'$.

If $i \leq r - 1$ and $a_{i+1} > 0$, we apply again Rule R_2 to move the “negative” grain from the place j into the place i , so that the (n, r) -partition $w' = a_r \cdots a_{i+1} 1 \cdots 0 \mid 0 \cdots 0b_{j+1} \cdots b_{n-r}$ is such that $\|w'\| = d$ and $w'' \sqsupseteq w'$. \square

An analogous statement of the previous theorem was proven in [36] for the sublattice of all the signed integer partitions of $P(n, d, r)$ having positive and negative parts all distinct between them. Below, we draw the Hasse diagram of the lattice $P(4, 3, 2)$ by using the evolution Rules R_1, R_2 , and R_3 starting from the minimum element of this lattice, which is $10 \mid 22$. We label a generic edge of the next diagram with the symbol k if it leads to a production that uses the Rule R_k , for $k \in \{1, 2, 3\}$



4. Sequential and Parallel Dynamics in $P(n, d, r)$

In this section, we study the parallel dynamic of the model $P(n, d, r)$. In such a context, we call *configuration* a generic element of $P(n, d, r)$. The initial configuration is $\widehat{0}$. Each configuration converges, in sequential and in parallel, toward the unique fixed point $\widehat{1}$ because of the lattice structure of the model. Let us note that if w is a configuration, when we use the evolution rules in parallel, on each column of w we can apply (due to the nature of the Rules R_1, R_2 , and R_3) at most one evolution rule, hence our model is deterministic. We denote by $T_{\text{seq}}(w)$ and $T_{\text{par}}(w)$ the number of time steps required to reach $\widehat{1}$ starting from the configuration w , using, respectively, the sequential or the parallel updating scheme. Obviously $T_{\text{seq}}(w)$ is independent of the order in which the sites are updated because $P(n, d, r)$ is a finite distributive, and hence also a graded lattice (let us note that $P(n, d, r)$ is distributive because it is a sublattice of the distributive lattice $P(n, r)$). In the next proposition, we determine the rank function of $P(n, d, r)$.

Proposition 8. *The map $\rho : P(n, d, r) \rightarrow \mathbb{N}$ such that $\rho(w) = \sum(w) - \sum(\widehat{0}) - (|w|_{>} - |\widehat{0}|_{>})$ is the rank function of $P(n, d, r)$.*

Proof. We denote by ϱ the rank function of the graded lattice $P(n, r)$. If w and w' are two (n, r) -signed partitions and $w > w'$, then $\sum(w) = \sum(w') + 1$. It follows that $\varrho(w) = \sum(w) - \sum(\widehat{0})$, for each $w \in P(n, r)$. Let now $w > w_t > \dots > w_1 > \widehat{0}$ be any saturated chain from $\widehat{0}$ to w in the lattice $P(n, d, r)$. Let us assume that in this chain w is obtained from $\widehat{0}$ with k applications of R_2 , for some integer $k \geq 0$. To each step $l \in \{1, \dots, t\}$ where we apply R_2 , there is the following situation: $w_l \sqsupset u_l \sqsupset w_{l-1}$, for one only element $u_l \in P(n, r) \setminus P(n, d, r)$. This means that $\varrho(w) = (t + 1) + k$, that is, $\rho(w) = t + 1 = \varrho(w) - k$. The integer k is also the difference between the number of positive parts of w and the number of positive parts of $\widehat{0}$. Hence, the thesis follows. \square

By using the rank function, we can compute the sequential convergence time of any element $w \in P(n, d, r)$ as in the following:

$$T_{\text{seq}}(w) = \rho(\widehat{1}) - \rho(w). \quad (20)$$

In particular, if we denote by $\text{rank}(P(n, d, r))$ the rank of the lattice $P(n, d, r)$, then $T_{\text{seq}}(\widehat{0}) = \rho(\widehat{1}) - \rho(\widehat{0}) = \rho(\widehat{1}) = \text{rank}(P(n, d, r))$. In the next proposition, we compute this number.

Proposition 9. *the following result holds:*

$$T_{\text{seq}}(\widehat{0}) = \begin{cases} d(n-1) & \text{if } d \leq r \text{ and } d \leq n-r, \\ dr - d + (n-r)^2 & \text{if } r \geq d \geq n-r, \\ d(n-r) - d + r^2 & \text{if } n-r \geq d \geq r, \\ r^2 + (n-r)^2 - d & \text{if } d \geq r \text{ and } d \geq n-r. \end{cases} \quad (21)$$

Proof. The minimal and maximal elements in $P(n, d, r)$ are, respectively,

$$\widehat{0} := \begin{cases} 0 \cdots 0 \mid 0 \cdots 0(n-r) \cdots (n-r) & \text{if } n-r \geq d, \\ 1 \cdots 1 \ 0 \cdots 0 \mid (n-r) \cdots (n-r) & \text{if } d \geq n-r, \end{cases} \quad (22)$$

$$\widehat{1} := \begin{cases} r \cdots r \ 0 \cdots 0 \mid 0 \cdots 0 & \text{if } r \geq d, \\ r \cdots r \mid 0 \cdots 0 \ 1 \cdots 1 & \text{if } d \geq r. \end{cases}$$

Then,

$$\sum(\widehat{0}) := \begin{cases} d(r-n) & \text{if } n-r \leq d, \\ d-n+r+(n-r)(r-n) & \text{if } d \leq n-r, \end{cases} \quad (23)$$

$$\sum(\widehat{1}) := \begin{cases} dr & \text{if } r \leq d, \\ r^2 + r - d & \text{if } d \leq r, \end{cases}$$

while the number of positive parts in $\widehat{0}$ and $\widehat{1}$ is, respectively,

$$|\widehat{0}|_{>} := \begin{cases} 0 & \text{if } n-r \leq d, \\ d-n+r & \text{if } d \leq n-r, \end{cases} \quad (24)$$

$$|\widehat{1}|_{>} := \begin{cases} d & \text{if } r \leq d, \\ r & \text{if } d \leq r. \end{cases}$$

The thesis follows then from simple arithmetic calculations and from Proposition 8, because $\text{rank}(P(n, d, r)) = \rho(\widehat{1})$. \square

The next theorem is the main result of this section. Here we determine the parallel rank of the lattice $P(n, d, r)$, for all the values of n, d , and r . In the proof of the next theorem, we also determine the fundamental sequence of $P(n, d, r)$ in almost all the values of n, d , and r . We do not provide the construction of the fundamental sequence for all the parameters n, d , and r because several cases have similar constructions. However, for completeness, in Tables 1 and 2 we provide a complete description of the parallel rank value and of the fundamental sequence for all the values n, d , and r .

Theorem 10.

- (I) If $d \leq \min\{r, n-r\}$, $T_{\text{par}}(\widehat{0}) = n + d - 2$.
- (II) If $r \geq d \geq 2(n-r)$, $T_{\text{par}}(\widehat{0}) = d + r - 2$.
- (III) If $r \geq d \geq n-r$ and $2(n-r) \geq d \geq n-r$, $T_{\text{par}}(\widehat{0}) = 2n - r - 2$.
- (IV) If $n-r \geq d \geq 2r$, $T_{\text{par}}(\widehat{0}) = d + n - r - 2$.
- (V) If $n-r \geq d \geq r$ and $2r \geq d \geq r$, $T_{\text{par}}(\widehat{0}) = n + r - 2$.
- (VI) If $\min\{2r, 2(n-r)\} \geq d \geq \max\{r, n-r\}$, $T_{\text{par}}(\widehat{0}) = 2n - d - 2$.
- (VII) If $2r > d \geq 2(n-r)$ and $d \geq \max\{r, n-r\}$, $T_{\text{par}}(\widehat{0}) = 2r - 2$.
- (VIII) If $2(n-r) > d \geq 2r$ and $d \geq \max\{r, n-r\}$, $T_{\text{par}}(\widehat{0}) = 2(n-r) - 2$.

Proof. In all cases, let us denote the sequence of configurations as in (1).

Case I ($d \leq \min\{r, n-r\}$). If $d \leq \min\{r, n-r\}$, then $\widehat{0} = 0 \cdots 0 \mid 0 \cdots 0 (n-r) \cdots (n-r)$ and $\widehat{1} = r \cdots r \ 0 \cdots 0 \mid 0 \cdots 0$. Therefore,

$$\widehat{0} \Rightarrow \langle (0, 0, 1), (0, 0, 2), \dots, (0, 0, d-1) \rangle \ 0 \cdots 0 \mid 0 \cdots 0 (n-r-d+1) \cdots (n-r) := w_1. \quad (25)$$

Starting from w_1 , we obtain

$$w_1 \Rightarrow \langle (0, 0, d), (0, 0, d), \dots, (0, 0, d) \rangle \ 0 \cdots 0 \mid 0 \cdots 0 \ 1 \cdots d := w_2, \quad (26)$$

where $(0, 0, d)$ is repeated $n-r-d$ times. The next steps are

$$w_2 \Rightarrow \langle (0, 1, d-1), (1, 1, d-2), \dots, (d-2, 1, 1), (d-1, 1, 0) \rangle \ d (d-1), \cdots \ 1 \ 0 \cdots 0 \mid 0 \cdots 0 := w_3, \quad (27)$$

$$w_3 \Rightarrow \langle (d, 0, 0), \dots, (d, 0, 0) \rangle \ r (r-1) \cdots (r-d+1) \ 0 \cdots 0 \mid 0 \cdots 0 := w_4,$$

where $(d, 0, 0)$ is repeated $r-d$ times. Finally, we have

$$w_4 \Rightarrow \langle (d-1, 0, 0), (d-2, 0, 0) \cdots, (2, 0, 0), (1, 0, 0) \rangle \ r \cdots r \ 0 \cdots 0 \mid 0 \cdots 0 = \widehat{1}. \quad (28)$$

From the previous computations, it follows that the fundamental sequence of $P(n, d, r)$ is

$$(1, 2, \dots, d-1, d, \dots, d, d-1, \dots, 1), \quad (29)$$

where the integer d is repeated exactly $n-d$ times; hence, $T_{\text{par}}(\widehat{0}) = n+d-2$. Let us note that the sequence (29) is unimodal and symmetric.

Case II ($r \geq d \geq 2(n-r)$). In this case, $\widehat{0} = 1 \cdots 1 \ 0 \cdots 0 \mid (n-r) \cdots (n-r)$ and $\widehat{1} = r \cdots r \ 0 \cdots 0 \mid 0 \cdots 0$. Since $d \geq 2(n-r)$, we have $d-n+r \geq n-r$ therefore,

$$\widehat{0} \Rightarrow \langle (1, 0, 1), (2, 0, 2), \dots, (n-r-1, 0, n-r-1) \rangle \ w_1, \quad (30)$$

with $w_1 := (n-r) (n-r-1) \cdots 1 \ 1 \cdots 1 \ 0 \cdots 0 \mid 1 \ 2 \cdots (n-r)$, where 1 is repeated exactly $d-2n+2r+1$ times.

Since $r \geq 2(n-r)$, by starting from w_1 we obtain

$$w_1 \Rightarrow \langle (n-r, 1, n-r-1), (n-r+1, 1, n-r-2), \dots, (2n-2r-1, 1, 0) \rangle \ w_2, \quad (31)$$

with $w_2 := (2n-2r) (2n-2r-1) \cdots 1 \ 1 \cdots 1 \ 0 \cdots 0 \mid 0 \cdots 0$, where 1 is repeated exactly $d-2n+2r+1$ times.

Now, from w_2 we have

$$w_2 \Rightarrow \langle (2n-2r, 0, 0), (2n-2r-1, 0, 0), \dots, (d-1, 0, 0) \rangle \ d (d-1) \cdots 1 \ 0 \cdots 0 \mid 0 \cdots 0 := w_3. \quad (32)$$

At this point we obtain

$$w_3 \Rightarrow \langle (d, 0, 0), \dots, (d, 0, 0) \rangle \ r (r-1) \cdots (r-d+1) \ 0 \cdots 0 \mid 0 \cdots 0 := w_4, \quad (33)$$

where $(d, 0, 0)$ is repeated $r-d$ times. Finally, we have

$$w_4 \Rightarrow \langle (d-1, 0, 0), (d-2, 0, 0) \cdots, (2, 0, 0), (1, 0, 0) \rangle \ r \cdots r \ 0 \cdots 0 \mid 0 \cdots 0 = \widehat{1}. \quad (34)$$

Thus, the parallel rank $T_{\text{par}}(\widehat{0})$ is equal to $d+r-2$, and the fundamental sequence of $P(n, d, r)$ is

$$(2, 4, \dots, 2(n-r-1), 2(n-r), \dots, 2(n-r), 2(n-r)+1, \dots, d-1, d, \dots, d, d-1, \dots, 1), \quad (35)$$

and so $P(n, d, r)$ in this case is parallel unimodal but not parallel symmetric.

In a similar way, we can compute in all other cases the parallel rank and the fundamental sequence. The details are left to the interested reader. Here are the results.

Case III ($r \geq d \geq n-r$ and $2(n-r) \geq d \geq n-r$). In this case, $T_{\text{par}}(\widehat{0}) = 2n-r-2$, and the fundamental sequence is

$$(2, 4, \dots, 2(d-n+r-1), 2(d-n+r), 2(d-n+r) + 1, \dots, d-1, d, \dots, d, d-1, \dots, 1). \quad (36)$$

It is unimodal and not symmetric.

Case IV ($n-r \geq d \geq 2r$). In this case, $T_{\text{par}}(\widehat{0}) = d+n-r-2$ and the fundamental sequence is

$$(1, 2, \dots, d-1, d, \dots, d, d-2, \dots, d-2r, d-2r-1, \dots, 1). \quad (37)$$

It is unimodal, and if $r \neq 0$, it is not symmetric.

Case V ($n-r \geq d \geq r$ and $2r \geq d \geq r$). In this case, $T_{\text{par}}(\widehat{0}) = n+r-2$. The fundamental sequence is

$$(1, 2, \dots, d-1, d, \dots, d, d-2, \dots, 2r-d, 2r-d-1, \dots, 1). \quad (38)$$

It is unimodal, and if $d \neq r$, it is not symmetric.

Case VI ($\min\{2r, 2(n-r)\} \geq d \geq \max\{r, n-r\}$). In this case, $T_{\text{par}}(\widehat{0}) = 2n-d-2$. The fundamental sequence depends on the relations between $r, n-r$, and d in the following way.

(VI.1) If $r \geq n - r$ and $2d - 1 \leq 3(n - r)$, the fundamental sequence is

$$(2, 4, \dots, 2(d - n + r), 2(d - n + r) + 1, \dots, d, \dots, \\ d, d - 1, \dots, n - r, \dots, n - r, n - r - 1, \dots, \\ n - d, \dots, n - d, n - d - 1, \dots, 1). \quad (39)$$

(VI.2) If $r \geq n - r$, $2d - 1 > 3(n - r)$, and $d \leq 3(n - r) - r + 1$, the fundamental sequence is

$$(2, \dots, 2(d - n + r), 2(d - n + r) + 1, \dots, d, \dots, d, \\ d - 1, \dots, 2(d - n + r), 2(d - n + r - 1), \dots, \\ 4n - 4r - 2d, 4n - 4r - 2d - 1, \dots, n - d, \dots, \\ n - d, n - d - 1, \dots, 1). \quad (40)$$

(VI.3) If $r \geq n - r$, $2d - 1 > 3(n - r)$ and $d > 3(n - r) - r + 1$, the fundamental sequence is

$$(2, \dots, 2(d - n + r), 2(d - n + r) + 1, \dots, d, \dots, d, d - 1, \\ \dots, 2(d - n + r), 2(d - n + r) - 1, 2(d - n + r) - 3, \\ \dots, 4r - 2n - 1, 4r - 2n - 2, \dots, n - d, \dots, n - d, \\ n - d - 1, \dots, 1). \quad (41)$$

(VI.4) If $r \leq n - r$ and $d \leq 2n - 3r + 1$, the fundamental sequence is

$$(2, 4, \dots, 2(d - n + r), 2(d - n + r) + 1, \dots, d - n + 2r - 1, \\ \dots, d - n + 2r - 1, d - n + 2r, \dots, n - r, \dots, n - r, \\ n - r - 1, \dots, 2n - d - 2r, 2n - d - 2r - 2, \dots, \\ 2r - d, 2r - d - 1, \dots, 1). \quad (42)$$

(VI.5) If $r \leq n - r$, $d > 2n - 3r + 1$, and $2d - 1 \leq 3(n - r)$, the fundamental sequence is

$$(2, 4, \dots, 2(d - n + r), 2(d - n + r) + 1, \dots, d - n + 2r - 1, \\ \dots, d - n + 2r - 1, d - n + 2r - 2, \dots, n - r, \dots, \\ n - r, n - r - 1, \dots, 2n - d - 2r, 2n - d - 2r - 2, \\ \dots, 2r - d, 2r - d - 1, \dots, 1). \quad (43)$$

(VI.6) If $r \leq n - r$, $d > 2n - 3r + 1$, and $2d - 1 > 3(n - r)$, the fundamental sequence is

$$(2, 4, \dots, 2(d - n + r), 2(d - n + r) + 1, \dots, d - n + 2r - 1, \\ \dots, d - n + 2r - 1, d - n + 2r - 2, \dots, n - r, \dots, n - r, \\ n - r - 1, \dots, 2n - d - 2r, 2n - d - 2r - 2, \dots, \\ 2r - d, 2r - d - 1, \dots, 1). \quad (44)$$

In those cases $P(n, d, r)$ is parallel unimodal but not parallel symmetric.

Case VII ($2r > d \geq 2(n - r)$ and $d \geq \max\{r, n - r\}$). In this case, the parallel rank is $T_{\text{par}}(\widehat{0}) = 2n - d - 2$. The fundamental sequence depends on the relations between r , $n - r$, and d in the following way.

(VII.1) If $3r \leq 2(n - r)$, the fundamental sequence is

$$(2, 4, \dots, 2(n - r), \dots, 2(n - r), 2(n - r) - 1, \dots, \\ 2(n - r) - 1, 2(n - r) - 2, 2(n - r) - 4, \dots, \\ 4r - 2n, 4r - 2n - 1, \dots, 1). \quad (45)$$

In this case $P(n, d, r)$ is parallel unimodal but not parallel symmetric.

(VII.2) If $3r > 2(n - r)$, the fundamental sequence is

$$(2, 4, \dots, 2(n - r), \dots, 2(n - r), 2(n - r) - 1, \dots, \\ 2(n - r) - 1, 2(n - r), 2(n - r) + 1, \dots, \\ r - 1, r - 1, \dots, 1). \quad (46)$$

Therefore, in this case, $P(n, d, r)$, it is not parallel unimodal and it is not parallel symmetric. This is the only case in which the fundamental sequence is not unimodal (see Example 11).

Case VIII ($2(n - r) > d \geq 2r$ and $d \geq \max\{r, n - r\}$). In this case, the parallel rank is $T_{\text{par}}(\widehat{0}) = 2n - d - 2$. The fundamental sequence depends on the relations between r , $n - r$, and d in the following way.

(VIII.1) If $d \leq 2n - 3r + 1$, the fundamental sequence is

$$(2, 4, \dots, 2(d - n + r), 2(d - n + r) + 1, \dots, d - n + 2r - 1, \\ \dots, d - n + 2r - 1, d - n + 2r, \dots, n - r, \dots, n - r, \\ n - r - 1, \dots, 2n - 2r - d, 2n - 2r - d - 2, \dots, d - 2r, \\ d - 2r - 1, \dots, 1). \quad (47)$$

(VIII.2) If $d > 2n - 3r + 1$ and $2d - 1 \geq 3(n - r)$, the fundamental sequence is

$$(2, 4, \dots, 2(d - n + r), 2(d - n + r) + 1, \dots, d - n + 2r - 1, \\ \dots, d - n + 2r - 1, d - n + 2r - 2, \dots, 2(d - n + r), \\ 2(d - n + r - 1), \dots, 2(2n - 2r - d), \\ 2(2n - 2r - d) - 1, \dots, 2n - 2r - d, \\ 2n - 2r - d - 2, \dots, d - 2r, d - 2r - 1, \dots, 1). \quad (48)$$


(VIII.3) If $d > 2n - 3r + 1$ and $2d - 1 > 3(n - r)$, the fundamental sequence is

$$\begin{aligned}
 &(2, 4, \dots, 2(d - n + r), 2(d - n + r) + 1, \dots, d - n + 2r - 1, \\
 &\dots, d - n + 2r - 1, d - n + 2r - 2, \dots, n - r + 1, \\
 &n - r, \dots, n - r, n - r - 1, \dots, 2n - 2r - d, \\
 &2n - 2r - d - 2, \dots, d - 2r, d - 2r - 1, \dots, 1).
 \end{aligned}
 \tag{49}$$

In these cases $P(n, d, r)$ is parallel unimodal but not parallel symmetric. \square

We summarize all the previous results in Tables 1 and 2.

Example 11. Let us consider the lattice $P(n, d, r)$ with $n = 7$, $d = 6$, and $r = 5$. In this case, $2r > d \geq 2(n - r)$, $d \geq \max\{r, n - r\}$, and $3r > 2(n - r)$. The fundamental chain is the following:



5. Conclusions and Future Research Directions

In this work, we have introduced a refinement of the concept of parallel time convergence for a finite deterministic discrete dynamical model X which also has a lattice structure. This new concept is the *fundamental sequence* of X . In particular, the length of the fundamental sequence is exactly the parallel time of convergence from the minimum to the maximum of X . In this paper, we have computed the fundamental sequence for two models whose configurations are signed integer partitions. The perspectives, for future research directions, are the following: to compute the fundamental sequence for more general models, to compare the properties of the fundamental sequence with the properties of other relevant combinatorial objects related to the order structure of X (e.g., the Whitney number sequence and the rank of X when X is graded), and to characterize the models whose fundamental sequence has some specific property (e.g., unimodality or symmetry).

References

- [1] É. Goles, M. Latapy, C. Magnien, M. Morvan, and H. D. Phan, "Sandpile models and lattices: a comprehensive survey," *Theoretical Computer Science*, vol. 322, no. 2, pp. 383–407, 2004.
- [2] T. Brylawski, "The lattice of integer partitions," *Discrete Mathematics*, vol. 6, pp. 201–219, 1973.
- [3] E. Goles, "Sand pile automata," *Annales de l'Institut Henri Poincaré A*, vol. 56, no. 1, pp. 75–90, 1992.
- [4] E. Goles and M. A. Kiwi, "Games on line graphs and sand piles," *Theoretical Computer Science*, vol. 115, no. 2, pp. 321–349, 1993.
- [5] P. Bak, C. Tang, and K. Wiesenfeld, "Self-organized criticality," *Physical Review A*, vol. 38, no. 1, pp. 364–374, 1988.
- [6] D. Dhar, "The abelian sandpiles and related models," *Physica A*, vol. 263, no. 1–4, pp. 4–25, 1999.
- [7] E. Goles and M. Margenstern, "Universality of the chip-firing game," *Theoretical Computer Science*, vol. 172, no. 1-2, pp. 121–134, 1997.
- [8] E. Goles, M. Morvan, and H. D. Phan, "Lattice structure and convergence of a game of cards," *Annals of Combinatorics*, vol. 6, no. 3-4, pp. 327–335, 2002.
- [9] G. Cattaneo, M. Comito, and D. Bianucci, "Sand piles: from physics to cellular automata models," *Theoretical Computer Science*, vol. 436, pp. 35–53, 2012.
- [10] A. Dennunzio, P. Guillon, and B. Masson, "Sand automata as cellular automata," *Theoretical Computer Science*, vol. 410, no. 38–40, pp. 3962–3974, 2009.
- [11] E. Goles and M. A. Kiwi, "One-dimensional sand piles, cellular automata and related models," in *Nonlinear Phenomena in Fluids, Solids and Other Complex Systems (Santiago, 1990)*, North-Holland Delta Series, pp. 169–185, North-Holland, Amsterdam, The Netherlands, 1991.
- [12] J. Cervelle and E. Formenti, "On sand automata," in *Mathematical Foundations of Computer Science 2003*, vol. 2607 of *Lecture Notes in Computer Science*, pp. 642–653, Springer, Berlin, Germany, 2003.
- [13] J. Cervelle, E. Formenti, and B. Masson, "Basic properties for sand automata," in *Mathematical Foundations of Computer Science 2005*, vol. 3618 of *Lecture Notes in Computer Science*, pp. 192–211, Springer, Berlin, Germany, 2005.
- [14] J. Cervelle, E. Formenti, and B. Masson, "From sandpiles to sand automata," *Theoretical Computer Science*, vol. 381, no. 1-3, pp. 1–28, 2007.
- [15] E. Formenti and B. Masson, "On computing the fixed points for generalized sandpiles," *International Journal of Unconventional Computing*, vol. 2, no. 1, pp. 13–25, 2005.
- [16] E. Formenti and B. Masson, "A note on fixed points of generalized ice piles models," *International Journal of Unconventional Computing*, vol. 2, no. 2, pp. 183–191, 2006.
- [17] E. Formenti, B. Masson, and T. Pisokas, "On symmetric sandpiles," in *Cellular Automata*, vol. 4173 of *Lecture Notes in Computer Science*, pp. 676–685, Springer, Berlin, Germany, 2006.
- [18] E. Formenti, B. Masson, and T. Pisokas, "Advances in symmetric sandpiles," *Fundamenta Informaticae*, vol. 76, no. 1-2, pp. 91–112, 2007.

- [19] E. Goles, M. Morvan, and H. D. Phan, "Sandpiles and order structure of integer partitions," *Discrete Applied Mathematics*, vol. 117, no. 1–3, pp. 51–64, 2002.
- [20] M. H. Le and T. H. D. Phan, "Strict partitions and discrete dynamical systems," *Theoretical Computer Science*, vol. 389, no. 1–2, pp. 82–90, 2007.
- [21] M. Latapy, "Partitions of an integer into powers," in *Discrete Models: Combinatorics, Computation, and Geometry (Paris, 2001)*, Discrete Mathematics and Theoretical Computer Science, pp. 215–228, 2001.
- [22] M. Latapy, "Integer partitions, tilings of 2D-gons and lattices," *RAIRO—Theoretical Informatics and Applications*, vol. 36, no. 4, pp. 389–399, 2002.
- [23] M. Latapy, R. Mantaci, M. Morvan, and H. D. Phan, "Structure of some sand piles model," *Theoretical Computer Science*, vol. 262, no. 1–2, pp. 525–556, 2001.
- [24] M. Latapy and H. D. Phan, "The lattice structure of chip firing games and related models," *Physica D*, vol. 155, no. 1–2, pp. 69–82, 2001.
- [25] M. Latapy and T. H. D. Phan, "The lattice of integer partitions and its infinite extension," *Discrete Mathematics*, vol. 309, no. 6, pp. 1357–1367, 2009.
- [26] K. Perrot and E. Rémila, "Transduction on Kadanoff sand pile model avalanches, application to wave pattern emergence," in *Mathematical Foundations of Computer Science 2011*, vol. 6907 of *Lecture Notes in Comput. Sci.*, pp. 508–519, Springer, Heidelberg, Germany, 2011.
- [27] K. Perrot, T. H. D. Phan, and T. van Pham, "On the set of fixed points of the parallel symmetric sand pile model," in *Automata 2011—17th International Workshop on Cellular Automata and Discrete Complex Systems*, Discrete Mathematics and Theoretical Computer Science, pp. 17–28, Discrete Mathematics and Theoretical Computer Science Proceedings, 2011.
- [28] T. H. D. Phan, "Two sided sand piles model and unimodal sequences," *RAIRO—Theoretical Informatics and Applications*, vol. 42, no. 3, pp. 631–646, 2008.
- [29] P. T. H. Duong and T. T. T. Huong, "On the stability of sand piles model," *Theoretical Computer Science*, vol. 411, no. 3, pp. 594–601, 2010.
- [30] G. E. Andrews, "Euler's 'de Partitio numerorum,'" *Bulletin of the American Mathematical Society*, vol. 44, no. 4, pp. 561–573, 2007.
- [31] W. J. Keith, "A bijective toolkit for signed partitions," *Annals of Combinatorics*, vol. 15, no. 1, pp. 95–117, 2011.
- [32] G. Chiaselotti, "On a problem concerning the weight functions," *European Journal of Combinatorics*, vol. 23, no. 1, pp. 15–22, 2002.
- [33] G. Chiaselotti, G. Infante, and G. Marino, "New results related to a conjecture of Manickam and Singhi," *European Journal of Combinatorics*, vol. 29, no. 2, pp. 361–368, 2008.
- [34] G. Chiaselotti, G. Marino, and C. Nardi, "A minimum problem for finite sets of real numbers with nonnegative sum," *Journal of Applied Mathematics*, vol. 2012, Article ID 847958, 15 pages, 2012.
- [35] G. Marino and G. Chiaselotti, "A method to count the positive 3-subsets in a set of real numbers with non-negative sum," *European Journal of Combinatorics*, vol. 23, no. 5, pp. 619–629, 2002.
- [36] C. Bisi, G. Chiaselotti, and P. A. Oliverio, "Sand piles models of signed partitions with d piles," *ISRN Combinatorics*, vol. 2013, Article ID 615703, 7 pages, 2013.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

