

Research Article

Nonlinear Integral Inequalities in Two Independent Variables and Their Applications

Kelong Zheng, Yu Wu, and Shengfu Deng

Received 10 June 2007; Accepted 27 July 2007

Recommended by Wing-Sum Cheung

This paper generalizes results of Cheung and Ma (2005) to more general inequalities with more than one distinct nonlinear term. From our results, some results of Cheung and Ma (2005) can be deduced as some special cases. Our results are also applied to show the boundedness of the solutions of a partial differential equation.

Copyright © 2007 Kelong Zheng et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

The integral inequalities play a fundamental role in the study of existence, uniqueness, boundedness, stability, invariant manifolds, and other qualitative properties of solutions of the theory of differential and integral equations. There are a lot of papers investigating them such as [1–8]. In particular, Pachpatte [2] discovered some new integral inequalities involving functions of two variables. These inequalities are applied to study the boundedness and uniqueness of the solutions of the following terminal value problem for the hyperbolic partial differential equation (1.1) with conditions (1.2):

$$D_1 D_2 u(x, y) = h(x, y, u(x, y)) + r(x, y), \quad (1.1)$$

$$u(x, \infty) = \sigma_\infty(x), \quad u(\infty, y) = \tau_\infty(y), \quad u(\infty, \infty) = k. \quad (1.2)$$

Cheung [9], and Dragomir and Kim [10, 11] established additional Gronwall-Ou-Iang type integral inequalities involving functions of two independent variables. Meng and Li [12] generalized the results of Pachpatte [2] to certain new integrals. Recently, Cheung

and Ma[13] discussed the following inequalities

$$\begin{aligned}
 u(x, y) &\leq a(x, y) + c(x, y) \int_0^x \int_y^\infty d(s, t)w(u(s, t)) dt ds, \\
 u(x, y) &\leq a(x, y) + c(x, y) \int_x^\infty \int_y^\infty d(s, t)w(u(s, t)) dt ds,
 \end{aligned}
 \tag{1.3}$$

where $a(x, y)$ and $c(x, y)$ have certain monotonicity.

Our main aim here, motivated by the work of Cheung and Ma [13], is to discuss more general integral inequalities with n nonlinear terms:

$$u(x, y) \leq a(x, y) + \sum_{i=1}^n \int_0^x \int_y^\infty d_i(x, y, s, t)w_i(u(s, t)) dt ds,
 \tag{1.4}$$

$$u(x, y) \leq a(x, y) + \sum_{i=1}^n \int_x^\infty \int_y^\infty d_i(x, y, s, t)w_i(u(s, t)) dt ds,
 \tag{1.5}$$

where we do not require the monotonicity of $a(x, y)$ and $d_i(x, y, s, t)$. Furthermore, we also show that some results of Cheung and Ma [13] can be deduced from our results as some special cases. Our results are also applied to show the boundedness of the solutions of a partial differential equation.

2. Main results

Let $\mathbb{R} = (-\infty, \infty)$ and $\mathbb{R}_+ = [0, \infty)$. $D_1z(x, y)$ and $D_2z(x, y)$ denote the first-order partial derivatives of $z(x, y)$ with respect to x and y , respectively.

As in [1, 5, 6], we define $w_1 \propto w_2$ for $w_1, w_2 : A \subset \mathbb{R} \rightarrow \mathbb{R} \setminus \{0\}$ if w_2/w_1 is nondecreasing on A . This concept helps us compare monotonicity of different functions. Suppose that

(C₁) $w_i(u)$ ($i = 1, \dots, n$) is a nonnegative, nondecreasing, and continuous function for $u \in \mathbb{R}_+$ with $w_i(u) > 0$ for $u > 0$ such that $w_1 \propto w_2 \propto \dots \propto w_n$;

(C₂) $a(x, y)$ is a nonnegative and continuous function for $x, y \in \mathbb{R}_+$;

(C₃) $d_i(x, y, s, t)$ ($i = 1, \dots, n$) is a continuous and nonnegative function for $x, y, s, t \in \mathbb{R}_+$.

Take the notation $W_i(u) := \int_{u_i}^u (dz/w_i(z))$, for $u \geq u_i$, where $u_i > 0$ is a given constant. Clearly, W_i is strictly increasing, so its inverse W_i^{-1} is well defined, continuous, and increasing in its corresponding domain.

THEOREM 2.1. *In addition to the assumptions (C₁), (C₂), and (C₃), suppose that $a(x, y)$ and $d_i(x, y, s, t)$ are bounded in $y \in \mathbb{R}_+$ for each fixed $x, s, t \in \mathbb{R}_+$. If $u(x, y)$ is a continuous and nonnegative function satisfying (1.4) for $x, y \in \mathbb{R}_+$, then*

$$u(x, y) \leq W_n^{-1} \left[W_n(b_n(x, y)) + \int_0^x \int_y^\infty \tilde{d}_n(x, y, s, t) dt ds \right]
 \tag{2.1}$$

for all $0 \leq x \leq x_1$, $y_1 \leq y < \infty$, where $b_n(x, y)$ is determined recursively by

$$\begin{aligned} b_1(x, y) &= \tilde{a}(x, y), \\ b_{i+1}(x, y) &= W_i^{-1} \left[W_i(b_i(x, y)) + \int_0^x \int_{y_1}^y \tilde{d}_i(x, y, s, t) dt ds \right], \\ \tilde{a}(x, y) &= \sup_{0 \leq \tau \leq x} \sup_{y \leq \mu < \infty} a(\tau, \mu), \quad \tilde{d}_i(x, y, s, t) = \sup_{0 \leq \tau \leq x} \sup_{y \leq \mu < \infty} d_i(\tau, \mu, s, t), \end{aligned} \quad (2.2)$$

$W_1(0) := 0$, and $x_1, y_1 \in \mathbb{R}_+$ are chosen such that

$$W_i(b_i(x_1, y_1)) + \int_0^{x_1} \int_{y_1}^y \tilde{d}_i(x, y, s, t) dt ds \leq \int_{u_i}^{\infty} \frac{dz}{w_i(z)} \quad (2.3)$$

for $i = 1, \dots, n$.

Remark 2.2. x_1 and y_1 are confined by (2.3). In particular, (2.1) is true for all $x, y \in \mathbb{R}_+$ when all w_i ($i = 1, \dots, n$) satisfy $\int_{u_i}^{\infty} (dz/w_i(z)) = \infty$.

Remark 2.3. As in [6, 5, 1], different choices of u_i in W_i do not affect our results.

Proof of Theorem 2.1. From the assumptions, we know that $\tilde{a}(x, y)$ and $\tilde{d}_i(x, y, s, t)$ are well defined. Moreover, $\tilde{a}(x, y)$ and $\tilde{d}_i(x, y, s, t)$ are nonnegative, nondecreasing in x , non-increasing in y ; and satisfy $\tilde{a}(x, y) \geq a(x, y)$ and $\tilde{d}_i(x, y, s, t) \geq d_i(x, y, s, t)$ for each $i = 1, \dots, n$.

We first discuss the case that $a(x, y) > 0$ for all $x, y \in \mathbb{R}_+$. Thus, $b_1(x, y)$ is positive, nondecreasing in x , nonincreasing in y ; and satisfies $b_1(x, y) \geq a(x, y)$ for all $x, y \in \mathbb{R}_+$. From (1.4), we have

$$u(x, y) \leq b_1(x, y) + \sum_{i=1}^n \int_0^x \int_y^{\infty} \tilde{d}_i(x, y, s, t) w_i(u(s, t)) dt ds. \quad (2.4)$$

Choose arbitrary \tilde{x}_1, \tilde{y}_1 such that $0 \leq \tilde{x}_1 \leq x_1$, $y_1 \leq \tilde{y}_1 < \infty$. From (2.4), we obtain

$$u(x, y) \leq b_1(\tilde{x}_1, \tilde{y}_1) + \sum_{i=1}^n \int_0^x \int_y^{\infty} \tilde{d}_i(\tilde{x}_1, \tilde{y}_1, s, t) w_i(u(s, t)) dt ds \quad (2.5)$$

for all $0 \leq x \leq \tilde{x}_1 \leq x_1$, $y_1 \leq \tilde{y}_1 \leq y < \infty$.

Having (2.5), we claim

$$u(x, y) \leq W_n^{-1} \left[W_n(\tilde{b}_n(\tilde{x}_1, \tilde{y}_1, x, y)) + \int_0^x \int_y^{\infty} \tilde{d}_n(\tilde{x}_1, \tilde{y}_1, s, t) dt ds \right] \quad (2.6)$$

for all $0 \leq x \leq \min\{\tilde{x}_1, x_2\}$, $\max\{\tilde{y}_1, y_2\} \leq y < \infty$, where

$$\begin{aligned} \tilde{b}_1(\tilde{x}_1, \tilde{y}_1, x, y) &= b_1(\tilde{x}_1, \tilde{y}_1), \\ \tilde{b}_{i+1}(\tilde{x}_1, \tilde{y}_1, x, y) &= W_i^{-1} \left[W_i(\tilde{b}_i(\tilde{x}_1, \tilde{y}_1, x, y)) + \int_0^x \int_y^{\infty} \tilde{d}_i(\tilde{x}_1, \tilde{y}_1, s, t) dt ds \right] \end{aligned} \quad (2.7)$$

for $i = 1, \dots, n - 1$ and $x_2, y_2 \in \mathbb{R}_+$ are chosen such that

$$W_i(\tilde{b}_i(\tilde{x}_1, \tilde{y}_1, x_2, y_2)) + \int_0^{x_2} \int_{y_2}^{\infty} \tilde{d}_i(\tilde{x}_1, \tilde{y}_1, s, t) dt ds \leq \int_{u_i}^{\infty} \frac{dz}{w_i(z)} \tag{2.8}$$

for $i = 1, \dots, n$.

Note that we may take $x_2 = x_1$ and $y_2 = y_1$. In fact, $\tilde{b}_i(\tilde{x}_1, \tilde{y}_1, x, y)$ and $\tilde{d}_i(\tilde{x}_1, \tilde{y}_1, x, y)$ are nondecreasing in \tilde{x}_1 , nonincreasing in \tilde{y}_1 for fixed x, y . Furthermore, it is easy to check that $\tilde{b}_i(\tilde{x}_1, \tilde{y}_1, \tilde{x}_1, \tilde{y}_1) = b_i(\tilde{x}_1, \tilde{y}_1)$ for $i = 1, \dots, n$. If x_2, y_2 are replaced by x_1, y_1 on the left side of (2.8), we have from (2.3)

$$\begin{aligned} W_i(\tilde{b}_i(\tilde{x}_1, \tilde{y}_1, x_1, y_1)) + \int_0^{x_1} \int_{y_1}^{\infty} \tilde{d}_i(\tilde{x}_1, \tilde{y}_1, s, t) dt ds \\ \leq W_i(\tilde{b}_i(x_1, y_1, x_1, y_1)) + \int_0^{x_1} \int_{y_1}^{\infty} \tilde{d}_i(x_1, y_1, s, t) dt ds \\ = W_i(b_i(x_1, y_1)) + \int_0^{x_1} \int_{y_1}^{\infty} \tilde{d}_i(x_1, y_1, s, t) dt ds \leq \int_{u_i}^{\infty} \frac{dz}{w_i(z)}. \end{aligned} \tag{2.9}$$

Thus, it means that we can take $x_2 = x_1, y_2 = y_1$.

In the following, we will use mathematical induction to prove (2.6).

For $n = 1$, let

$$z(x, y) = \int_0^x \int_y^{\infty} \tilde{d}_1(\tilde{x}_1, \tilde{y}_1, s, t) w_1(u(s, t)) dt ds. \tag{2.10}$$

Then $z(x, y)$ is differentiable, nonnegative, nondecreasing for $x \in [0, \tilde{x}_1]$, and nonincreasing for $y \in [\tilde{y}_1, \infty)$ and $z(0, y) = z(x, \infty) = 0$. From (2.5), we have the following:

$$\begin{aligned} u(x, y) &\leq b_1(\tilde{x}_1, \tilde{y}_1) + z(x, y), \\ D_1 z(x, y) &= \int_y^{\infty} \tilde{d}_1(\tilde{x}_1, \tilde{y}_1, x, t) w_1(u(x, t)) dt \\ &\leq \int_y^{\infty} \tilde{d}_1(\tilde{x}_1, \tilde{y}_1, x, t) w_1(b_1(\tilde{x}_1, \tilde{y}_1) + z(x, t)) dt \\ &\leq w_1(b_1(\tilde{x}_1, \tilde{y}_1) + z(x, y)) \int_y^{\infty} \tilde{d}_1(\tilde{x}_1, \tilde{y}_1, x, t) dt. \end{aligned} \tag{2.11}$$

Since w_1 is nondecreasing and $b_1(\tilde{x}_1, \tilde{y}_1) + z(x, y) > 0$, we get

$$\begin{aligned} \frac{D_1(b_1(\tilde{x}_1, \tilde{y}_1) + z(x, y))}{w_1(b_1(\tilde{x}_1, \tilde{y}_1) + z(x, y))} &= \frac{D_1 z(x, y)}{w_1(b_1(\tilde{x}_1, \tilde{y}_1) + z(x, y))} \\ &\leq \frac{w_1(b_1(\tilde{x}_1, \tilde{y}_1) + z(x, y)) \int_y^{\infty} \tilde{d}_1(\tilde{x}_1, \tilde{y}_1, x, t) dt}{w_1(b_1(\tilde{x}_1, \tilde{y}_1) + z(x, y))} \\ &= \int_y^{\infty} \tilde{d}_1(\tilde{x}_1, \tilde{y}_1, x, t) dt. \end{aligned} \tag{2.12}$$

Integrating both sides of the above inequality from 0 to x , we obtain

$$\begin{aligned} W_1(b_1(\tilde{x}_1, \tilde{y}_1) + z(x, y)) &\leq W_1(b_1(\tilde{x}_1, \tilde{y}_1) + z(0, y)) + \int_0^x \int_y^\infty \tilde{d}_1(\tilde{x}_1, \tilde{y}_1, s, t) dt ds \\ &= W_1(b_1(\tilde{x}_1, \tilde{y}_1)) + \int_0^x \int_y^\infty \tilde{d}_1(\tilde{x}_1, \tilde{y}_1, s, t) dt ds. \end{aligned} \quad (2.13)$$

Thus the monotonicity of W_1^{-1} implies

$$u(x, y) \leq b_1(\tilde{x}_1, \tilde{y}_1) + z(x, y) \leq W_1^{-1} \left[W_1(b_1(\tilde{x}_1, \tilde{y}_1)) + \int_0^x \int_y^\infty \tilde{d}_1(\tilde{x}_1, \tilde{y}_1, s, t) dt ds \right], \quad (2.14)$$

that is, (2.6) is true for $n = 1$.

Assume that (2.6) is true for $n = m$. Consider

$$u(x, y) \leq b_1(\tilde{x}_1, \tilde{y}_1) + \sum_{i=1}^{m+1} \int_0^x \int_y^\infty \tilde{d}_i(\tilde{x}_1, \tilde{y}_1, s, t) w_i(u(s, t)) dt ds \quad (2.15)$$

for all $0 \leq x \leq \tilde{x}_1, \tilde{y}_1 \leq y < \infty$. Let

$$z(x, y) = \sum_{i=1}^{m+1} \int_0^x \int_y^\infty \tilde{d}_i(\tilde{x}_1, \tilde{y}_1, s, t) w_i(u(s, t)) dt ds. \quad (2.16)$$

Then $z(x, y)$ is differentiable, nonnegative, nondecreasing for $x \in [0, \tilde{x}_1]$, and nonincreasing for $y \in [\tilde{y}_1, \infty)$. Obviously, $z(0, y) = z(x, \infty) = 0$ and $u(x, y) \leq b_1(\tilde{x}_1, \tilde{y}_1) + z(x, y)$. Since w_1 is nondecreasing and $b_1(\tilde{x}_1, \tilde{y}_1) + z(x, y) > 0$, we have

$$\begin{aligned} &\frac{D_1(b_1(\tilde{x}_1, \tilde{y}_1) + z(x, y))}{w_1(b_1(\tilde{x}_1, \tilde{y}_1) + z(x, y))} \\ &\leq \frac{\sum_{i=1}^{m+1} \int_y^\infty \tilde{d}_i(\tilde{x}_1, \tilde{y}_1, x, t) w_i(u(x, t)) dt}{w_1(b_1(\tilde{x}_1, \tilde{y}_1) + z(x, y))} \\ &\leq \frac{\sum_{i=1}^{m+1} \int_y^\infty \tilde{d}_i(\tilde{x}_1, \tilde{y}_1, x, t) w_i(b_1(\tilde{x}_1, \tilde{y}_1) + z(x, t)) dt}{w_1(b_1(\tilde{x}_1, \tilde{y}_1) + z(x, y))} \\ &\leq \int_y^\infty \tilde{d}_1(\tilde{x}_1, \tilde{y}_1, x, t) dt + \sum_{i=2}^{m+1} \int_y^\infty \tilde{d}_i(\tilde{x}_1, \tilde{y}_1, x, t) \phi_i(b_1(\tilde{x}_1, \tilde{y}_1) + z(x, t)) dt \\ &\leq \int_y^\infty \tilde{d}_1(\tilde{x}_1, \tilde{y}_1, x, t) dt + \sum_{i=1}^m \int_y^\infty \tilde{d}_{i+1}(\tilde{x}_1, \tilde{y}_1, x, t) \phi_{i+1}(b_1(\tilde{x}_1, \tilde{y}_1) + z(x, t)) dt, \end{aligned} \quad (2.17)$$

where $\phi_{i+1}(u) = w_{i+1}(u)/w_1(u)$, $i = 1, \dots, m$. Integrating the above inequality from 0 to x , we obtain

$$W_1(b_1(\tilde{x}_1, \tilde{y}_1) + z(x, y)) \leq W_1(b_1(\tilde{x}_1, \tilde{y}_1)) + \int_0^x \int_y^\infty \tilde{d}_1(\tilde{x}_1, \tilde{y}_1, s, t) dt ds + \sum_{i=1}^m \int_0^x \int_y^\infty \tilde{d}_{i+1}(\tilde{x}_1, \tilde{y}_1, s, t) \phi_{i+1}(b_1(\tilde{x}_1, \tilde{y}_1) + z(s, t)) dt ds, \tag{2.18}$$

or

$$\xi(x, y) \leq c_1(x, y) + \sum_{i=1}^m \int_0^x \int_y^\infty \tilde{d}_{i+1}(\tilde{x}_1, \tilde{y}_1, s, t) \phi_{i+1}(W_1^{-1}(\xi(s, t))) dt ds \tag{2.19}$$

for $0 \leq x \leq \tilde{x}_1$ and $\tilde{y}_1 \leq y < \infty$, the same as (2.6) for $n = m$, where $\xi(x, y) = W_1(b_1(\tilde{x}_1, \tilde{y}_1) + z(x, y))$ and $c_1(x, y) = W_1(b_1(\tilde{x}_1, \tilde{y}_1)) + \int_0^x \int_y^\infty \tilde{d}_1(\tilde{x}_1, \tilde{y}_1, s, t) dt ds$.

From the assumption (C_1) , each $\phi_{i+1}(W_1^{-1}(u))$, $i = 1, \dots, m$, is continuous and non-decreasing for u . Moreover, $\phi_2(W_1^{-1}) \circ \phi_3(W_1^{-1}) \circ \dots \circ \phi_{m+1}(W_1^{-1})$. By the inductive assumption, we have

$$\xi(x, y) \leq \Phi_{m+1}^{-1} \left[\Phi_{m+1}(c_m(x, y)) + \int_0^x \int_y^\infty \tilde{d}_{m+1}(\tilde{x}_1, \tilde{y}_1, s, t) dt ds \right] \tag{2.20}$$

for all $0 \leq x \leq \min\{\tilde{x}_1, x_3\}$, $\max\{\tilde{y}_1, y_3\} \leq y < \infty$, where $\Phi_{i+1}(u) = \int_{\tilde{u}_{i+1}}^u (dz/\phi_{i+1}(W_1^{-1}(z)))$, $u > 0, \tilde{u}_{i+1} = W_1(u_{i+1})$, Φ_{i+1}^{-1} is the inverse of Φ_{i+1} , $i = 1, \dots, m$,

$$c_{i+1}(x, y) = \Phi_{i+1}^{-1} \left[\Phi_{i+1}(c_i(x, y)) + \int_0^x \int_y^\infty \tilde{d}_{i+1}(\tilde{x}_1, \tilde{y}_1, s, t) dt ds \right], \quad i = 1, \dots, m, \tag{2.21}$$

and $x_3, y_3 \in \mathbb{R}_+$ are chosen such that

$$\Phi_{i+1}(c_i(x_3, y_3)) + \int_0^{x_3} \int_{y_3}^\infty \tilde{d}_{i+1}(\tilde{x}_1, \tilde{y}_1, s, t) dt ds \leq \int_{\tilde{u}_{i+1}}^{W_1(\infty)} \frac{dz}{\phi_{i+1}(W_1^{-1}(z))} \tag{2.22}$$

for $i = 1, \dots, m$.

Note that

$$\begin{aligned} \Phi_i(u) &= \int_{\tilde{u}_i}^u \frac{dz}{\phi_i(W_1^{-1}(z))} = \int_{W_1(u_i)}^u \frac{w_1(W_1^{-1}(z)) dz}{w_i(W_1^{-1}(z))} \\ &= \int_{u_i}^{W_1^{-1}(u)} \frac{dz}{w_i(z)} = W_i \circ W_1^{-1}(u), \quad i = 2, \dots, m+1. \end{aligned} \tag{2.23}$$

From (2.20), we have

$$\begin{aligned} u(x, y) &\leq b_1(\tilde{x}_1, \tilde{y}_1) + z(x, y) = W_1^{-1}(\xi(x, y)) \\ &\leq W_{m+1}^{-1} \left[W_{m+1}(W_1^{-1}(c_m(x, y))) + \int_0^x \int_y^\infty \tilde{d}_{m+1}(\tilde{x}_1, \tilde{y}_1, s, t) dt ds \right] \end{aligned} \tag{2.24}$$

for all $0 \leq x \leq \min\{\tilde{x}_1, x_3\}$, $\max\{\tilde{y}_1, y_3\} \leq y < \infty$. Let $\tilde{c}_i(x, y) = W_1^{-1}(c_i(x, y))$. Then,

$$\begin{aligned}\tilde{c}_1(x, y) &= W_1^{-1}(c_1(x, y)) \\ &= W_1^{-1}\left[W_1(b_1(\tilde{x}_1, \tilde{y}_1)) + \int_0^x \int_y^\infty \tilde{d}_1(\tilde{x}_1, \tilde{y}_1, s, t) dt ds\right] \\ &= \tilde{b}_2(\tilde{x}_1, \tilde{y}_1, x, y).\end{aligned}\quad (2.25)$$

Moreover, with the assumption that $\tilde{c}_m(x, y) = \tilde{b}_{m+1}(\tilde{x}_1, \tilde{y}_1, x, y)$, we have

$$\begin{aligned}\tilde{c}_{m+1}(x, y) &= W_1^{-1}\left[\Phi_{m+1}^{-1}(\Phi_{m+1}(c_m(x, y))) + \int_0^x \int_y^\infty \tilde{d}_{m+1}(\tilde{x}_1, \tilde{y}_1, s, t) dt ds\right] \\ &= W_{m+1}^{-1}\left[W_{m+1}(W_1^{-1}(c_m(x, y))) + \int_0^x \int_y^\infty \tilde{d}_{m+1}(\tilde{x}_1, \tilde{y}_1, s, t) dt ds\right] \\ &= W_{m+1}^{-1}\left[W_{m+1}(\tilde{c}_m(x, y)) + \int_0^x \int_y^\infty \tilde{d}_{m+1}(\tilde{x}_1, \tilde{y}_1, s, t) dt ds\right] \\ &= W_{m+1}^{-1}\left[W_{m+1}(\tilde{b}_{m+1}(\tilde{x}_1, \tilde{y}_1, x, y)) + \int_0^x \int_y^\infty \tilde{d}_{m+1}(\tilde{x}_1, \tilde{y}_1, s, t) dt ds\right] \\ &= \tilde{b}_{m+2}(\tilde{x}_1, \tilde{y}_1, x, y).\end{aligned}\quad (2.26)$$

This proves that

$$\tilde{c}_i(x, y) = \tilde{b}_{i+1}(\tilde{x}_1, \tilde{y}_1, x, y), \quad i = 1, \dots, m. \quad (2.27)$$

Therefore, (2.22) becomes

$$\begin{aligned}W_{i+1}(\tilde{b}_{i+1}(\tilde{x}_1, \tilde{y}_1, x_3, y_3)) + \int_0^{x_3} \int_{y_3}^\infty \tilde{d}_{i+1}(\tilde{x}_1, \tilde{y}_1, s, t) dt ds \\ \leq \int_{\tilde{u}_{i+1}}^{W_1(\infty)} \frac{dz}{\phi_{i+1}(W_1^{-1}(z))} = \int_{u_{i+1}}^\infty \frac{dz}{w_{i+1}(z)}, \quad i = 1, \dots, m.\end{aligned}\quad (2.28)$$

The above inequalities and (2.8) imply that we may take $x_2 = x_3$, $y_2 = y_3$. From (2.24), we get

$$u(x, y) \leq W_{m+1}^{-1}\left[W_{m+1}(\tilde{b}_{m+1}(\tilde{x}_1, \tilde{y}_1, x, y)) + \int_0^x \int_y^\infty \tilde{d}_{m+1}(\tilde{x}_1, \tilde{y}_1, s, t) dt ds\right] \quad (2.29)$$

for all $0 \leq x \leq \tilde{x}_1 \leq x_2$, $y_2 \leq \tilde{y}_1 \leq y < \infty$. This proves (2.6) by mathematical induction.

Taking $x = \tilde{x}_1$, $y = \tilde{y}_1$, $x_2 = x_1$, and $y_2 = y_1$, we have

$$u(\tilde{x}_1, \tilde{y}_1) \leq W_n^{-1}\left[W_n(\tilde{b}_n(\tilde{x}_1, \tilde{y}_1, \tilde{x}_1, \tilde{y}_1)) + \int_0^{\tilde{x}_1} \int_{\tilde{y}_1}^\infty \tilde{d}_n(\tilde{x}_1, \tilde{y}_1, s, t) dt ds\right] \quad (2.30)$$

for $0 \leq \tilde{x}_1 \leq x_1, y_1 \leq \tilde{y}_1 < \infty$. It is easy to verify $\tilde{b}_n(\tilde{x}_1, \tilde{y}_1, \tilde{x}_1, \tilde{y}_1) = b_n(\tilde{x}_1, \tilde{y}_1)$. Thus, (2.30) can be written as

$$u(\tilde{x}_1, \tilde{y}_1) \leq W_n^{-1} \left[W_n(b_n(\tilde{x}_1, \tilde{y}_1)) + \int_0^{\tilde{x}_1} \int_{\tilde{y}_1}^{\infty} \tilde{d}_n(\tilde{x}_1, \tilde{y}_1, s, t) dt ds \right]. \tag{2.31}$$

Since \tilde{x}_1, \tilde{y}_1 are arbitrary, replace \tilde{x}_1 and \tilde{y}_1 by x and y respectively and we have

$$u(x, y) \leq W_n^{-1} \left[W_n(b_n(x, y)) + \int_0^x \int_y^{\infty} \tilde{d}_n(x, y, s, t) dt ds \right] \tag{2.32}$$

for all $0 \leq x \leq x_1, y_1 \leq y < \infty$.

In case $a(x, y) = 0$ for some $x, y \in \mathbb{R}_+$. Let $b_{1,\epsilon}(x, y) := b_1(x, y) + \epsilon$ for all $x, y \in \mathbb{R}_+$, where $\epsilon > 0$ is arbitrary, and then $b_{1,\epsilon}(x, y) > 0$. Using the same arguments as above, where $b_1(x, y)$ is replaced with $b_{1,\epsilon}(x, y) > 0$, we get

$$u(x, y) \leq W_n^{-1} \left[W_n(b_{n,\epsilon}(x, y)) + \int_0^x \int_y^{\infty} \tilde{d}_n(x, y, s, t) dt ds \right]. \tag{2.33}$$

Letting $\epsilon \rightarrow 0^+$, we obtain (2.1) by the continuity of $b_{1,\epsilon}$ in ϵ and the continuity of W_i and W_i^{-1} under the notation $W_1(0) := 0$. □

THEOREM 2.4. *In addition to the assumptions (C_1) , (C_2) , and (C_3) , suppose that $a(x, y)$ and $d_i(x, y, s, t)$ are bounded in $x, y \in \mathbb{R}_+$ for each fixed $s, t \in \mathbb{R}_+$. If $u(x, y)$ is a continuous and nonnegative function satisfying (1.5) for $x, y \in \mathbb{R}_+$, then*

$$u(x, y) \leq W_n^{-1} \left[W_n(b_n(x, y)) + \int_x^{\infty} \int_y^{\infty} \hat{d}_n(x, y, s, t) dt ds \right] \tag{2.34}$$

for all $x_4 \leq x < \infty, y_4 \leq y < \infty$, where $b_n(x, y)$ is determined recursively by

$$\begin{aligned} b_1(x, y) &= \hat{a}(x, y), \\ b_{i+1}(x, y) &= W_i^{-1} \left[W_i(b_i(x, y)) + \int_x^{\infty} \int_y^{\infty} \hat{d}_i(x, y, s, t) dt ds \right], \end{aligned} \tag{2.35}$$

$$\begin{aligned} \hat{a}(x, y) &= \sup_{x \leq \tau < \infty} \sup_{y \leq \mu < \infty} a(\tau, \mu), \\ \hat{d}_i(x, y, s, t) &= \sup_{x \leq \tau < \infty} \sup_{y \leq \mu < \infty} d_i(\tau, \mu, s, t), \end{aligned} \tag{2.36}$$

$W_1(0) := 0$, and $x_4, y_4 \in \mathbb{R}_+$ are chosen such that

$$W_i(b_i(x_4, y_4)) + \int_{x_4}^{\infty} \int_{y_4}^{\infty} \hat{d}_i(x, y, s, t) dt ds \leq \int_{u_i}^{\infty} \frac{dz}{w_i(z)} \tag{2.37}$$

for $i = 1, \dots, n$.

The proof is similar to the argument in the proof of Theorem 2.1 with suitable modification. We omit the details here.

Remark 2.5. Take $d_1(x, y, s, t) = c(x, y)d(s, t)$ and $n = 1$ in (1.4). Suppose that $a(x, y)$ and $c(x, y)$ are continuous, nonnegative, nondecreasing in x and nonincreasing in y ; and $d(s, t)$ is nonnegative and continuous. We note that

$$b_1(x, y) = a(x, y), \quad \tilde{d}_1(x, y, s, t) = c(x, y)d(s, t). \quad (2.38)$$

From Theorem 2.1, we get

$$u(x, y) \leq W_1^{-1} \left[W_1(a(x, y)) + c(x, y) \int_0^x \int_y^\infty d(s, t) dt ds \right], \quad (2.39)$$

which is exactly (2.6) of Lemma 2.2 in [13].

Remark 2.6. Take $d_1(x, y, s, t) = c(x, y)d(s, t)$ and $n = 1$ in (1.5). Suppose that $a(x, y)$ and $c(x, y)$ are continuous, nonnegative, nonincreasing in x, y ; and $d(s, t)$ is nonnegative and continuous. It is easy to check that

$$b_1(x, y) = a(x, y), \quad \hat{d}_1(x, y, s, t) = c(x, y)d(s, t). \quad (2.40)$$

From Theorem 2.4, we get

$$u(x, y) \leq W_1^{-1} [W_1(a(x, y)) + c(x, y) \int_x^\infty \int_y^\infty d(s, t) dt ds] \quad (2.41)$$

which is (2.10) of Lemma 2.2 in [13].

3. Applications

Consider the partial differential equation

$$D_1 D_2 v(x, y) = \frac{1}{(x+1)^2(y+1)^2} + \exp(-x) \exp(-y) \sqrt{|v(x, y)| + 1} + x \exp(-x) \exp(-y) \mathfrak{T}v(x, y), \quad (3.1)$$

$$v(x, \infty) = \sigma(x), v(0, y) = \tau(y), v(0, \infty) = k \quad (3.2)$$

for $x, y \in \mathbb{R}_+$, where $\sigma, \tau \in C(\mathbb{R}_+, \mathbb{R})$, $\sigma(x)$ is nondecreasing in x , $\tau(y)$ is nonincreasing in y , k is a real constant, and \mathfrak{T} is a continuous operator on $C(\mathbb{R}_+ \times \mathbb{R}_+, \mathbb{R})$ such that $|\mathfrak{T}v| \leq c_0|v|$ for a constant $c_0 > 0$. Integrating (3.1) with respect to x and y and using the initial conditions (3.2), we get

$$\begin{aligned} v(x, y) = & \sigma(x) + \tau(y) - k - \frac{x}{(x+1)(y+1)} \\ & - \int_0^x \int_y^\infty \exp(-s) \exp(-t) \sqrt{|v(s, t)| + 1} dt ds \\ & - \int_0^x \int_y^\infty s \exp(-s) \exp(-t) \mathfrak{T}v(s, t) dt ds. \end{aligned} \quad (3.3)$$

Thus,

$$\begin{aligned}
 |v(x, y)| &\leq |\sigma(x) + \tau(y) - k| + \frac{x}{(x+1)(y+1)} \\
 &\quad + \int_0^x \int_y^\infty \exp(-s) \exp(-t) \sqrt{|v(s, t)| + 1} dt ds \\
 &\quad + \int_0^x \int_y^\infty s \exp(-s) \exp(-t) c_0 |v(s, t)| dt ds.
 \end{aligned} \tag{3.4}$$

Letting $u(x, y) = |v(x, y)|$, we have

$$u(x, y) \leq a(x, y) + \int_0^x \int_y^\infty d_1(x, y, s, t) w_1(u) dt ds + \int_0^x \int_y^\infty d_2(x, y, s, t) w_2(u) dt ds, \tag{3.5}$$

where $a(x, y) = |\sigma(x) + \tau(y) - k| + x/(x+1)(y+1)$, $w_1(u) = \sqrt{u+1}$, $w_2(u) = c_0 u$, $d_1(x, y, s, t) = \exp(-s) \exp(-t)$, $d_2(x, y, s, t) = s \exp(-s) \exp(-t)$. Clearly, $w_2(u)/w_1(u) = c_0(u/\sqrt{u+1})$ is nondecreasing for $u > 0$, that is, $w_1 \propto w_2$. Then for $u_1, u_2 > 0$,

$$\begin{aligned}
 b_1(x, y) &= a(x, y), \quad \tilde{d}_1(x, y, s, t) = d_1(x, y, s, t), \quad \tilde{d}_2(x, y, s, t) = d_2(x, y, s, t), \\
 W_1(u) &= \int_{u_1}^u \frac{dz}{\sqrt{z+1}} = 2(\sqrt{u+1} - \sqrt{u_1+1}), \quad W_1^{-1}(u) = \left(\frac{u}{2} + \sqrt{u_1+1}\right)^2 - 1, \\
 W_2(u) &= \int_{u_2}^u \frac{dz}{c_0 z} = \frac{1}{c_0} \ln \frac{u}{u_2}, \quad W_2^{-1}(u) = u_2 \exp(c_0 u), \\
 b_2(x, y) &= W_1^{-1} [W_1(b_1(x, y)) + \int_0^x \int_y^\infty \tilde{d}_1(x, y, s, t) dt ds] \\
 &= W_1^{-1} [2(\sqrt{b_1(x, y)+1} - \sqrt{u_1+1}) + (1 - \exp(-x)) \exp(-y)] \\
 &= \left[\sqrt{b_1(x, y)+1} + \frac{1 - \exp(-x)}{2} \exp(-y)\right]^2 - 1.
 \end{aligned} \tag{3.6}$$

By Theorem 2.1, we have

$$\begin{aligned}
 |v(x, y)| &\leq W_2^{-1} [W_2(b_2(x, y)) + \int_0^x \int_y^\infty \tilde{d}_2(x, y, s, t) dt ds] \\
 &= W_2^{-1} \left[\frac{1}{c_0} \ln \frac{b_2(x, y)}{u_2} + (1 - (x+1) \exp(-x)) \exp(-y) \right] \\
 &= u_2 \exp \left[c_0 \left(\frac{1}{c_0} \ln \frac{b_2(x, y)}{u_2} + (1 - (x+1) \exp(-x)) \exp(-y) \right) \right] \\
 &= b_2(x, y) \exp [c_0 (1 - (x+1) \exp(-x)) \exp(-y)] \\
 &= \left[\left(\sqrt{|\sigma(x) + \tau(y) - k| + \frac{x}{(x+1)(y+1)}} + 1 + \frac{1 - \exp(-x)}{2} \exp(-y) \right)^2 - 1 \right] \\
 &\quad \times \exp [c_0 (1 - (x+1) \exp(-x)) \exp(-y)].
 \end{aligned} \tag{3.7}$$

This implies that the solution of (3.1) is bounded for $x, y \in \mathbb{R}_+$ provided that $\sigma(x) + \tau(y) - k$ is bounded for all $x, y \in \mathbb{R}_+$.

References

- [1] M. Pinto, "Integral inequalities of Bihari-type and applications," *Funkcialaj Ekvacioj*, vol. 33, no. 3, pp. 387–403, 1990.
- [2] B. G. Pachpatte, "On some fundamental integral inequalities and their discrete analogues," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 2, no. 2, article 15, pp. 1–13, 2001.
- [3] B. G. Pachpatte, "On some new inequalities related to certain inequalities in the theory of differential equations," *Journal of Mathematical Analysis and Applications*, vol. 189, no. 1, pp. 128–144, 1995.
- [4] O. Lipovan, "A retarded integral inequality and its applications," *Journal of Mathematical Analysis and Applications*, vol. 285, no. 2, pp. 436–443, 2003.
- [5] S. K. Choi, S. Deng, N. J. Koo, and W. Zhang, "Nonlinear integral inequalities of Bihari-type without class H ," *Mathematical Inequalities & Applications*, vol. 8, no. 4, pp. 643–654, 2005.
- [6] R. P. Agarwal, S. Deng, and W. Zhang, "Generalization of a retarded Gronwall-like inequality and Its applications," *Applied Mathematics and Computation*, vol. 165, no. 3, pp. 599–612, 2005.
- [7] S. G. Hristova, "Nonlinear delay integral inequalities for piecewise continuous functions and applications," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 5, no. 4, article 88, pp. 1–14, 2004.
- [8] W. Zhang and S. Deng, "Projected Gronwall-Bellman's inequality for integrable functions," *Mathematical and Computer Modelling*, vol. 34, no. 3-4, pp. 393–402, 2001.
- [9] W.-S. Cheung, "Some retarded Gronwall-Bellman-Ou-Iang-type inequalities and applications to initial boundary value problems," in preparation.
- [10] S. S. Dragomir and Y.-H. Kim, "On certain new integral inequalities and their applications," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 3, no. 4, article 65, pp. 1–8, 2002.
- [11] S. S. Dragomir and Y.-H. Kim, "Some integral inequalities for functions of two variables," *Electronic Journal of Differential Equations*, vol. 2003, no. 10, pp. 1–13, 2003.
- [12] F. W. Meng and W. N. Li, "On some new integral inequalities and their applications," *Applied Mathematics and Computation*, vol. 148, no. 2, pp. 381–392, 2004.
- [13] W.-S. Cheung and Q.-H. Ma, "On certain new Gronwall-Ou-Iang type integral inequalities in two variables and their applications," *Journal of Inequalities and Applications*, vol. 2005, no. 4, pp. 347–361, 2005.

Kelong Zheng: College of Science, Southwest University of Science and Technology,
Mianyang, Sichuan 621010, China
Email address: zhengkelong@swust.edu.cn

Yu Wu: Yibin University, Yibin, Sichuan 644007, China
Email address: wuyu003@yahoo.com.cn

Shengfu Deng: Department of Mathematics, Virginia Polytechnical Institute and State University,
Blacksburg, VA 24061, USA
Email address: sfdeng@vt.edu