

Research Article

Slowly Oscillating Solutions for Differential Equations with Strictly Monotone Operator

Chuanyi Zhang and Yali Guo

Received 2 August 2006; Accepted 28 February 2007

Recommended by Ondrej Dosly

The authors discuss necessary and sufficient conditions for the existence and uniqueness of slowly oscillating solutions for the differential equation $u' + F(u) = h(t)$ with strictly monotone operator. Particularly, the authors give necessary and sufficient conditions for the existence and uniqueness of slowly oscillating solutions for the differential equation $u' + \nabla\Phi(u) = h(t)$, where $\nabla\Phi$ denotes the gradient of the convex function Φ on \mathbb{R}^N .

Copyright © 2007 C. Zhang and Y. Guo. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

In this paper, we will consider the following differential equation:

$$u' + F(u) = h(t), \quad (1.1)$$

where the maps $h : \mathbb{R} \rightarrow \mathbb{R}^N$ and $F : \mathbb{R}^N \rightarrow \mathbb{R}^N$ are continuous. A special class, of the dissipative equation (1.1), is the case where the field F is derived from a convex potential Φ :

$$u' + \nabla\Phi(u) = h(t). \quad (1.2)$$

For the dissipative equation (1.1), Biroli [1], Dafermos [2], Haraux [3], Huang [4], and Ishii [5] have given important contributions to the question of almost periodic solutions which are valid even for the abstract evolution equations. In [6], Philippe Cieutat gives necessary and sufficient conditions for the existence and uniqueness of the bounded (resp., almost periodic) solution of (1.2) when the forcing term $h(t)$ is bounded (resp., almost periodic). In the scalar case $N = 1$, Slyusarchuk established similar results in [7]. But the conditions which are established in [6] for (1.2) do not hold for (1.1), even in

the linear case. So in [6] Cieutat also gives a sufficient condition, then a necessary condition, for the existence and uniqueness of the bounded (resp., almost periodic) solution of (1.1).

The numerical space \mathbb{R}^N is endowed with its standard inner product $\sum_{k=1}^N x_k y_k$, $|\cdot|$ denotes the associated Euclidian norm. We denote by $BC(\mathbb{R}^N)$ the Banach space of continuous bounded functions from \mathbb{R} to \mathbb{R}^N endowed with the norm $\|u\|_\infty := \sup_{t \in \mathbb{R}} |u(t)|$. When k is a positive integer, $BC^k(\mathbb{R}^N)$ is the space of functions in $BC(\mathbb{R}^N) \cap C^k(\mathbb{R}^N)$ such that all their derivatives, up to order k , are bounded functions. When $u \in BC^1(\mathbb{R}^N)$, we set $\|u\|_{C^1} = \|u\|_\infty + \|u'\|_\infty$ and when $u \in BC^2(\mathbb{R}^N)$, we set $\|u\|_{C^2} = \|u\|_\infty + \|u'\|_\infty + \|u''\|_\infty$.

In 1984, Sarason in [8] extended almost periodic functions and introduced the definition of remotely periodic functions. The space of remotely periodic functions, as a C^* -subalgebra of $BC(\mathbb{R}^N)$, is generated by almost periodic functions and slowly oscillating functions which are defined as following.

Definition 1.1 [8]. A function $f \in BC(\mathbb{R}^N)$ is said to be slowly oscillating if

$$\lim_{|t| \rightarrow +\infty} |f(t+a) - f(t)| = 0, \quad \text{for each } a \in \mathbb{R}, \tag{1.3}$$

the set of all these functions is denoted by $SO(\mathbb{R}^N)$.

Comparing with the space $AP(\mathbb{R})$ of almost periodic functions, the space of slowly oscillating functions is quite large. In fact, $AP(\mathbb{R}) = \overline{\text{span}}\{e^{i\lambda t} : \lambda \in \mathbb{R}\}$, where the closure is taken in $BC(\mathbb{R})$ (e.g., see [9] for details). $SO(\mathbb{R})$ not only contains such space as $C_0(\mathbb{R})$ which consists of all the functions f such that $f(t) \rightarrow 0$ as $|t| \rightarrow \infty$, but also properly contains $X = \overline{\text{span}}\{e^{i\lambda t^\alpha} : \lambda \in \mathbb{R}, 0 < \alpha < 1\}$ (see [10–12] for details). The only functions in $AP(\mathbb{R}) \cap SO(\mathbb{R})$ are the constant functions on \mathbb{R} . We also point out that the slowly oscillating functions $SO(\mathbb{R}^N)$ studied here form a strict subset of the slowly oscillating functions studied on [13, page 250, Definition 4.2.1]. Thus, all functions in $SO(\mathbb{R}^N)$ are uniformly continuous.

To our knowledge, nobody has investigated the existence and uniqueness of slowly oscillating solutions for the differential equation (1.1). So in this paper, we give a sufficient, then a necessary condition for the existence and uniqueness of slowly oscillating solutions for the differential equation (1.1). We will give sufficient and necessary conditions for the existence and uniqueness of slowly oscillating solutions for differential equation (1.2).

To show the main results of the paper, we need the following definition and lemma.

Definition 1.2. A function $F : \mathbb{R}^N \rightarrow \mathbb{R}^N$ is said to be strictly monotone on \mathbb{R}^N if $(F(x_1) - F(x_2), x_1 - x_2) > 0$ for all $x_1, x_2 \in \mathbb{R}^N$ such that $x_1 \neq x_2$.

LEMMA 1.3 [6]. *Let $F : \mathbb{R}^N \rightarrow \mathbb{R}^N$ be a continuous and strictly monotone map. Then for every compact subset K of \mathbb{R}^N and for every $\varepsilon > 0$, there exists $c > 0$ such that*

$$(F(x_1) - F(x_2), x_1 - x_2) > c |x_1 - x_2|^2 \tag{1.4}$$

for all $x_1, x_2 \in K$ such that

$$|x_1 - x_2| \geq \varepsilon. \tag{1.5}$$

2. Main results

For each $u \in BC^1(\mathbb{R}^N)$, the function $t \rightarrow u'(t) + F(u(t))$ belongs to $BC(\mathbb{R}^N)$, so we can define the operator $\mathcal{F}_1 : BC^1(\mathbb{R}^N) \rightarrow BC^0(\mathbb{R}^N)$ with $\mathcal{F}_1(u)(t) := u'(t) + F(u(t))$ for all $u \in BC^1(\mathbb{R}^N)$ and $t \in \mathbb{R}$. Let $SO^1(\mathbb{R}^N) = SO(\mathbb{R}^N) \cap C^1(\mathbb{R}^N)$ and $\|u\|_{C^1} = \|u\|_\infty + \|u'\|_\infty$ for $u \in SO^1(\mathbb{R}^N)$. Set $\mathcal{F}_2 = \mathcal{F}_1|_{SO^1(\mathbb{R}^N)}$.

Consider the following assertions:

(A) F is a strictly monotone map on \mathbb{R}^N such that

$$\lim_{|x| \rightarrow \infty} \frac{F(x), x}{|x|} = +\infty; \quad (2.1)$$

(B) $\mathcal{F}_2 : (SO^1(\mathbb{R}^N), \|\cdot\|_{C^1}) \rightarrow (SO(\mathbb{R}^N), \|\cdot\|_\infty)$ is a homeomorphism;

(C) $F : (\mathbb{R}^N, |\cdot|) \rightarrow (\mathbb{R}^N, |\cdot|)$ is a homeomorphism.

THEOREM 2.1. *Let $F : \mathbb{R}^N \rightarrow \mathbb{R}^N$ be a continuous map. Then the following implications hold: (A) \Rightarrow (B) \Rightarrow (C).*

Proof. By [6], (A) implies that $\mathcal{F}_1 : (BC^1(\mathbb{R}^N), \|\cdot\|_{C^1}) \rightarrow (BC(\mathbb{R}^N), \|\cdot\|_\infty)$ is a homeomorphism. That is, (1.1) has for each h from $SO(\mathbb{R}^N)$ a unique solution $u(t)$ from $SO^1(\mathbb{R}^N)$, which depends continuously on h . To show (A) \Rightarrow (B), it remains to show $u \in SO(\mathbb{R}^N)$ if $h \in SO(\mathbb{R}^N)$.

Suppose, by the way of contradiction, $u(t) \notin SO(\mathbb{R}^N)$. Then there exist $a_0, \varepsilon_0 > 0$ and sequence $t_n \rightarrow \infty$ such that

$$|u(t_n + a_0) - u(t_n)| \geq \varepsilon_0. \quad (2.2)$$

Without loss of generality, we can assume $t_n - t_{n-1} \rightarrow +\infty$.

Since $u(t) \in BC^1(\mathbb{R}^N)$, $u(t)$ is uniformly continuous on \mathbb{R} . Then there exists $\delta > 0$ such that

$$|u(t + a_0) - u(t)| \geq \frac{\varepsilon_0}{2}, \quad \forall t \in P'_n, \quad (2.3)$$

where $P'_n = (t_n - \delta, t_n + \delta)$.

Set

$$\begin{aligned} P_n &= (t_n - \delta, t_n), & I_n &= [t_{n-1}, t_n], \\ C_n &= \left\{ s \in I_n : |u(s + a_0) - u(s)| \geq \frac{\varepsilon_0}{2} \right\}, & C'_n &= \left\{ s \in I_n : |u(s + a_0) - u(s)| < \frac{\varepsilon_0}{2} \right\} \end{aligned} \quad (2.4)$$

and put

$$\Phi(t) = u(t + a_0) - u(t). \quad (2.5)$$

Obviously $P_n \subset C_n$.

4 Journal of Inequalities and Applications

Let $K = \overline{u(\mathbb{R})}$. Note that $u(t)$ is bounded on \mathbb{R}^N and therefore, K is a compact subset of \mathbb{R}^N . By Lemma 1.3 there exists c_o such that

$$(F(u(s+a_o)) - F(u(s)), u(s+a_o) - u(s)) \geq c_o |u(s+a_o) - u(s)|^2 \quad (2.6)$$

for each $s \in C_n$.

Note

$$\int_{t_{n-1}}^{t_n} \frac{d}{ds} \left(\frac{1}{2} |\Phi(s)|^2 e^{2c_0 s} \right) ds = \frac{1}{2} |\Phi(t_n)|^2 \cdot e^{2c_0 t_n} - \frac{1}{2} |\Phi(t_{n-1})|^2 \cdot e^{2c_0 t_{n-1}}. \quad (2.7)$$

At the same time, we also have

$$\int_{t_{n-1}}^{t_n} \frac{d}{ds} \left(\frac{1}{2} |\Phi(s)|^2 e^{2c_0 s} \right) ds = \int_{C_n} \frac{d}{ds} \left(\frac{1}{2} |\Phi(s)|^2 e^{2c_0 s} \right) ds + \int_{C'_n} \frac{d}{ds} \left(\frac{1}{2} |\Phi(s)|^2 e^{2c_0 s} \right) ds. \quad (2.8)$$

Case 1. $s \in C_n$, that is,

$$|u(s+a_o) - u(s)| \geq \frac{\varepsilon_0}{2}. \quad (2.9)$$

We can get

$$\begin{aligned} \frac{d}{ds} \left(\frac{1}{2} |\Phi(s)|^2 \right) &= (\Phi(s)', \Phi(s)) \\ &= (h(s+a_o) - h(s), \Phi(s)) - (F(u(s+a_o)) - F(u(s)), \Phi(s)). \end{aligned} \quad (2.10)$$

By (2.6), we can obtain

$$\frac{d}{ds} \left(\frac{1}{2} |\Phi(s)|^2 \right) \leq |h(s+a_o) - h(s)| \cdot |\Phi(s)| - c_o |\Phi(s)|^2. \quad (2.11)$$

Also we can see

$$\frac{d}{ds} \left(\frac{1}{2} |\Phi(s)|^2 e^{2c_0 s} \right) = (\Phi'(s), \Phi(s)) \cdot e^{2c_0 s} + c_0 e^{2c_0 s} \cdot |\Phi(s)|^2. \quad (2.12)$$

By (2.11), we deduce

$$\frac{d}{ds} \left(\frac{1}{2} |\Phi(s)|^2 e^{2c_0 s} \right) \leq |h(s+a_o) - h(s)| \cdot |\Phi(s)| \cdot e^{2c_0 s}. \quad (2.13)$$

Case 2. $s \in C'_n$, that is,

$$|u(s+a_o) - u(s)| < \frac{\varepsilon_0}{2}. \quad (2.14)$$

Moreover, one has

$$\frac{d}{ds} \left(\frac{1}{2} |\Phi(s)|^2 e^{2c_0 s} \right) = (\Phi'(s), \Phi(s)) \cdot e^{2c_0 s} + c_0 e^{2c_0 s} \cdot |\Phi(s)|^2, \quad (2.15)$$

$$(\Phi(s)', \Phi(s)) = (h(s+a_o) - h(s), \Phi(s)) - (F(u(s+a_o)) - F(u(s)), \Phi(s)). \quad (2.16)$$

For F is strictly monotone on \mathbb{R}^N , we can deduce

$$(F(u(s+a_0)) - F(u(s)), \Phi(s)) > 0. \quad (2.17)$$

Moreover, by (2.17) and (2.16) one has

$$(\Phi(s)', \Phi(s)) < |\Phi(s)| \cdot |h(s+a_0) - h(s)| < \frac{\varepsilon_0}{2} |h(s+a_0) - h(s)|. \quad (2.18)$$

By (2.15) and (2.18), we can get

$$\frac{d}{ds} \left(\frac{1}{2} |\Phi(s)|^2 e^{2c_0 s} \right) < \frac{\varepsilon_0}{2} |h(s+a_0) - h(s)| \cdot e^{2c_0 s} + \frac{\varepsilon_0^2}{4} c_0 e^{2c_0 s}. \quad (2.19)$$

Considering the above two cases, one has

$$\begin{aligned} & \frac{1}{2} |\Phi(t_n)|^2 \cdot e^{2c_0 t_n} - \frac{1}{2} |\Phi(t_{n-1})|^2 \cdot e^{2c_0 t_{n-1}} \\ &= \int_{t_{n-1}}^{t_n} \frac{d}{ds} \left(\frac{1}{2} |\Phi(s)|^2 e^{2c_0 s} \right) ds \\ &< \int_{C_n} |h(s+a_0) - h(s)| \cdot |\Phi(s)| \cdot e^{2c_0 s} ds \\ &\quad + \int_{C'_n} \left(\frac{\varepsilon_0}{2} |h(s+a_0) - h(s)| + \frac{\varepsilon_0^2}{4} c_0 \right) \cdot e^{2c_0 s} ds \\ &\leq \sup_{t \in I_n} |h(t+a_0) - h(t)| \cdot \sup_{t \in I_n} |\Phi(t)| \cdot \int_{C_n} e^{2c_0 s} ds \\ &\quad + \left(\frac{\varepsilon_0}{2} \sup_{t \in I_n} |h(t+a_0) - h(t)| + \frac{\varepsilon_0^2}{4} c_0 \right) \cdot \int_{C'_n} e^{2c_0 s} ds. \end{aligned} \quad (2.20)$$

Since

$$\begin{aligned} \int_{C_n} e^{2c_0 s} ds &\leq \int_{t_{n-1}}^{t_n} e^{2c_0 s} ds, \\ \int_{C'_n} e^{2c_0 s} ds &\leq \int_{t_{n-1}}^{t_n} e^{2c_0 s} ds - \int_{t_{n-\delta}}^{t_{n-1}} e^{2c_0 s} ds, \end{aligned} \quad (2.21)$$

one has

$$\begin{aligned} & \frac{1}{2} |\Phi(t_n)|^2 \cdot e^{2c_0 t_n} - \frac{1}{2} |\Phi(t_{n-1})|^2 \cdot e^{2c_0 t_{n-1}} \\ &\leq \sup_{t \in I_n} |h(t+a_0) - h(t)| \cdot \sup_{t \in I_n} |\Phi(t)| \cdot \int_{t_{n-1}}^{t_n} e^{2c_0 s} ds \\ &\quad + \left(\frac{\varepsilon_0}{2} \sup_{t \in I_n} |h(t+a_0) - h(t)| + \frac{\varepsilon_0^2}{4} c_0 \right) \cdot \left(\int_{t_{n-1}}^{t_n} e^{2c_0 s} ds - \int_{t_{n-\delta}}^{t_{n-1}} e^{2c_0 s} ds \right). \end{aligned} \quad (2.22)$$

So, we can get

$$\begin{aligned} & \frac{1}{2} |\Phi(t_n)|^2 \cdot e^{2c_0 t_n} - \frac{1}{2} |\Phi(t_{n-1})|^2 \cdot e^{2c_0 t_{n-1}} \\ & \leq \sup_{t \in I_n} |h(t+a_0) - h(t)| \cdot \sup_{t \in I_n} |\Phi(t)| \cdot \frac{1}{2c_0} (e^{2c_0 t_n} - e^{2c_0 t_{n-1}}) \\ & \quad + \left(\frac{\varepsilon_0}{2} \sup_{t \in I_n} |h(t+a_0) - h(t)| + \frac{\varepsilon_0^2}{4} c_0 \right) \cdot \left[\frac{1}{2c_0} (e^{2c_0 t_n} - e^{2c_0 t_{n-1}}) - e^{2c_0(t_n-\delta)} \cdot (e^{2c_0 \delta} - 1) \cdot \frac{1}{2c_0} \right]. \end{aligned} \tag{2.23}$$

That is,

$$\begin{aligned} & \frac{1}{2} |\Phi(t_n)|^2 \cdot e^{2c_0 t_n} - \frac{1}{2} |\Phi(t_{n-1})|^2 \cdot e^{2c_0 t_{n-1}} \\ & \leq \sup_{t \in I_n} |h(t+a_0) - h(t)| \\ & \quad \cdot \left[\frac{e^{2c_0 t_n} - e^{2c_0 t_{n-1}}}{2c_0} \cdot \sup_{t \in I_n} |\Phi(t)| + \frac{\varepsilon_0}{2} \cdot \left(\frac{e^{2c_0 t_n} - e^{2c_0 t_{n-1}}}{2c_0} - \frac{(e^{2c_0 \delta} - 1)}{2c_0} \cdot e^{2c_0(t_n-\delta)} \right) \right] \\ & \quad + \left(\frac{\varepsilon_0^2 (e^{2c_0 t_n} - e^{2c_0 t_{n-1}})}{8} - \frac{\varepsilon_0^2 (e^{2c_0 \delta} - 1)}{8} \cdot e^{2c_0(t_n-\delta)} \right). \end{aligned} \tag{2.24}$$

Thus,

$$\begin{aligned} & \sup_{t \in I_n} |h(t+a_0) - h(t)| \\ & \geq \frac{\frac{1}{2} |\Phi(t_n)|^2 \cdot e^{2c_0 t_n} - \frac{1}{2} |\Phi(t_{n-1})|^2 \cdot e^{2c_0 t_{n-1}} - \frac{\varepsilon_0^2 (e^{2c_0 t_n} - e^{2c_0 t_{n-1}})}{8} + \frac{\varepsilon_0^2 (e^{2c_0 \delta} - 1)}{8} \cdot e^{2c_0(t_n-\delta)}}{\sup_{t \in I_n} |\Phi(t)| \cdot \frac{1}{2c_0} (e^{2c_0 t_n} - e^{2c_0 t_{n-1}}) + \frac{\varepsilon_0}{2} \cdot \left[\frac{1}{2c_0} (e^{2c_0 t_n} - e^{2c_0 t_{n-1}}) - e^{2c_0(t_n-\delta)} \cdot (e^{2c_0 \delta} - 1) \cdot \frac{1}{2c_0} \right]} \\ & = \frac{(4c_0 |\Phi(t_n)|^2 - \varepsilon_0^2 c_0) \cdot e^{2c_0 t_n} - (4c_0 |\Phi(t_{n-1})|^2 - \varepsilon_0^2 c_0) \cdot e^{2c_0 t_{n-1}} + \varepsilon_0^2 c_0 (e^{2c_0 \delta} - 1) \cdot e^{2c_0(t_n-\delta)}}{(4 \sup_{t \in I_n} |\Phi(t)| + 2\varepsilon_0) \cdot (e^{2c_0 t_n} - e^{2c_0 t_{n-1}}) - 2\varepsilon_0 (e^{2c_0 \delta} - 1) \cdot e^{2c_0(t_n-\delta)}} \\ & \geq \frac{(4c_0 |\Phi(t_n)|^2 - \varepsilon_0^2 c_0) \cdot e^{2c_0 t_n} - (4c_0 |\Phi(t_{n-1})|^2 - \varepsilon_0^2 c_0) \cdot e^{2c_0 t_{n-1}} + \varepsilon_0^2 c_0 (e^{2c_0 \delta} - 1) \cdot e^{2c_0(t_n-\delta)}}{(4 \sup_{t \in I_n} |\Phi(t)| + 2\varepsilon_0) \cdot (e^{2c_0 t_n} - e^{2c_0 t_{n-1}})} \\ & = \frac{(4c_0 |\Phi(t_n)|^2 - \varepsilon_0^2 c_0) - (4c_0 |\Phi(t_{n-1})|^2 - \varepsilon_0^2 c_0) \cdot e^{2c_0(t_{n-1}-t_n)} + \varepsilon_0^2 c_0 (e^{2c_0 \delta} - 1) \cdot e^{-2c_0 \delta}}{(4 \sup_{t \in I_n} |\Phi(t)| + 2\varepsilon_0) \cdot (1 - e^{2c_0(t_{n-1}-t_n)})}. \end{aligned} \tag{2.25}$$

Since $\Phi(t) = u(t+a_0) - u(t)$ and the solution $u(t)$ is bounded, then we can assume $\exists M > 0$, such that

$$|\Phi(t)| < M, \quad \text{for each } t \in \mathbb{R}. \tag{2.26}$$

Also we have

$$|\Phi(t_n)|^2 \geq \frac{\varepsilon_0^2}{4}, \quad \text{for } t_n \in C_n. \quad (2.27)$$

Then

$$\sup_{t \in I_n} |h(t+a_0) - h(t)| \geq \frac{-(4c_0M^2 - \varepsilon_0^2c_0) \cdot e^{2c_0(t_{n-1}-t_n)} + \varepsilon_0^2c_0(e^{2c_0\delta} - 1) \cdot e^{-2c_0\delta}}{(4M + 2\varepsilon_0) \cdot (1 - e^{2c_0(t_{n-1}-t_n)})}. \quad (2.28)$$

When $n \rightarrow +\infty$, one has

$$t_n - t_{n-1} \rightarrow +\infty, \quad e^{2c_0(t_{n-1}-t_n)} \rightarrow 0. \quad (2.29)$$

So

$$\lim_{t \rightarrow +\infty} |h(t+a_0) - h(t)| > \frac{1}{2} \cdot \frac{\varepsilon_0^2c_0(e^{2c_0\delta} - 1) \cdot e^{-2c_0\delta}}{4M + 2\varepsilon_0} = \frac{\varepsilon_0^2c_0(e^{2c_0\delta} - 1)}{2e^{2c_0\delta}(4M + 2\varepsilon_0)} > 0. \quad (2.30)$$

This contradicts the fact $h(t) \in SO(\mathbb{R}^N)$. We must have $u(t) \in SO(\mathbb{R}^N)$.

Finally we show that

$$(B) \implies (C). \quad (2.31)$$

If we denote by \mathcal{C} the set of constant mapping from \mathbb{R} to \mathbb{R}^N , one has $\mathcal{C} \subset SO^1(\mathbb{R}^N)$ and for $u \in \mathcal{C}$, the function $\mathcal{F}_3(u) \in SO(\mathbb{R}^N)$ ($\mathcal{F}_3(u) = F(u(0))$), for all $t \in \mathbb{R}$, so we can define the restriction operator of \mathcal{F}_3 to \mathcal{C} by $\mathcal{F}_4: \mathcal{C} \rightarrow \mathcal{C}$ with $\mathcal{F}_4(u) = F(u(0))$ for all $u \in \mathcal{C}$ and all $t \in \mathbb{R}$. For $u \in \mathcal{C}$, one has $\|u\|_{C^1} = |u(0)|$ and $\|\mathcal{F}_3(u)\|_\infty = |F(u(0))|$; then it is equivalent to prove \mathcal{F}_4 or F is a homeomorphism. It remains to prove that \mathcal{F}_4 is surjective. Let $h \in \mathcal{C}$. By hypothesis, there exists $u \in SO^1(\mathbb{R}^N)$ such that $\mathcal{F}_3(u) = h$. we want to prove that $u \in \mathcal{C}$. For that we denote by $u_a(t) = u(t+a)$ for all t and $a \in \mathbb{R}$. Note that $\mathcal{F}_3(u_a) = h$ for all $a \in \mathbb{R}$. By injectivity of \mathcal{F}_3 , we deduce that $u_a(t) = u(t)$ for all $a \in \mathbb{R}$, therefore $u \in \mathcal{C}$.

Remark 2.2. The following example constructed in [6] can be used to show that Assertion (A) is not a necessary condition for the existence or the uniqueness of a bounded or slowly oscillating solution of (1.1). Consider the map $F: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $F(x_1, x_2) = Bx = (-x_2, x_1 + x_2)$. The map F is monotone and does not satisfy (A). However, the eigenvalues of B are conjugate and their real parts are equal to $1/2$, therefore the linear system $u' + Bu = 0$ has an exponential dichotomy: namely, there exists $k > 0$ such that $\|\exp(-Bt)\|_{L(\mathbb{R})} \leq k \exp(-t/2)$ for all $t \geq 0$. As a consequence, the system $u' + Bu = 0$ has precisely one bounded solution on \mathbb{R} : $u = 0$; this implies the injectivity of \mathcal{F}_1 . Moreover, the following function $u(t) := \int_{-\infty}^t \exp(-B(t-s))h(s)ds$ is a solution of $u' + Bu = h$ for $h \in BC(\mathbb{R}^N)$ and satisfies $\|u(t)\| \leq 2k\|h\|_\infty$ for all $t \in \mathbb{R}$; this implies the surjectivity of \mathcal{F}_1 . Since \mathcal{F}_1 is a bounded linear map between Banach spaces, which is bijective, then \mathcal{F}_1 is an isomorphism between $BC^1(\mathbb{R}^2)$ and $BC(\mathbb{R}^N)$. To show that (B) holds in this case, it

remains to show that $u \in SO(\mathbb{R}^2)$ if $h \in SO(\mathbb{R}^2)$. In fact, for $a \in \mathbb{R}$

$$\begin{aligned} |u(t+a) - u(t)| &= \left| \int_{-\infty}^{t+a} \exp(-B(t+a-s))h(s)ds - \int_{-\infty}^t \exp(-B(t-s))h(s)ds \right| \\ &\leq \int_{-\infty}^t \exp(-B(t-s)) |h(s+a) - h(s)| ds. \end{aligned} \tag{2.32}$$

It follows that $|u(t+a) - u(t)| \rightarrow 0$ as $t \rightarrow -\infty$. Since $h \in SO(\mathbb{R}^2)$, for $\varepsilon > 0$ there exists $t_0 > 0$ such that $|h(t+a) - h(t)| < \varepsilon$ for all $t > t_0$. Now

$$\begin{aligned} |u(t+a) - u(t)| &\leq \left(\int_{-\infty}^{t_0} + \int_{t_0}^t \right) \exp(-B(t-s)) |h(s+a) - h(s)| ds \\ &\leq 2\|h\|_\infty \int_{-\infty}^{t_0} \exp(-B(t-s)) ds + \varepsilon \int_{t_0}^t \exp(-B(t-s)) ds \end{aligned} \tag{2.33}$$

and therefore, $u(t+a) - u(t) \rightarrow 0$ as $t \rightarrow +\infty$. This shows that $u \in SO(\mathbb{R}^2)$.

Remark 2.3. The following example constructed also in [6] can be used to show that (C) is not a sufficient condition for the existence of slowly oscillating solution of (1.1) even when F is a linear monotone map. Consider the map $F: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $F(x_1, x_2) := Ax = (-x_2, x_1)$. F is a homeomorphism and a monotone map. Let $v = (\sin t, \cos t)$. Then $v' + Av = 0$. Let f be any continuously differentiable function on \mathbb{R} such that $f(t) = t^{1/3}$ for $|t| > 1$ and let $h(t) = f'(t)v(t)$. Since $h(t) \rightarrow 0$ as $|t| \rightarrow \infty$, $h \in SO(\mathbb{R}^2)$. The equation $u' + Au = h$ has no bounded solution, because $u(t) = f(t)v(t)$ is an unbounded solution, therefore \mathcal{F}_2 is not surjective.

Nevertheless, for (1.2) we have the following result.

THEOREM 2.4. *Let Φ be a convex and continuously differentiable function on \mathbb{R}^N . Assume that $F = \nabla\Phi$. Then (A), (B), and (C) are equivalent.*

We have already shown that (A) \Rightarrow (B) \Rightarrow (C) in Theorem 2.1. The equivalence of (A) and (C) is [6, Theorem 1.1]. □

Acknowledgments

The authors would like to thank the referee for the valuable comments. The authors also would like to thank Professor Ondrej Dosly for his regards. The research is supported by the NSF of China (no. 10671046).

References

- [1] M. Biroli, "Sur les solutions bornées ou presque périodiques des équations d'évolution multivoques sur un espace de Hilbert," *Ricerche di Matematica*, vol. 21, pp. 17–47, 1972.
- [2] C. M. Dafermos, "Almost periodic processes and almost periodic solutions of evolution equations," in *Dynamical Systems (Proc. Internat. Sympos., Univ. Florida, Gainesville, Fla., 1976)*, pp. 43–57, Academic Press, New York, NY, USA, 1977.
- [3] A. Haraux, *Nonlinear Evolution Equations—Global Behavior of Solutions*, vol. 841 of *Lecture Notes in Mathematics*, Springer, Berlin, Germany, 1981.

- [4] F. L. Huang, “Existence of almost periodic solutions for dissipative differential equations,” *Annals of Differential Equations*, vol. 6, no. 3, pp. 271–279, 1990.
- [5] H. Ishii, “On the existence of almost periodic complete trajectories for contractive almost periodic processes,” *Journal of Differential Equations*, vol. 43, no. 1, pp. 66–72, 1982.
- [6] P. Cieutat, “Necessary and sufficient conditions for existence and uniqueness of bounded or almost-periodic solutions for differential systems with convex potential,” *Differential and Integral Equations*, vol. 18, no. 4, pp. 361–378, 2005.
- [7] V. E. Slyusarchuk, “Necessary and sufficient conditions for existence and uniqueness of bounded and almost-periodic solutions of nonlinear differential equations,” *Acta Applicandae Mathematicae*, vol. 65, no. 1–3, pp. 333–341, 2001.
- [8] D. Sarason, “Remotely almost periodic functions,” in *Proceedings of the Conference on Banach Algebras and Several Complex Variables (New Haven, Conn., 1983)*, vol. 32 of *Contemporary Mathematics*, pp. 237–242, American Mathematical Society, Providence, RI, USA, 1984.
- [9] C. Zhang, *Almost Periodic Type Functions and Ergodicity*, Science Press, Beijing, China, 2003.
- [10] C. Zhang, “New limit power function spaces,” *IEEE Transactions on Automatic Control*, vol. 49, no. 5, pp. 763–766, 2004.
- [11] C. Zhang, “Strong limit power functions,” *The Journal of Fourier Analysis and Applications*, vol. 12, no. 3, pp. 291–307, 2006.
- [12] C. Zhang and C. Meng, “ C^* -algebra of strong limit power functions,” *IEEE Transactions on Automatic Control*, vol. 51, no. 5, pp. 828–831, 2006.
- [13] W. Arendt, C. J. K. Batty, M. Hieber, and F. Neubrander, *Vector-Valued Laplace Transforms and Cauchy Problems*, vol. 96 of *Monographs in Mathematics*, Birkhäuser, Basel, Switzerland, 2001.

Chuanyi Zhang: Department of Mathematics, Harbin Institute of Technology, Harbin 150001, China
Email address: czhang@hit.edu.cn

Yali Guo: Department of Mathematics, Harbin Institute of Technology, Harbin 150001, China
Email address: yali0520@sina.com