

Research Article

An Optimal Double Inequality between Power-Type Heron and Seiffert Means

Yu-Ming Chu,¹ Miao-Kun Wang,² and Ye-Fang Qiu²

¹ Department of Mathematics, Huzhou Teachers College, Huzhou 313000, China

² Department of Mathematics, Zhejiang Sci-Tech University, Hangzhou 310018, China

Correspondence should be addressed to Yu-Ming Chu, chuyuming2005@yahoo.com.cn

Received 29 August 2010; Accepted 16 November 2010

Academic Editor: Alexander I. Domoshnitsky

Copyright © 2010 Yu-Ming Chu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

For $k \in [0, +\infty)$, the power-type Heron mean $H_k(a, b)$ and the Seiffert mean $T(a, b)$ of two positive real numbers a and b are defined by $H_k(a, b) = ((a^k + (ab)^{k/2} + b^k)/3)^{1/k}$, $k \neq 0$; $H_k(a, b) = \sqrt{ab}$, $k = 0$ and $T(a, b) = (a - b)/2 \arctan((a - b)/(a + b))$, $a \neq b$; $T(a, b) = a$, $a = b$, respectively. In this paper, we find the greatest value p and the least value q such that the double inequality $H_p(a, b) < T(a, b) < H_q(a, b)$ holds for all $a, b > 0$ with $a \neq b$.

1. Introduction

For $k \in [0, +\infty)$, the power-type Heron mean $H_k(a, b)$ and the Seiffert mean $T(a, b)$ of two positive real numbers a and b are defined by

$$H_k(a, b) = \begin{cases} \left(\frac{a^k + (ab)^{k/2} + b^k}{3} \right)^{1/k}, & k \neq 0, \\ \sqrt{ab}, & k = 0, \end{cases} \quad (1.1)$$

$$T(a, b) = \begin{cases} \frac{a - b}{2 \arctan((a - b)/(a + b))}, & a \neq b, \\ a, & a = b, \end{cases} \quad (1.2)$$

respectively.

Recently, the means of two variables have been the subject of intensive research [1–15]. In particular, many remarkable inequalities for $H_k(a, b)$ and $T(a, b)$ can be found in the literature [16–20].

It is well known that $H_k(a, b)$ is continuous and strictly increasing with respect to $k \in [0, +\infty)$ for fixed $a, b > 0$ with $a \neq b$. Let $A(a, b) = (a + b)/2$, $I(a, b) = (1/e)(b^b/a^a)^{1/(b-a)}$, $L(a, b) = (b - a)/(\log b - \log a)$, $G(a, b) = \sqrt{ab}$, and $H(a, b) = 2ab/(a + b)$ be the arithmetic, identric, logarithmic, geometric, and harmonic means of two positive numbers a and b with $a \neq b$, respectively. Then

$$\min\{a, b\} < H(a, b) < G(a, b) < L(a, b) < I(a, b) < A(a, b) < \max\{a, b\}. \quad (1.3)$$

For $p \in \mathbb{R}$, the power mean $M_p(a, b)$ of order p of two positive numbers a and b is defined by

$$M_p(a, b) = \begin{cases} \left(\frac{a^p + b^p}{2}\right)^{1/p}, & p \neq 0, \\ \sqrt{ab}, & p = 0. \end{cases} \quad (1.4)$$

The main properties for power mean are given in [21].

In [16], Jia and Cao presented the inequalities

$$\begin{aligned} H_0(a, b) = G(a, b) < L(a, b) < H_p(a, b) < M_q(a, b), \\ A(a, b) < H_{\log 3 / \log 2}(a, b) \end{aligned} \quad (1.5)$$

for all $a, b > 0$ with $a \neq b$, $p \geq 1/2$, and $q \geq (2/3)p$.

Sándor [22] proved that

$$I(a, b) > H_1(a, b) \quad (1.6)$$

for all $a, b > 0$ with $a \neq b$.

In [19], Seiffert established that

$$M_1(a, b) < T(a, b) < M_2(a, b) \quad (1.7)$$

for all $a, b > 0$ with $a \neq b$.

The purpose of this paper is to present the optimal upper and lower power-type Heron mean bounds for the Seiffert mean $T(a, b)$. Our main result is the following Theorem 1.1.

Theorem 1.1. *For all $a, b > 0$ with $a \neq b$, one has*

$$H_{\log 3 / \log(\pi/2)}(a, b) < T(a, b) < H_{5/2}(a, b), \quad (1.8)$$

and $H_{\log 3 / \log(\pi/2)}(a, b)$ and $H_{5/2}(a, b)$ are the best possible lower and upper power-type Heron mean bounds for the Seiffert mean $T(a, b)$, respectively.

2. Lemmas

In order to prove our main result, Theorem 1.1, we need two lemmas which we present in this section.

Lemma 2.1. *If $k = \log 3 / \log(\pi/2) = 2.43\dots$ and $t > 1$, then*

$$\begin{aligned} & -24(k-2)(k+3)(k+4)(3k+2)t^{k+8} + 48k(k-1)(k+3)(3k-2)t^{k+6} \\ & - (k+4)(k+6)(k+8)(4k-7)t^8 < 0. \end{aligned} \quad (2.1)$$

Proof. For $t > 1$, we clearly see that

$$\begin{aligned} & -24(k-2)(k+3)(k+4)(3k+2)t^{k+8} + 48k(k-1)(k+3) \\ & \times (3k-2)t^{k+6} - (k+4)(k+6)(k+8)(4k-7)t^8 \\ & < t^8 \left[-24(k-2)(k+3)(k+4)(3k+2)t^2 + 48k(k-1)(k+3) \right. \\ & \left. \times (3k-2)t - (k+4)(k+6)(k+8)(4k-7) \right]. \end{aligned} \quad (2.2)$$

Let

$$\begin{aligned} h(t) &= -24(k-2)(k+3)(k+4)(3k+2)t^2 + 48k(k-1)(k+3) \\ & \times (3k-2)t - (k+4)(k+6)(k+8)(4k-7). \end{aligned} \quad (2.3)$$

Then

$$h(1) = 68k^4 - 281k^3 - 1010k^2 + 2072k + 2496 = -104.992\dots < 0 \quad (2.4)$$

and $h(t)$ is strictly decreasing in $[1, +\infty)$ because of $k(k-1)(k+3)(3k-2)/(k-2)(k+3)(k+4)(3k+2) < 1$ for $k = \log 3 / \log(\pi/2)$.

Therefore, Lemma 2.1 follows from (2.2)–(2.4) together with the monotonicity of $h(t)$. \square

Lemma 2.2. *If $k = \log 3 / \log(\pi/2) = 2.43\dots$, $t \in [1, +\infty)$, and $g(t) = -8t^{4k-4} + 8t^{4k-6} + (2-k)t^{3k+2} + 2kt^{3k} - 2(k+2)t^{3k-2} + 2(k-4)t^{3k-4} + (10-k)t^{3k-6} + (7-4k)t^{2k+2} + 2(4k-1)t^{2k} - 2(4k+5)t^{2k-2} + 2(4k-1)t^{2k-4} + (7-4k)t^{2k-6} + (10-k)t^{k+2} + 2(k-4)t^k - 2(k+2)t^{k-2} + 2kt^{k-4} + (2-k)t^{k-6} + 8t^2 - 8$, then there exists $\lambda \in (1, +\infty)$ such that $g(t) > 0$ for $t \in (1, \lambda)$ and $g(t) < 0$ for $t \in (\lambda, \infty)$.*

Proof. Let $g_1(t) = g'(t)/t$, $g_2(t) = t^{9-k}g'_1(t)$, $g_3(t) = g'_2(t)/2t$, $g_4(t) = g'_3(t)/2t$, $g_5(t) = g'_4(t)/kt$, $g_6(t) = g'_5(t)/t$, and $g_7(t) = t^{9-k}g'_6(t)$. Then elaborated computations lead to

$$g(1) = 0, \quad (2.5)$$

$$\lim_{t \rightarrow +\infty} g(t) = -\infty, \quad (2.6)$$

$$\begin{aligned} g_1(t) = & -32(k-1)t^{4k-6} + 16(2k-3)t^{4k-8} + (2-k)(3k+2)t^{3k} \\ & + 6k^2t^{3k-2} - 2(k+2)(3k-2)t^{3k-4} + 2(k-4)(3k-4)t^{3k-6} \\ & + 3(k-2)(10-k)t^{3k-8} + 2(k+1)(7-4k)t^{2k} + 4k(4k-1)t^{2k-2} \\ & - 4(k-1)(4k+5)t^{2k-4} + 4(k-2)(4k-1)t^{2k-6} + 2(7-4k) \\ & \times (k-3)t^{2k-8} + (k+2)(10-k)t^k + 2k(k-4)t^{k-2} \\ & - 2(k-2)(k+2)t^{k-4} + 2k(k-4)t^{k-6} + (2-k)(k-6)t^{k-8} + 16, \end{aligned} \quad (2.7)$$

$$g_1(1) = 0, \quad (2.8)$$

$$\lim_{t \rightarrow +\infty} g_1(t) = -\infty, \quad (2.9)$$

$$\begin{aligned} g_2(t) = & -64(k-1)(2k-3)t^{3k+2} + 64(k-2)(2k-3)t^{3k} + 3k(2-k) \\ & \times (3k+2)t^{2k+8} + 6k^2(3k-2)t^{2k+6} - 2(k+2)(3k-2)(3k-4)t^{2k+4} \\ & + 6(k-2)(k-4)(3k-4)t^{2k+2} + 3(k-2)(10-k)(3k-8)t^{2k} \\ & + 4k(k+1)(7-4k)t^{k+8} + 8k(k-1)(4k-1)t^{k+6} - 8(k-1) \\ & \times (k-2)(4k+5)t^{k+4} + 8(k-2)(k-3)(4k-1)t^{k+2} + 4(k-4) \\ & \times (k-3)(7-4k)t^k + k(k+2)(10-k)t^8 + 2k(k-2)(k-4)t^6 \\ & - 2(k-2)(k-4)(k+2)t^4 + 2k(k-4)(k-6)t^2 + (2-k)(k-6)(k-8), \end{aligned} \quad (2.10)$$

$$g_2(1) = 144(5-2k) > 0, \quad (2.11)$$

$$\lim_{t \rightarrow +\infty} g_2(t) = -\infty, \quad (2.12)$$

$$\begin{aligned} g_3(t) = & -32(k-1)(2k-3)(3k+2)t^{3k} + 96k(k-2)(2k-3)t^{3k-2} \\ & + 3k(2-k)(k+4)(3k+2)t^{2k+6} + 6k^2(k+3)(3k-2)t^{2k+4} \\ & - 2(k+2)^2(3k-2)(3k-4)t^{2k+2} + 6(k+1)(k-2)(k-4) \\ & \times (3k-4)t^{2k} + 3k(k-2)(10-k)(3k-8)t^{2k-2} + 2k(k+1) \\ & \times (7-4k)(k+8)t^{k+6} + 4k(k-1)(k+6)(4k-1)t^{k+4} \\ & - 4(k-1)(k-2)(k+4)(4k+5)t^{k+2} + 4(k-2)(k-3)(k+2) \\ & \times (4k-1)t^k + 2k(k-4)(k-3)(7-4k)t^{k-2} + 4k(k+2) \\ & \times (10-k)t^6 + 6k(k-2)(k-4)t^4 - 4(k-2)(k-4)(k+2)t^2 \\ & + 2k(k-4)(k-6), \end{aligned} \quad (2.13)$$

$$g_3(1) = 72(5k - 2)(5 - 2k) > 0, \quad (2.14)$$

$$\lim_{t \rightarrow +\infty} g_3(t) = -\infty, \quad (2.15)$$

$$\begin{aligned} g_4(t) = & -48k(k-1)(2k-3)(3k+2)t^{3k-2} + 48k(k-2)(2k-3) \\ & \times (3k-2)t^{3k-4} + 3k(2-k)(k+3)(k+4)(3k+2)t^{2k+4} \\ & + 6k^2(k+2)(k+3)(3k-2)t^{2k+2} - 2(k+1)(k+2)^2(3k-2) \\ & \times (3k-4)t^{2k} + 6k(k+1)(k-2)(k-4)(3k-4)t^{2k-2} \\ & + 3k(k-1)(k-2)(10-k)(3k-8)t^{2k-4} + k(k+1)(k+6) \end{aligned} \quad (2.16)$$

$$\begin{aligned} & \times (7-4k)(k+8)t^{k+4} + 2k(k-1)(k+4)(k+6)(4k-1)t^{k+2} \\ & - 2(k-1)(k-2)(k+2)(k+4)(4k+5)t^k + 2k(k-2)(k-3) \\ & \times (k+2)(4k-1)t^{k-2} + k(k-2)(k-3)(k-4)(7-4k)t^{k-4} \\ & + 12k(k+2)(10-k)t^4 + 12k(k-2)(k-4)t^2 - 4(k-2)(k-4)(k+2), \end{aligned}$$

$$g_4(1) = 4(-318k^3 + 885k^2 - 210k - 72) = 304.99 \dots > 0, \quad (2.17)$$

$$\lim_{t \rightarrow +\infty} g_4(t) = -\infty, \quad (2.18)$$

$$\begin{aligned} g_5(t) = & -48(k-1)(2k-3)(3k-2)(3k+2)t^{3k-4} + 48(k-2)(2k-3) \\ & \times (3k-2)(3k-4)t^{3k-6} + 6(2-k)(k+2)(k+3)(k+4)(3k+2) \\ & \times t^{2k+2} + 12k(k+1)(k+2)(k+3)(3k-2)t^{2k} - 4(k+1)(k+2)^2 \\ & \times (3k-2)(3k-4)t^{2k-2} + 12(k-1)(k-2)(k-4)(k+1)(3k-4) \\ & \times t^{2k-4} + 6(k-1)(k-2)^2(10-k)(3k-8)t^{2k-6} + (k+1)(k+4) \end{aligned} \quad (2.19)$$

$$\begin{aligned} & \times (k+6)(k+8)(7-4k)t^{k+2} + 2(k-1)(k+2)(k+4)(k+6) \\ & \times (4k-1)t^k - 2(k-1)(k-2)(k+2)(k+4)(4k+5)t^{k-2} \\ & + 2(k-2)^2(k-3)(k+2)(4k-1)t^{k-4} + (k-2)(k-3)(k-4)^2 \\ & \times (7-4k)t^{k-6} + 48(k+2)(10-k)t^2 + 24(k-2)(k-4), \end{aligned}$$

$$g_5(1) = 4(-1038k^3 + 3549k^2 - 3360k + 2196) = 323.50 \dots > 0, \quad (2.20)$$

$$\lim_{t \rightarrow +\infty} g_5(t) = -\infty, \quad (2.21)$$

$$\begin{aligned}
g_6(t) &= -48(k-1)(2k-3)(3k-2)(3k-4)(3k+2)t^{3k-6} + 144(k-2)^2 \\
&\quad \times (2k-3)(3k-2)(3k-4)t^{3k-8} + 12(2-k)(k+1)(k+2) \\
&\quad \times (k+3)(k+4)(3k+2)t^{2k} + 24k^2(k+1)(k+2)(k+3) \\
&\quad \times (3k-2)t^{2k-2} - 8(k-1)(k+1)(k+2)^2(3k-2)(3k-4)t^{2k-4} \\
&\quad + 24(k-1)(k-2)^2(k-4)(k+1)(3k-4)t^{2k-6} + 12(k-1) \\
&\quad \times (k-2)^2(k-3)(10-k)(3k-8)t^{2k-8} + (k+1)(k+2) \\
&\quad \times (k+4)(k+6)(k+8)(7-4k)t^k + 2k(k-1)(k+2)(k+4) \\
&\quad \times (k+6)(4k-1)t^{k-2} - 2(k-1)(k-2)^2(k+2)(k+4) \\
&\quad \times (4k+5)t^{k-4} + 2(k-2)^2(k-3)(k-4)(k+2)(4k-1)t^{k-6} \\
&\quad + (k-2)(k-3)(k-4)^2(k-6)(7-4k)t^{k-8} + 96(k+2)(10-k), \\
g_6(1) &= 4(-3348k^4 + 16233k^3 - 30204k^2 + 28092k - 6768) = -2933.37\dots < 0, \tag{2.22}
\end{aligned}$$

$$\begin{aligned}
g_7(t) &= -144(k-1)(k-2)(2k-3)(3k-2)(3k-4)(3k+2)t^{2k+2} \\
&\quad - 144(k-2)^2(2k-3)(3k-2)(3k-4)(8-3k)t^{2k} - 24k(k-2) \\
&\quad \times (k+1)(k+2)(k+3)(k+4)(3k+2)t^{k+8} + 48k^2(k-1)(k+1) \\
&\quad \times (k+2)(k+3)(3k-2)t^{k+6} - 16(k-1)(k-2)(k+1)(k+2)^2 \\
&\quad \times (3k-2)(3k-4)t^{k+4} + 48(k-1)(k-2)^2(3-k)(4-k)(k+1) \\
&\quad \times (3k-4)t^{k+2} - 24(k-1)(k-2)^2(3-k)(4-k)(10-k)(8-3k) \\
&\quad \times t^k - k(k+1)(k+2)(k+4)(k+6)(k+8)(4k-7)t^8 + 2k \\
&\quad \times (k-1)(k-2)(k+2)(k+4)(k+6)(4k-1)t^6 + 2(k-1) \\
&\quad \times (k-2)^2(4-k)(k+2)(k+4)(4k+5)t^4 - 2(k-2)^2(3-k) \\
&\quad \times (4-k)(6-k)(k+2)(4k-1)t^2 + (k-2)(3-k)(k-4)^2 \\
&\quad \times (6-k)(4k-7)(8-k). \tag{2.23}
\end{aligned}$$

From the expression of $g_7(t)$ and Lemma 2.1, we get

$$\begin{aligned}
g_7(t) &< \left[-144(k-1)(k-2)(2k-3)(3k-2)(3k-4)(3k+2) + 48(k-1) \right. \\
&\quad \times (k-2)^2(3-k)(4-k)(k+1)(3k-4) + 2k(k-1)(k-2)(k+2) \\
&\quad \times (k+4)(k+6)(4k-1) + 2(k-1)(k-2)^2(4-k)(k+2)(k+4) \\
&\quad \left. \times (4k+5) + (k-2)(3-k)(k-4)^2(6-k)(8-k)(4k-7) \right] t^{2k+2} \\
&\quad + k(k+1)(k+2) \left[-24(k-2)(k+3)(k+4)(3k+2)t^{k+8} + 48k \right. \\
&\quad \left. \times (k-1)(k+3)(3k-2)t^{k+6} - (k+4)(k+6)(k+8)(4k-7)t^8 \right]
\end{aligned}$$

$$\begin{aligned}
&= \left(140k^7 - 9353k^6 + 52543k^5 - 103636k^4 + 51700k^3 + 88448k^2 \right. \\
&\quad \left. - 131968k + 54016\right)t^{2k+2} + k(k+1)(k+2) \\
&\quad \times \left[-24(k-2)(k+3)(k+4)(3k+2)t^{k+8} + 48k(k-1)(k+3)(3k-2)t^{k+6} \right. \\
&\quad \left. - (k+4)(k+6)(k+8)(4k-7)t^8\right] \\
&= (-20221.36\dots)t^{2k+2} + k(k+1)(k+2) \\
&\quad \times \left[-24(k-2)(k+3)(k+4)(3k+2)t^{k+8} + 48k(k-1)(k+3)(3k-2)t^{k+6} \right. \\
&\quad \left. - (k+4)(k+6)(k+8)(4k-7)t^8\right] \\
&< 0.
\end{aligned} \tag{2.24}$$

From (2.24), we know that $g_6(t)$ is strictly decreasing in $[1, \infty)$. Then (2.22) implies that $g_5(t)$ is strictly decreasing in $[1, \infty)$.

From (2.20) and (2.21) together with the monotonicity of $g_5(t)$, we clearly see that there exists $\lambda_1 \in (1, \infty)$ such that $g_4(t)$ is strictly increasing in $[1, \lambda_1]$ and strictly decreasing in $[\lambda_1, \infty)$.

Inequality (2.17) and (2.18) together with the piecewise monotonicity of $g_4(t)$ imply that there exists $\lambda_2 \in (1, \infty)$ such that $g_3(t)$ is strictly increasing in $[1, \lambda_2]$ and strictly decreasing in $[\lambda_2, \infty)$.

The piecewise monotonicity of $g_3(t)$ together with (2.14) and (2.15) leads to the fact that there exists $\lambda_3 \in (1, \infty)$ such that $g_2(t)$ is strictly increasing in $[1, \lambda_3]$ and strictly decreasing in $[\lambda_3, \infty)$.

From (2.11) and (2.12) together with the piecewise monotonicity of $g_2(t)$, we conclude that there exists $\lambda_4 \in (1, \infty)$ such that $g_1(t)$ is strictly increasing in $[1, \lambda_4]$ and strictly decreasing in $[\lambda_4, \infty)$.

Equations (2.8) and (2.9) together with the piecewise monotonicity of $g_1(t)$ imply that there exists $\lambda_5 \in (1, \infty)$ such that $g(t)$ is strictly increasing in $[1, \lambda_5]$ and strictly decreasing in $[\lambda_5, \infty)$.

Therefore, Lemma 2.2 follows from (2.5) and (2.6) together with the piecewise monotonicity of $g(t)$. \square

3. Proof of Theorem 1.1

Proof of Theorem 1.1. Without loss of generality, we assume that $a > b$. We first prove that $T(a, b) < H_{5/2}(a, b)$. Let $t = \sqrt[4]{a/b} > 1$, then from (1.1) and (1.2) we have

$$\log T(a, b) - \log H_{5/2}(a, b) = \log \frac{t^4 - 1}{2 \arctan((t^4 - 1)/(t^4 + 1))} - \frac{2}{5} \log \frac{t^{10} + t^5 + 1}{3}. \tag{3.1}$$

Let

$$f(t) = \log \frac{t^4 - 1}{2 \arctan((t^4 - 1)/(t^4 + 1))} - \frac{2}{5} \log \frac{t^{10} + t^5 + 1}{3}. \quad (3.2)$$

Then simple computations lead to

$$\begin{aligned} \lim_{t \rightarrow 1} f(t) &= 0, \\ f'(t) &= \frac{2t^3(2t^6 + t^5 + t + 2)}{(t^4 - 1)(t^{10} + t^5 + 1)(\arctan((t^4 - 1)/(t^4 + 1)))} f_1(t), \end{aligned} \quad (3.3)$$

where $f_1(t) = \arctan((t^4 - 1)/(t^4 + 1)) - 2(t^4 - 1)(t^{10} + t^5 + 1)/(t^8 + 1)(2t^6 + t^5 + t + 2)$. Note that

$$\begin{aligned} \lim_{t \rightarrow 1} f_1(t) &= 0, \\ f_1'(t) &= -\frac{2(t^2 + 1)(t + 1)^2(t - 1)^4}{(1 + t^8)^2(2t^6 + t^5 + t + 2)^2} f_2(t), \end{aligned} \quad (3.4)$$

where

$$\begin{aligned} f_2(t) &= t^{18} + 2t^{17} + 4t^{16} + 6t^{15} - 8t^{12} + 6t^{11} + 21t^{10} + 28t^9 \\ &\quad + 21t^8 + 6t^7 - 8t^6 + 6t^3 + 4t^2 + 2t + 1 > 0 \end{aligned} \quad (3.5)$$

for $t > 1$.

Therefore, $T(a, b) < H_{5/2}(a, b)$ follows from (3.1)–(3.5).

Next, we prove that $T(a, b) > H_{\log 3 / \log(\pi/2)}(a, b)$. Let $k = \log 3 / \log(\pi/2) = 2.43\dots$ and $t = \sqrt{a/b} > 1$, then (1.1) and (1.2) lead to

$$\log T(a, b) - \log H_k(a, b) = \log \frac{t^2 - 1}{2 \arctan((t^2 - 1)/(t^2 + 1))} - \frac{1}{k} \log \frac{t^{2k} + t^k + 1}{3}. \quad (3.6)$$

Let

$$F(t) = \log \frac{t^2 - 1}{2 \arctan((t^2 - 1)/(t^2 + 1))} - \frac{1}{k} \log \frac{t^{2k} + t^k + 1}{3}. \quad (3.7)$$

Then simple computations lead to

$$\lim_{t \rightarrow 1} F(t) = \lim_{t \rightarrow +\infty} F(t) = 0, \quad (3.8)$$

$$F'(t) = \frac{2t^{2k-1} + t^{k+1} + t^{k-1} + 2t}{(t^2 - 1)(t^{2k} + t^k + 1) \arctan((t^2 - 1)/(t^2 + 1))} F_1(t), \quad (3.9)$$

where $F_1(t) = \arctan((t^2 - 1)/(t^2 + 1)) - 2(t^2 - 1)(t^{2k} + t^k + 1)/(t^4 + 1)(2t^{2k-2} + t^k + t^{k-2} + 2)$. Note that

$$\lim_{t \rightarrow 1} F_1(t) = 0, \tag{3.10}$$

$$\lim_{t \rightarrow +\infty} F_1(t) = \frac{\pi}{4} - 1 < 0, \tag{3.11}$$

$$F'(t) = \frac{2t^3}{(t^4 + 1)^2(2t^{2k-2} + t^k + t^{k-2} + 2)^2} F_2(t), \tag{3.12}$$

where

$$\begin{aligned} F_2(t) = & -8t^{4k-4} + 8t^{4k-6} + (2 - k)t^{3k+2} + 2kt^{3k} - 2(k + 2)t^{3k-2} \\ & + 2(k - 4)t^{3k-4} + (10 - k)t^{3k-6} + (7 - 4k)t^{2k+2} \\ & + 2(4k - 1)t^{2k} - 2(4k + 5)t^{2k-2} + 2(4k - 1)t^{2k-4} \\ & + (7 - 4k)t^{2k-6} + (10 - k)t^{k+2} + 2(k - 4)t^k \\ & - 2(k + 2)t^{k-2} + 2kt^{k-4} + (2 - k)t^{k-6} + 8t^2 - 8. \end{aligned} \tag{3.13}$$

From (3.12) and (3.13) together with Lemma 2.2, we clearly see that there exists $\lambda \in (1, \infty)$ such that $F_1(t)$ is strictly increasing in $[1, \lambda]$ and strictly decreasing in $[\lambda, \infty)$.

Equations (3.9)–(3.11) and the piecewise monotonicity of $F_1(t)$ imply that there exists $\mu \in (1, \infty)$ such that $F(t)$ is strictly increasing in $[1, \mu]$ and strictly decreasing in $[\mu, \infty)$. Then from (3.8) we get

$$F(t) > 0 \tag{3.14}$$

for $t > 1$.

Therefore, $T(a, b) > H_{\log 3 / \log(\pi/2)}(a, b)$ follows from (3.6) and (3.7) together with (3.14).

At last, we prove that $H_{\log 3 / \log(\pi/2)}(a, b)$ and $H_{5/2}(a, b)$ are the best possible lower and upper power-type Heron mean bounds for the Seiffert mean $T(a, b)$, respectively.

For any $0 < \varepsilon < k = \log 3 / \log(\pi/2) = 2.43\dots$ and $x > 0$, from (1.1) and (1.2), one has

$$[T(1, 1 + x)]^{5/2-\varepsilon} - [H_{5/2-\varepsilon}(1, 1 + x)]^{5/2-\varepsilon} = \frac{J(x)}{3(2^{5/2-\varepsilon})(\arctan(x/(x + 2)))^{5/2-\varepsilon}}, \tag{3.15}$$

$$\lim_{x \rightarrow +\infty} \frac{H_{k+\varepsilon}(x, 1)}{T(x, 1)} = \frac{\pi}{2} \cdot 3^{-1/(k+\varepsilon)} > \frac{\pi}{2} \cdot 3^{-1/k} = 1, \tag{3.16}$$

where $J(x) = 3x^{5/2-\varepsilon} - [(1 + x)^{5/2-\varepsilon} + (1 + x)^{5/4-\varepsilon/2} + 1][2 \arctan(x/(x + 2))]^{5/2-\varepsilon}$.

Let $x \rightarrow 0$, making use of Taylor extension, we get

$$\begin{aligned}
 J(x) &= 3x^{5/2-\varepsilon} - 2^{5/2-\varepsilon} \left[\frac{x}{2} - \frac{x^2}{4} + \frac{x^3}{12} + o(x^3) \right]^{5/2-\varepsilon} \\
 &\quad \times \left[3 + \left(\frac{15}{4} - \frac{3}{2}\varepsilon \right)x + \left(\frac{13}{8} - \frac{5}{4}\varepsilon \right) \left(\frac{5}{4} - \frac{\varepsilon}{2} \right)x^2 + o(x^2) \right] \quad (3.17) \\
 &= \frac{1}{8}\varepsilon(5-2\varepsilon)x^{9/2-\varepsilon} + o(x^{9/2-\varepsilon}).
 \end{aligned}$$

Equations (3.15) and (3.17) together with inequality (3.16) imply that for any $0 < \varepsilon < \log 3 / \log(\pi/2)$, there exist $\delta = \delta(\varepsilon) > 0$ and $X = X(\varepsilon) > 1$ such that $T(1, 1+x) > H_{5/2-\varepsilon}(1, 1+x)$ for $x \in (0, \delta)$ and $H_{\log 3 / \log(\pi/2)+\varepsilon}(1, x) > T(1, x)$ for $x \in (X, +\infty)$. \square

Acknowledgments

This work was supported by the Natural Science Foundation of China under Grant no. 11071069, the Natural Science Foundation of Zhejiang Province under Grant no. Y7080106, and the Innovation Team Foundation of the Department of Education of Zhejiang Province under Grant no. T200924.

References

- [1] M.-K. Wang, Y.-M. Chu, and Y.-F. Qiu, "Some comparison inequalities for generalized Muirhead and identric means," *Journal of Inequalities and Applications*, vol. 2010, Article ID 295620, 10 pages, 2010.
- [2] B.-Y. Long and Y.-M. Chu, "Optimal inequalities for generalized logarithmic, arithmetic, and geometric means," *Journal of Inequalities and Applications*, vol. 2010, Article ID 806825, 10 pages, 2010.
- [3] B.-Y. Long and Y.-M. Chu, "Optimal power mean bounds for the weighted geometric mean of classical means," *Journal of Inequalities and Applications*, vol. 2010, Article ID 905679, 6 pages, 2010.
- [4] Y.-M. Chu and B.-Y. Long, "Best possible inequalities between generalized logarithmic mean and classical means," *Abstract and Applied Analysis*, vol. 2010, Article ID 303286, 13 pages, 2010.
- [5] Y.-M. Chu, Y.-F. Qiu, M.-K. Wang, and G.-D. Wang, "The optimal convex combination bounds of arithmetic and harmonic means for the Seiffert's mean," *Journal of Inequalities and Applications*, vol. 2010, Article ID 436457, 7 pages, 2010.
- [6] B. Long, W. Xia, and Y. Chu, "An optimal inequality for power mean, geometric mean and harmonic mean," *International Journal of Modern Mathematics*, vol. 5, no. 2, pp. 149–155, 2010.
- [7] Y.-M. Chu and W.-f. Xia, "Two optimal double inequalities between power mean and logarithmic mean," *Computers & Mathematics with Applications*, vol. 60, no. 1, pp. 83–89, 2010.
- [8] X.-M. Zhang, B.-Y. Xi, and Y.-M. Chu, "A new method to prove and find analytic inequalities," *Abstract and Applied Analysis*, vol. 2010, Article ID 128934, 19 pages, 2010.
- [9] X.-M. Zhang and Y.-M. Chu, "A new method to study analytic inequalities," *Journal of Inequalities and Applications*, vol. 2010, Article ID 698012, 13 pages, 2010.
- [10] Y.-M. Chu and W.-f. Xia, "Inequalities for generalized logarithmic means," *Journal of Inequalities and Applications*, vol. 2009, Article ID 763252, 7 pages, 2009.
- [11] M.-y. Shi, Y.-M. Chu, and Y.-p. Jiang, "Optimal inequalities among various means of two arguments," *Abstract and Applied Analysis*, vol. 2009, Article ID 694394, 10 pages, 2009.
- [12] Y.-M. Chu and W.-f. Xia, "Two sharp inequalities for power mean, geometric mean, and harmonic mean," *Journal of Inequalities and Applications*, Article ID 741923, 6 pages, 2009.

- [13] Y. Chu and W. Xia, "Solution of an open problem for Schur convexity or concavity of the Gini mean values," *Science in China A*, vol. 52, no. 10, pp. 2099–2106, 2009.
- [14] Y. Chu and X. Zhang, "Necessary and sufficient conditions such that extended mean values are Schur-convex or Schur-concave," *Journal of Mathematics of Kyoto University*, vol. 48, no. 1, pp. 229–238, 2008.
- [15] Y. Chu, X. Zhang, and G. Wang, "The Schur geometrical convexity of the extended mean values," *Journal of Convex Analysis*, vol. 15, no. 4, pp. 707–718, 2008.
- [16] G. Jia and J. Cao, "A new upper bound of the logarithmic mean," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 4, no. 4, article 80, 2003.
- [17] Z.-H. Zhang, V. Lokesh, and Y.-d. Wu, "The new bounds of the logarithmic mean," *Advanced Studies in Contemporary Mathematics*, vol. 11, no. 2, pp. 185–191, 2005.
- [18] Z.-H. Zhang and Y.-d. Wu, "The generalized Heron mean and its dual form," *Applied Mathematics E-Notes*, vol. 5, pp. 16–23, 2005.
- [19] J. Seiffert, "Aufgabe β 16," *Die Wurzel*, vol. 29, pp. 221–222, 1995.
- [20] P. A. Hästö, "A monotonicity property of ratios of symmetric homogeneous means," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 3, no. 5, article 71, 2002.
- [21] P. S. Bullen, D. S. Mitrinović, and P. M. Vasić, *Means and Their Inequalities*, D. Reidel, the Netherlands, 1988.
- [22] J. Sándor, "A note on some inequalities for means," *Archiv der Mathematik*, vol. 56, no. 5, pp. 471–473, 1991.