

Research Article

Tail Dependence for Regularly Varying Time Series

Ai-Ju Shi^{1,2} and Jin-Guan Lin¹

¹ Department of Mathematics, Southeast University, Nanjing 210096, China

² College of Science, Nanjing University of Posts and Telecommunications, Nanjing 210003, China

Correspondence should be addressed to Jin-Guan Lin, jglin@seu.edu.cn

Received 9 January 2012; Accepted 14 March 2012

Academic Editor: Ming Li

Copyright © 2012 A.-J. Shi and J.-G. Lin. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We use tail dependence functions to study tail dependence for regularly varying (RV) time series. First, tail dependence functions about RV time series are deduced through the intensity measure. Then, the relation between the tail dependence function and the intensity measure is established: they are biuniquely determined. Finally, we obtain the expressions of the tail dependence parameters based on the expectation of the RV components of the time series. These expressions are coincided with those obtained by the conditional probability. Some simulation examples are demonstrated to verify the results we established in this paper.

1. Introduction

Copula is a useful tool for handling multivariate distributions with given univariate margins. A copula C is a distribution function, defined on the unit cube $[0, 1]^d$, with uniform one-dimensional margins U_i . For any $(u_1, \dots, u_d) \in [0, 1]^d$, $C(u_1, \dots, u_d) = P\{U_1 \leq u_1, \dots, U_d \leq u_d\}$; the survival copula is $\hat{C}(u_1, \dots, u_d) = P\{U_1 \geq 1 - u_1, \dots, U_d \geq 1 - u_d\}$, the joint survival function of copula C is $\bar{C}(u_1, \dots, u_d) = \hat{C}(1 - u_1, \dots, 1 - u_d)$. Given a copula C , let

$$F(t_1, \dots, t_d) = C(F_1(t_1), \dots, F_d(t_d)), \quad \text{where } (t_1, \dots, t_d) \in \mathbb{R}^d, \quad (1.1)$$

then F is a multivariate distribution with univariate margins F_1, \dots, F_d . On the other hand, given a distribution F with margins F_1, \dots, F_d , there exists a copula C such that (1.1) holds. And copula C is unique if F_1, \dots, F_d are all continuous (Sklar [1], Nelsen [2]).

In generally, copula forms a natural way to describe the dependence between series when making abstraction of their marginal distributions. Overviews of the probabilistic and statistical properties of copula are to be found in [1–6].

Tail dependence plays an important role among dependence measures due to its ability to describe dependence among extreme values (Frahm et al. [7], Resnick [8, 9], and Nikoloulopoulos et al. [10]) which is introduced by Joe [4]. The issue of tail dependence is mainly for heavy tailed phenomena, heavy tailed phenomenon in fractal time series. It is extensively studied and applied in insurance, risk management, traffic management and engineering management, and so forth. [11–27].

Researchers find various multivariate distributions with heavy tails to describe the extremal or tail dependence, see, Pisarenko and Rodkin [13], Hult and Lindskog [28], and Fang et al. [29]. Many interesting tail quantities have been derived via standard methods: coefficients of tail dependence [30–37] and tail dependence copulas (Charpentier and Segers [38]).

In this paper, we are interested in the tail behavior of the time series X_1, \dots, X_t which have the form:

$$\mathbf{X} = (X_1, \dots, X_t) = (RZ_1, \dots, RZ_t), \quad (1.2)$$

where the scale variable R is independent of random vector (Z_1, \dots, Z_t) . And \mathbf{X} is multivariate regularly varying with distribution function F having copula C .

This distribution is a generalized class, including, for example, multivariate Pareto and multivariate elliptical distribution as special ones. Especially, the multivariate t distribution is included in it. As an example, we will justify the results through multivariate t copula.

In order to analyze the tail dependence behavior of (1.2), we first study the tail dependence functions via intensity measure. Then using the relation between tail dependence parameter and the tail dependence functions, we explore the explicit representations of the tail dependence parameters.

The outline of this paper is as follows. After some preliminaries about multivariate regularly varying series and dependence functions in Section 2, detailed results for the tail dependence functions are discussed in Section 3, the expressions of tail dependence parameters for RV time series are demonstrated in Section 4, and multivariate t distribution is demonstrated as an example in Section 5.

Throughout, (X_1, \dots, X_t) is a random vector with joint distribution function F and copula C . Minima and maxima will be denoted by \wedge and \vee , respectively. The Cartesian product $\prod_{i=1}^t [a_i, b_i]$ is denoted by $[\mathbf{a}, \mathbf{b}]$ for any $\mathbf{a}, \mathbf{b} \in \overline{R}^t$.

2. Preliminaries

Definition 2.1. The t -dimensional random vector \mathbf{X} is said to be regularly varying with index $\alpha \geq 0$ if there exists a random vector Θ with values in \mathbb{S}^{t-1} a.s., where \mathbb{S}^{t-1} denotes the unit sphere in R^d with respect to the norm $|\cdot|$, such that, for all $u > 0$,

$$\frac{P\{|\mathbf{X}| > ux, \mathbf{X}/|\mathbf{X}| \in \cdot\}}{P\{|\mathbf{X}| > x\}} \xrightarrow{v} u^{-\alpha} P\{\Theta \in \cdot\}, \quad (2.1)$$

as $x \rightarrow \infty$. The symbol \xrightarrow{v} stands for vague convergence on \mathbb{S}^{t-1} ; vague convergence of measures is treated in detail in Kallenberg [39]. The distribution of Θ is referred to as the

spectral measure of \mathbf{X} . For further information on multivariate regular variation we refer to Resnick [8, 9].

In fact, (2.1) is equivalent to the following expression

$$nP\{a_n^{-1}\mathbf{X} \in \cdot\} \xrightarrow{v} \mu(\cdot), \quad (2.2)$$

where μ is an intensity measure or Radon measure on $R/\{0\}$ and a_n is a sequence of nonnegative numbers.

From the Definition 2.1, we can see that the regularly varying distribution is connected with intensity measure μ . The following lemma yields the explicit relation between them which can be found in [8].

Lemma 2.2. *Let random vector \mathbf{X} be regularly varying with index $\alpha \geq 0$ and distribution function F , then it is equivalent to the following.*

- (1) *There exists an intensity measure μ on $R^t/\{0\}$, such that for every Borel set $B \subset R^t/\{0\}$ bounded away from the origin that satisfies $\mu(\partial B) = 0$,*

$$\lim_{u \rightarrow \infty} \frac{P\{\mathbf{X} \in uB\}}{P\{|\mathbf{X}| > u\}} = \mu(B), \quad (2.3)$$

with the homogeneous condition $\mu(uB) = u^{-\alpha}\mu(B)$.

- (2) *There exists an intensity measure μ on $R^t/\{0\}$, such that*

$$\lim_{u \rightarrow \infty} \frac{1 - F(ux_1, ux_2, \dots, ux_t)}{1 - F(u, u, \dots, u)} = \frac{P\{\mathbf{X}/u \in [0, \mathbf{x}]^c\}}{P\{\mathbf{X}/u \in [0, 1]^c\}} = \mu([0, \mathbf{x}]^c), \quad (2.4)$$

for all continuous points \mathbf{x} of μ . According to Lemma 2.2, one notices that for any nonnegative multivariate regularly varying random vector \mathbf{X} , its nondegenerate univariate margins X_i have regularly varying right tails and with the same index of \mathbf{X} also, that is,

$$\bar{F}_i(x) = P\{X_i > x\} = x^{-\alpha}L_i(x), \quad x > 0, \quad (2.5)$$

where $L_i(x)$ is a slowly varying function.

Lemma 2.3 (Breiman [40]). *Let ξ and η be two independent nonnegative random variables, η be regularly varying with index α . If there exists a $\gamma > \alpha$, such that $E\xi^\gamma < \infty$, then*

$$P\{\xi\eta > x\} \sim E(\xi^\alpha)P\{\eta > x\}. \quad (2.6)$$

The multivariate version of the Lemma belongs to Basrak et al. [41]. It is said that, if \mathbf{X} is regularly varying in the sense of (2.2), \mathbf{A} is a random $t \times t$ matrix, independent of \mathbf{X} , with $0 < E\|\mathbf{A}\|^\gamma < \infty$ for some $\gamma > \alpha$, then

$$nP\{a_n^{-1}\mathbf{A}\mathbf{X} \in \cdot\} \xrightarrow{v} \tilde{\mu}(\cdot) := E(\mu \circ \mathbf{A}^{-1}(\cdot)), \quad (2.7)$$

where \xrightarrow{v} denotes vague convergence on $\mathbb{R}^t / \{0\}$.

Definition 2.4 (Kluppelberg et al. [42]). Let F be the distribution function of random vector \mathbf{X} with continuous margins $F_i, 1 \leq i \leq t$ and copula C . For any $\mathbf{w} = (w_1, w_2, \dots, w_t) \in \mathbb{R}_+^t$, the lower dependence function is defined as

$$l(\mathbf{w}; C) = \lim_{x \rightarrow 0^+} \frac{C(xw_1, xw_2, \dots, xw_t)}{x}, \quad (2.8)$$

and the upper dependence function is defined as

$$u(\mathbf{w}; C) = \lim_{x \rightarrow 0^+} \frac{\bar{C}(1 - xw_1, 1 - xw_2, \dots, 1 - xw_t)}{x}. \quad (2.9)$$

The upper exponent function is defined as

$$u^*(\mathbf{w}; C) = \sum_{\emptyset \neq S \subset I} (-1)^{|S|-1} u_S(\mathbf{w}_S; C_S), \quad (2.10)$$

where $u_S(\mathbf{w}_S; C_S) = \lim_{x \rightarrow 0^+} \bar{C}(1 - xw_j, \forall j \in S) / x$.

From the definition, we can verify the elementary properties listed in Proposition 2.5 of the tail dependence function. We denote $\tau_J = \lim_{x \rightarrow 1^-} P\{F_j(X_j) > x, \forall j \notin J \mid F_i(X_i) > x, \forall i \in J\}$ and $\xi_J = \lim_{x \rightarrow 0^+} P\{F_j(X_j) < x, \forall j \notin J \mid F_i(X_i) < x, \forall i \in J\}$ are the upper tail and lower dependence parameters of \mathbf{X} , respectively, where J is a nonempty subset of $I = \{1, \dots, t\}$. C_J is the margin of C with component indexes in J .

Proposition 2.5. (1) For any $1 \leq i, j \leq t$,

$$\tau_{ij} = u(1, 1; C_{ij}); \quad \xi_{ij} = l(1, 1; C_{ij}), \quad (2.11)$$

where C_{ij} is the margin copula of X_i, X_j .

(2) For any nonempty $J \subset I$,

$$\tau_J = \frac{u(1, 1, \dots, 1; C)}{u(1, 1, \dots, 1; C_J)}; \quad \xi_J = \frac{l(1, 1, \dots, 1; C)}{l(1, 1, \dots, 1; C_J)}, \quad (2.12)$$

(3)

$$u(\mathbf{w}; C) = \lim_{x \rightarrow 0^+} \frac{\widehat{C}(xw_1, xw_2, \dots, xw_t)}{x} = l(\mathbf{w}; \widehat{C}). \quad (2.13)$$

Proof. (1) According to the definition of τ_{ij} , we get

$$\begin{aligned} \tau_{ij} &= \lim_{x \rightarrow 1^-} P\{F_j(X_j) > x \mid F_i(X_i) > x\} = \lim_{x \rightarrow 0^+} \frac{P\{F_j(X_j) > 1-x, F_i(X_i) > 1-x\}}{P\{F_i(X_i) > 1-x\}} \\ &= \lim_{x \rightarrow 0^+} \frac{\overline{C}_{ij}(1-x, 1-x)}{x} = u(1, 1; C_{ij}); \end{aligned} \quad (2.14)$$

similarly,

$$\begin{aligned} \xi_{ij} &= \lim_{x \rightarrow 0^+} P\{F_j(X_j) < x \mid F_i(X_i) < x\} = \lim_{x \rightarrow 0^+} \frac{P\{F_j(X_j) < x, F_i(X_i) < x\}}{P\{F_i(X_i) < x\}} \\ &= \lim_{x \rightarrow 0^+} \frac{C_{ij}(x, x)}{x} = l(1, 1; C_{ij}). \end{aligned} \quad (2.15)$$

(2) Note that

$$\tau_j = \lim_{x \rightarrow 1^-} \frac{P\{F_j(X_j) > x, \forall j \in I\}}{P\{F_i(X_i) > x, \forall i \in J\}} = \lim_{x \rightarrow 0^+} \frac{\overline{C}(1-x, \dots, 1-x)/x}{\overline{C}_J(1-x, \dots, 1-x)/x} \quad (2.16)$$

combined with (2.9), the first part is determined. The second part can be verified similarly.

(3) We can obtain the proof only paying attention to $\overline{C}(u_1, \dots, u_t) = \widehat{C}(1-u_1, \dots, 1-u_t)$.

From the proposition, the upper tail dependence function of copula C is the lower one of its survival copula \widehat{C} . And in most fractal time series, from the point of view of either theory or applications, people only need to understand the right tail of the data, so we focus on the upper tail function $u(\mathbf{w}; C)$ and coefficient τ_j in the following.

We first study the upper tail dependence function of multivariate regularly varying time series in (1.2) using the intensity measure. \square

3. The Upper Tail Dependence Function for RV Time Series

Theorem 3.1. *Let X_1, \dots, X_t be RV time series with regularly varying index α , distribution function F , copula C , and the stochastic representation as (1.2). If the margins are tail equivalent as $x \rightarrow \infty$, then the upper tail dependence function can be written as*

$$u(\mathbf{w}; C) = \frac{\mu\left(\prod_{i=1}^t [w_i^{-1/\alpha}, \infty)\right)}{\mu\left([1, \infty] \times \overline{R}^{t-1}\right)}, \quad (3.1)$$

and the upper exponent function can be written as

$$u^*(\mathbf{w}; C) = \frac{\mu\left(\left(\prod_{i=1}^t [0, w_i^{-1/\alpha}]\right)^c\right)}{\mu\left(\left([0, 1] \times \bar{R}^{t-1}\right)^c\right)}. \quad (3.2)$$

Proof. For any $\mathbf{w} = (w_1, \dots, w_t) \in \mathbb{R}_+^t$,

$$u(\mathbf{w}; C) = \lim_{x \rightarrow 0^+} \frac{P\{\bar{F}_i(X_i) \leq x w_i, \forall i \in I\}}{P\{\bar{F}_1(X_1) \leq x\}} = \lim_{x \rightarrow 0^+} \frac{P\{X_i > \bar{F}_i^{-1}(x w_i), \forall i \in I\}}{P\{X_1 > \bar{F}_1^{-1}(x)\}}. \quad (3.3)$$

Since every margin F_i is regularly varying with the same index α , we obtain that

$$\bar{F}_i(y) = \frac{L_i(y)}{y^\alpha}, \quad y > 0, \quad (3.4)$$

where $L_i(y)$ is slowly varying function. So for any $w_i > 0$, as $x \rightarrow 0^+$,

$$\bar{F}_i(w_i^{1/\alpha} y) = \frac{L_i(w_i^{1/\alpha} y)}{w_i y^\alpha} = \frac{1}{w_i} \cdot \frac{L_i(w_i^{1/\alpha} y)}{L_i(y)} \cdot \frac{L_i(y)}{y^\alpha} = \frac{1}{w_i} \cdot h_i(w_i, y) \cdot \bar{F}_i(y), \quad (3.5)$$

where $h_i(w_i, y) = L_i(w_i^{1/\alpha} y)/L_i(y) \rightarrow 1$ as $y \rightarrow \infty$. So the equation becomes

$$\bar{F}_i(w_i^{1/\alpha} y) = \frac{1}{w_i} \cdot h_i(w_i, y) \cdot \bar{F}_i(y), \quad (3.6)$$

in other words,

$$w_i^{1/\alpha} y = \bar{F}_i^{-1}\left(\frac{1}{w_i} \bar{F}_i(y) h_i(w_i, y)\right). \quad (3.7)$$

Now we let $\bar{F}_i(y) = x w_i$, then

$$w_i^{1/\alpha} \bar{F}_i^{-1}(x w_i) = \bar{F}_i^{-1}\left(x h_i\left(w_i, \bar{F}_i^{-1}(x w_i)\right)\right), \quad (3.8)$$

so, $\bar{F}_i^{-1}(x w_i) = w_i^{-1/\alpha} \bar{F}_i^{-1}\left(x h_i\left(w_i, \bar{F}_i^{-1}(x w_i)\right)\right)$.

As $x \rightarrow 0^+$, $h_i(w_i, \bar{F}_i^{-1}(x w_i)) \rightarrow 1$, so we get that

$$\bar{F}_i^{-1}(x w_i) \approx w_i^{-1/\alpha} \bar{F}_i^{-1}(x). \quad (3.9)$$

And since the margins are equivalent, that is, $\bar{F}_i(y)/\bar{F}_1(y) \rightarrow 1$ as $y \rightarrow \infty$. We have $\bar{F}_i^{-1}(x)/\bar{F}_1^{-1}(x) \rightarrow 1$ as $x \rightarrow 0^+$ (Resnick [8]). So for sufficient small x , $\bar{F}_i^{-1}(x) \approx \bar{F}_1^{-1}(x)$, and $z = \bar{F}_1^{-1}(x)$, combining (3.3) and (2.3), we obtain that

$$\begin{aligned} u(\mathbf{w}; C) &= \lim_{x \rightarrow 0^+} \frac{P\{X_i > w_i^{-1/\alpha} \bar{F}_i^{-1}(x), \forall i \in I\}}{P\{X_1 > \bar{F}_1^{-1}(x)\}} = \lim_{z \rightarrow \infty} \frac{P\{X_i > w_i^{-1/\alpha} z, \forall i \in I\}}{P\{X_1 > z\}} \\ &= \frac{\mu\left(\prod_{i=1}^t [w_i^{-1/\alpha}, \infty)\right)}{\mu\left([1, \infty] \times \bar{R}^{t-1}\right)}. \end{aligned} \quad (3.10)$$

In order to calculate $u^*(\mathbf{w}; C)$, we recall *the inclusion-exclusion formula*, it says that

$$P\{\cap_{i \in I} A_i\} = \sum_{\emptyset \neq S \subset I} (-1)^{|S|-1} P\{\cup_{j \in S} A_j\} \quad (3.11)$$

is valid for any finite set I and arbitrary events A_i , where $i \in I$.

Using this formula, (2.10) becomes

$$\begin{aligned} u^*(\mathbf{w}; C) &= \lim_{x \rightarrow 0^+} \frac{P\{F_j(X_j) > 1 - xw_j, \exists j \in I\}}{x} = \lim_{x \rightarrow 0^+} \frac{P\{\bar{F}_j(X_j) \leq xw_j, \exists j \in I\}}{P\{\bar{F}_1(X_1) \leq x\}} \\ &= \lim_{x \rightarrow 0^+} \frac{P\{X_j > \bar{F}_j^{-1}(xw_j), \exists j \in I\}}{P\{X_1 > \bar{F}_1^{-1}(x)\}}. \end{aligned} \quad (3.12)$$

By using the same method of (3.3), the following equation holds:

$$u^*(\mathbf{w}; C) = \lim_{z \rightarrow \infty} \frac{P\{X_i > w_i^{-1/\alpha} z, \exists i \in I\}}{P\{X_1 > z\}} = \frac{\mu\left(\left(\prod_{i=1}^t [0, w_i^{-1/\alpha}]\right)^c\right)}{\mu\left([1, \infty] \times \bar{R}^{t-1}\right)}. \quad (3.13)$$

□

Corollary 3.2. *Under the same conditions as Theorem 3.1, the following result holds*

$$\mu\left([1, \infty] \times \bar{R}^{t-1}\right) = \frac{1}{u^*(1, \dots, 1; C)}. \quad (3.14)$$

Proof. By (2.4), one can see that $\mu([0, 1]^c) = 1$. So we can get the result immediately by letting all $w_i = 1$, $1 \leq i \leq t$ in (3.2).

According to Theorem 3.1 and Corollary 3.2, we can represent the intensity measure through the tail dependence function as the following Corollary. □

Corollary 3.3. *Under the same conditions as Theorem 3.1, one has*

$$\begin{aligned}\mu([\mathbf{w}, \infty]) &= \frac{u(w_1^{-\alpha}, \dots, w_t^{-\alpha}; C)}{u^*(1, \dots, 1; C)}, \\ \mu([0, \mathbf{w}]^c) &= \frac{u^*(w_1^{-\alpha}, \dots, w_t^{-\alpha}; C)}{u^*(1, \dots, 1; C)}.\end{aligned}\quad (3.15)$$

4. The Upper Tail Dependence Parameters for Regularly Varying Time Series

According to Proposition 2.5 and Theorem 3.1, we can express the tail dependence parameters by their tail dependence functions. In this section, we will deduce the upper tail dependence parameters of time series with multivariate varying distribution in (1.2) by this method. Hereafter, we let μ be the intensity measure of $\mathbf{R} = (R, R, \dots, R)$ with copula C^R . Where R is regularly varying at ∞ with index α , with survival function $\bar{F}_R(r) = L(r)/r^\alpha$, and $L(\cdot)$ is a slowly varying function. So for any nonnegative vector $\mathbf{w} = (w_1, \dots, w_t)$, we have

$$\mu([0, \mathbf{w}]^c; C^R) = \lim_{r \rightarrow \infty} \frac{P\{R > r \wedge_{i=1}^t w_i\}}{P\{R > r\}} = \lim_{r \rightarrow \infty} \frac{\bar{F}_R(r \wedge_{i=1}^t w_i)}{\bar{F}_R(r)}, \quad (4.1)$$

by inserting $\bar{F}_R(r \wedge_{i=1}^t w_i) = L(r \wedge_{i=1}^t w_i) / (r \wedge_{i=1}^t w_i)^\alpha$ and $\bar{F}_R(r) = L(r) / r^\alpha$ into the representation, then,

$$\mu([0, \mathbf{w}]^c; C^R) = \frac{1}{(\wedge_{i=1}^t w_i)^\alpha} = \left(\bigvee_{i=1}^t w_i \right)^{-\alpha}. \quad (4.2)$$

Similarly, we have,

$$\mu([\mathbf{w}, \infty]; C^R) = \bigwedge_{i=1}^t w_i^{-\alpha}. \quad (4.3)$$

Consequently, we get the main result as follows.

Theorem 4.1. *Let X_1, \dots, X_t be regularly varying time series with the same regularly varying index α and the stochastic representation given in (1.2), the margins are tail equivalent as $x \rightarrow \infty$. If there exists a $\gamma > \alpha$ holds for $0 < E(Z_{i+}^\gamma) < \infty$, then the upper tail dependence parameter of X_1, \dots, X_t is*

$$\tau_J = \frac{E\left(\bigwedge_{i=1}^t (Z_{i+}^\alpha / E(Z_{i+}^\alpha))\right)}{E\left(\bigwedge_{i \in J} (Z_{i+}^\alpha / E(Z_{i+}^\alpha))\right)}. \quad (4.4)$$

Proof. We first calculate the tail dependence function of $\mathbf{X} = (RZ_1, \dots, RZ_t)$. In the following, let C^X and C^Y be the copula of \mathbf{X} and \mathbf{Y} , respectively. Denote

$$(Y_1, \dots, Y_t)^T = \mathbf{A}(R, \dots, R)^T, \quad (4.5)$$

where

$$\mathbf{A} = \text{diag} \left(\frac{Z_{1+}}{(E(Z_{1+}^\alpha))^{1/\alpha}}, \dots, \frac{Z_{t+}}{(E(Z_{t+}^\alpha))^{1/\alpha}} \right). \quad (4.6)$$

Note that $Y_i = (Z_{i+}/(E(Z_{i+}^\alpha))^{1/\alpha})R$ is strictly increasing transformation of $X_i > 0$, for all $i \in I$, and the tail dependence function and the parameter are all copula properties. Hence \mathbf{Y} and \mathbf{X} have the same tail dependence functions. By Lemma 2.3, one can see that the marginal variables Y_i of vector \mathbf{Y} are tail equivalent and regularly varying with the same index as \mathbf{X} as $x \rightarrow \infty$. Denote the intensity measures of \mathbf{Y} and \mathbf{R} by $\tilde{\mu}(\cdot)$ and $\mu(\cdot)$, respectively. According to (2.7),

$$\tilde{\mu}(\cdot) = E\left(\mu\left(\mathbf{A}^{-1}\cdot\right)\right). \quad (4.7)$$

Now by (4.6), we see that,

$$\mathbf{A}^{-1} = \text{diag} \left(\frac{(E(Z_{1+}^\alpha))^{1/\alpha}}{Z_{1+}}, \dots, \frac{(E(Z_{t+}^\alpha))^{1/\alpha}}{Z_{t+}} \right), \quad (4.8)$$

combining this with (4.3), for any nonnegative \mathbf{w} , we obtain the intensity measure given by

$$\begin{aligned} \tilde{\mu}([\mathbf{w}, \infty]) &= E\left(\mu\left(\mathbf{A}^{-1}[\mathbf{w}, \infty]\right)\right) = E\left(\mu\left(\prod_{i=1}^t \left[\frac{(E(Z_{i+}^\alpha))^{1/\alpha}}{Z_{i+}} w_i, \infty\right]; C^R\right)\right) \\ &= E\left(\bigwedge_{i=1}^t \frac{Z_{i+}^\alpha}{E(Z_{i+}^\alpha)} w_i^{-\alpha}\right). \end{aligned} \quad (4.9)$$

Hence, we have

$$\tilde{\mu}\left(\prod_{i=1}^t [w_i^{-\alpha}, \infty]\right) = E\left(\bigwedge_{i=1}^t \frac{Z_{i+}^\alpha}{E(Z_{i+}^\alpha)} w_i\right). \quad (4.10)$$

Substituting this measure into (3.1), we get the upper tail dependence function of vector \mathbf{Y} as follows:

$$u(\mathbf{w}; C^Y) = E\left(\bigwedge_{i=1}^t \frac{Z_{i+}^\alpha}{E(Z_{i+}^\alpha)} w_i\right). \quad (4.11)$$

Since \mathbf{Y} and \mathbf{X} have the same tail dependence functions, we have

$$u(\mathbf{w}; C^{\mathbf{X}}) = E\left(\bigwedge_{i=1}^t \frac{Z_{i+}^{\alpha}}{E(Z_{i+}^{\alpha})} w_i\right). \quad (4.12)$$

By (2) in Proposition 2.5, we obtain the upper tail dependence parameters of vector \mathbf{X} . \square

5. Examples

Let \mathbf{Z} in (1.2) be $\mathbf{Z} = A(U_1, \dots, U_n)$, where A is a $t \times n$ matrix with $AA^T = \Sigma$, and Σ is a $t \times t$ semidefinite matrix, $U = (U_1, \dots, U_n)$ is uniformly distributed on the unit sphere (with respect to Euclidean distance) in R^n . We know that \mathbf{X} conforms to an elliptical contoured distribution (Fang et al. [43]). The tail dependence of the elliptical contoured distribution has been discussed in Schmidt [33]. Here we select the t distribution to display our results in Theorem 4.1 as a special case.

If $\mathbf{X} \sim \mathfrak{t}_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \nu)$, then \mathbf{X} has the stochastic representation ([43]):

$$\mathbf{X} = \boldsymbol{\mu} + \frac{\sqrt{\nu}}{\sqrt{S}} \mathbf{Z}, \quad (5.1)$$

where $S \sim \chi_{\nu}^2$ and $\mathbf{Z} \sim \mathbb{N}_n(\mathbf{0}, \boldsymbol{\Sigma})$ are independent, $\boldsymbol{\mu} \in R^n$.

Let $R = \sqrt{\nu/S}$. Then $R^2 \sim IG(\nu/2, \nu/2)$ and R is regularly varying with index ν at ∞ . So the vector (X_1, \dots, X_n) is regularly varying according to Schmidt [33].

For the upper tail dependence that only relies on the tail behavior of the random vector, we can focus, without loss of generality, on the random vector \mathbf{X} with zero mean vector. Furthermore, since the strictly increasing transformation of (X_1, \dots, X_n) does not change the copula, $\Delta^{-1/2}\mathbf{X}$ has the same copula as \mathbf{X} , where $\boldsymbol{\Sigma} = (\sigma_{ij})$ and $\Delta = \text{diag}(\sigma_{11}, \sigma_{22}, \dots, \sigma_{nn})$. Thus $\Delta^{-1/2}\mathbf{X} \sim \mathfrak{t}_n(0, \Delta^{-1/2}\boldsymbol{\Sigma}\Delta^{-1/2}, \nu)$. It is evident that $\Delta^{-1/2}\boldsymbol{\Sigma}\Delta^{-1/2}$ becomes the correlation matrix of the random vector. Consequently, we may assume that the covariance matrix $\boldsymbol{\Sigma}$ is the correlation matrix. In this situation, all Z_i 's have the same margins as $N(0, 1)$. So $E(Z_{i+}^{\nu})$ are all equal for any $1 \leq i \leq n$. Under these assumptions, using (4.4), we get the upper tail dependence parameter of $\mathfrak{t}_n(0, \boldsymbol{\Sigma}, \nu)$ as

$$\tau_J = \frac{E\left(\bigwedge_{i=1}^n Z_{i+}^{\nu}\right)}{E\left(\bigwedge_{i \in J} Z_{i+}^{\nu}\right)}. \quad (5.2)$$

This is coincided to the one obtained in Shi and Lin [34].

6. Simulations

In Section 4, we obtain the expressions of the tail dependence indexes about RV time series in (1.2). In Section 5, we display our result in the multivariate t distribution as example. In this Section, we will illustrate these results by some Monte Carlo simulated numerical examples.




Figure 1: The estimation of τ_2, τ_{12} under AR(1) (the left one) and EX (the right one) correlation structure.

Given that $\mathbf{y}^{(1)}, \mathbf{y}^{(2)}, \dots, \mathbf{y}^{(m)}$ be generated from the multivariate normal distribution $\mathbf{N}_n(\mathbf{0}, \boldsymbol{\rho})$, then the upper tail dependence indices of $\mathbf{t}_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \nu)$ can be estimated by

$$\hat{\tau}_J = \frac{\sum_{k=1}^m \left(\bigwedge_{j=1}^n |y_j^{(k)}| \right)^\nu I\{\mathbf{y}^{(k)} > 0 \text{ or } \mathbf{y}^{(k)} < 0\}}{\sum_{k=1}^m \left(\bigwedge_{i \in J} |y_i^{(k)}| \right)^\nu I\{\mathbf{y}^{(k)} > 0 \text{ or } \mathbf{y}^{(k)} < 0\}}. \quad (6.1)$$

We estimate the upper tail dependence parameter of 3-dimensional t distribution under autoregressive of order 1 (AR(1)), exchangeable(EX), Toeplitz(TOEP), and unstructured(UN) correlation structure, respectively. For each correlation matrix, we first generate 80,000 pseudorandom vectors, then use (5.2) to estimate tail dependence parameter for different ν . Specifically, we do the following simulations.

$$\boldsymbol{\Sigma}_1 = \begin{pmatrix} 1 & -0.3 & 0.09 \\ -0.3 & 1 & -0.3 \\ 0.09 & -0.3 & 1 \end{pmatrix}, \quad \boldsymbol{\Sigma}_2 = \begin{pmatrix} 1 & -0.3 & -0.3 \\ -0.3 & 1 & -0.3 \\ -0.3 & -0.3 & 1 \end{pmatrix}. \quad (6.2)$$

Let $\mathbf{J} = \{2\}$ and $\{1, 2\}$, respectively. The corresponding upper tail dependence parameters are denoted by τ_2 and τ_{12} . $\boldsymbol{\Sigma}_1$ and $\boldsymbol{\Sigma}_2$ are under AR(1) and EX correlation structure, respectively, the simulated values of τ_2, τ_{12} about different ν are computed and plotted in Figure 1. $\boldsymbol{\Sigma}_3$ and $\boldsymbol{\Sigma}_4$ are under TOEP and UN correlation structure, the corresponding results are demonstrated in Figure 2.

From the two figures, in spite of the correlation structure, τ_j decreased and approached 0 quickly as ν increased to ∞ , which is the tail dependence index for multivariate normal copula.

Many researchers try to discuss the monotonicity of the tail dependence parameter about the regular varying index. Embrechts et al. [11] proved that the tail dependence of the bivariate t distribution is decreasing about the regular varying index ν , and demonstrated




Figure 2: The estimation of τ_2, τ_{12} under TOEP (the left one) and UN (the right one) correlation structure.

that the tail dependence parameter τ_1 is decreasing in ν by numerical results. But From the right graph in Figure 2., these conclusions are not always correct when $t \geq 3$.

$$\sum_3 = \begin{pmatrix} 1 & -0.3 & 0.5 \\ -0.3 & 1 & -0.3 \\ 0.5 & -0.3 & 1 \end{pmatrix}, \quad \sum_4 = \begin{pmatrix} 1 & 0.3 & 0.5 \\ 0.3 & 1 & 0.7 \\ 0.5 & 0.7 & 1 \end{pmatrix}. \quad (6.3)$$

7. Conclusion

In the paper, we mainly study tail dependence of RV time series in (1.2). We use tail dependence function and intensity measure to express tail dependence parameters. Using tail dependence function, we do not need to consider the explicit representation of the copula. We first discuss the tail dependence function of the RV time series due to the propositions of the regularly varying function, connecting the biuniquely determined property between the tail dependence function and the intensity measure. Then we calculate the explicit formula of the upper tail dependence parameter about the RV time series under some conditions. In fact, we can obtain the extreme upper tail dependence index (Shi and Lin [34]) very similarly to Theorem 4.1, for concise, we omit it here.

Copula of continuous variables is invariant under strictly increasing transformation (Nelsen [2]). In order to obtain the tail dependence function of random vector \mathbf{X} , we shift to solve that of \mathbf{Y} in (4.5), which is just a strictly increasing transformation of \mathbf{X} .

At last, we select the t distribution as a special case to display our result, they are coincided to the one given in [34]. The monotonicity of the tail dependence parameters about the regular varying index is still an open problem. Under what constraints the tail dependence parameters will be decreasing in the variation index? We are still interested in the problem. We will discuss it in the following work in details. In engineering application, when we confront fractal time series and seasonal data, we can model the tail dependence property via the tail dependence function if the data is consistent with the constraint conditions in our work.

Acknowledgment

The authors are very grateful to the referees for their comments and suggestions on the earlier version of the paper, which led to a much improved paper. This project is supported by MSRFSEU 3207011102, NSFC 11171065, and NSFJS BK2011058.

References

- [1] M. Sklar, "Fonctions de répartition à n dimensions et leurs marges," vol. 8, pp. 229–231, 1959.
- [2] R. B. Nelsen, *An introduction to Copulas*, Springer Series in Statistics, Springer, New York, NY, USA, 2nd edition, 2006.
- [3] B. Schweizer and A. Sklar, *Probabilistic Metric Spaces*, North-Holland Series in Probability and Applied Mathematics, North-Holland Publishing, New York, NY, USA, 1983.
- [4] H. Joe, *Multivariate Models and dependence concepts*, vol. 73 of *Monographs on Statistics and Applied Probability*, Chapman & Hall, London, 1997.
- [5] C. Genest and A. C Favre, "Everything you always wanted to know about copula modeling but were afraid to ask," *Journal of Hydrological Engineering*, vol. 12, pp. 347–368, 2007.
- [6] C. Klüppelberg and G. Kuhn, "Copula structure analysis," *Journal of the Royal Statistical Society. Series B. Statistical Methodology*, vol. 71, no. 3, pp. 737–753, 2009.
- [7] G. Frahm, M. Junker, and R. Schmidt, "Estimating the tail-dependence coefficient: properties and pitfalls," *Insurance: Mathematics & Economics*, vol. 37, no. 1, pp. 80–100, 2005.
- [8] S. I. Resnick, *Extreme Values, Regular Variation, and Point Processes*, vol. 4 of *Applied Probability. A Series of the Applied Probability Trust*, Springer, New York, NY, USA, 1987.
- [9] S. I. Resnick, *Heavy-Tail Phenomena: Probabilistic and Statistical Modeling*, Springer Series in Operations Research and Financial Engineering, Springer, New York, NY, USA, 2007.
- [10] A. K. Nikoloulopoulos, H. Joe, and H. Li, "Extreme value properties of multivariate t copulas," *Extremes. Statistical Theory and Applications in Science, Engineering and Economics*, vol. 12, no. 2, pp. 129–148, 2009.
- [11] P. Embrechts, F. Lindskog, and A. McNeil, "Modelling dependence with copulas and applications to risk management," in *Handbook of Heavy Tailed Distributions in Finance*, S. Rachev, Ed., Elsevier, New York, NY, USA, 2003.
- [12] P. Embrechts, A. J. McNeil, and D. Straumann, "Correlation and dependence in risk management: properties and pitfalls," in *Risk Management: Value at Risk and Beyond*, M. A. H. Dempster, Ed., pp. 176–223, Cambridge University Press, Cambridge, UK, 1999.
- [13] V. Pisarenko and M. Rodkin, *Heavy-Tailed Distributions in Disaster Analysis*, Springer, 2010.
- [14] E. Kole, K. Koedijk, and M. Verbeek, "Selecting copulas for risk management," Tech. Rep., Erasmus School of Economics and Business Economics, Erasmus University Rotterdam, Rotterdam, The Netherlands, 2006.
- [15] S. Ghosh, "Modelling bivariate rainfall distribution and generating bivariate correlated rainfall data in neighbouring meteorological subdivisions using copula," *Hydrological Processes*, vol. 24, pp. 3558–3567, 2010.
- [16] V. Ganti, K. M. Straub, E. Foufoula-Georgiou, and C. Paola, "Space-time dynamics of depositional systems: experimental evidence and theoretical modeling of heavy-tailed statistics," *Journal of Geophysical Research*, vol. 116, no. F2, Article ID F02011, 17 pages, 2011.
- [17] V. Fasen, "Modeling network traffic by a cluster Poisson input process with heavy and light-tailed file sizes," *Queueing Systems*, vol. 66, no. 4, pp. 313–350, 2010.
- [18] P. Loiseau, P. Goncalves, G. Dewaele, P. Borgnat, P. Abry, and P. Vicat-Blanc, "Investigating self-similarity and heavy-tailed distributions on a large-scale experimental facility," *IEEE/ACM Transactions on Networking*, vol. 18, no. 4, pp. 1261–1274, 2010.
- [19] M. Li and W. Zhao, "Representation of a stochastic traffic bound," *IEEE Transactions on Parallel and Distributed Systems*, vol. 21, no. 9, pp. 1368–1372, 2010.
- [20] P. Abry, P. Borgnat, F. Ricciato, A. Scherrer, and D. Veitch, "Revisiting an old friend: on the observability of the relation between long range dependence and heavy tail," *Telecommunication Systems*, vol. 43, no. 3-4, pp. 147–165, 2010.

- [21] J. A. Hernández, I. W. Phillips, and J. Aracil, "Discrete-time heavy-tailed chains, and their properties in modeling network traffic," *ACM Transactions on Modeling and Computer Simulation*, vol. 17, no. 4, 11 pages, 2007.
- [22] C. Cattani and G. Pierro, "Complexity on acute myeloid leukemia mRNA transcript variant," *Mathematical Problems in Engineering*, Article ID 379873, 16 pages, 2011.
- [23] M. Li, C. Cattani, and S. Y. Chen, "Viewing sea level by a one-dimensional random function with long memory," *Mathematical Problems in Engineering*, vol. 2011, Article ID 654284, 13 pages, 2011.
- [24] M. Li and W. Zhao, "Visiting power laws in cyber-physical networking systems," *Mathematical Problems in Engineering*, vol. 2012, Article ID 302786, 13 pages, 2012.
- [25] S. Y. Chen, H. Tong, Z. Wang, S. Liu, M. Li, and B. Zhang, "Improved generalized belief propagation for vision processing," *Mathematical Problems in Engineering*, Article ID 416963, 12 pages, 2011.
- [26] M. Li, "Fractal time series—a tutorial review," *Mathematical Problems in Engineering*, Article ID 157264, 26 pages, 2010.
- [27] H. Sheng, Y. Chen, and T. Qiu, *Fractional Processes and Fractional-Order Signal Processing*, Signals and Communication Technology, Springer, London, UK, 2012, <http://www.springer.com/engineering/signals/book/978-1-4471-2232-6>.
- [28] H. Hult and F. Lindskog, "Multivariate extremes, aggregation and dependence in elliptical distributions," *Advances in Applied Probability*, vol. 34, no. 3, pp. 587–608, 2002.
- [29] H.-B. Fang, K.-T. Fang, and S. Kotz, "The meta-elliptical distributions with given marginals," *Journal of Multivariate Analysis*, vol. 82, no. 1, pp. 1–16, 2002.
- [30] S. G. Coles, J. Heffernan, and J. A. Tawn, "Dependence measures for multivariate extremes," *Extremes*, vol. 2, pp. 339–365, 1999.
- [31] A. W. Ledford and J. A. Tawn, "Statistics for near independence in multivariate extreme values," *Biometrika*, vol. 83, no. 1, pp. 169–187, 1996.
- [32] M. Mesfioui and J.-F. Quessy, "Dependence structure of conditional Archimedean copulas," *Journal of Multivariate Analysis*, vol. 99, no. 3, pp. 372–385, 2008.
- [33] R. Schmidt, "Tail dependence for elliptically contoured distributions," *Mathematical Methods of Operations Research*, vol. 55, no. 2, pp. 301–327, 2002.
- [34] A. J. Shi and J. G. Lin, "Monotonicity of the tail dependence for multivariate t-copula," *Journal of Southeast University*, vol. 27, no. 4, pp. 466–470, 2011.
- [35] H. Li, "Orthant tail dependence of multivariate extreme value distributions," *Journal of Multivariate Analysis*, vol. 100, no. 1, pp. 243–256, 2009.
- [36] C. Arthur and S. Johan, "Tails of multivariate Archimedean copulas," *Journal of Multivariate Analysis*, vol. 100, no. 7, pp. 1521–1537, 2009.
- [37] G. Frahm, "On the extremal dependence coefficient of multivariate distributions," *Statistics & Probability Letters*, vol. 76, no. 14, pp. 1470–1481, 2006.
- [38] A. Charpentier and J. Segers, "Lower tail dependence for Archimedean copulas: characterizations and pitfalls," *Insurance: Mathematics & Economics*, vol. 40, no. 3, pp. 525–532, 2007.
- [39] O. Kallenberg, *Random Measures*, Academic Press, Berlin, Germany, 3rd edition, 1983.
- [40] L. Breiman, "On some limit theorems similar to the arc-sin law," *Theory of Probability and Its Applications*, vol. 10, pp. 323–331, 1965.
- [41] B. Basrak, R. A. Davis, and T. Mikosch, "Regular variation of GARCH processes," *Stochastic Processes and their Applications*, vol. 99, no. 1, pp. 95–116, 2002.
- [42] C. Kluppelberg, G. Kuhn, and L. Peng, "Multivariate tail copula: modeling and estimation," Tech. Rep. number 468, Center for Mathematical Sciences, Munich University of Technology, 2006, <http://epub.ub.uni-muenchen.de/>.
- [43] K. T. Fang, S. Kotz, and K. W. Ng, *Symmetric Multivariate and Related Distributions*, vol. 36 of *Monographs on Statistics and Applied Probability*, Chapman & Hall, London, UK, 1990.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

