



**ON ASYMPTOTIC CONSTANTS RELATED TO PRODUCTS OF  
BERNOULLI NUMBERS AND FACTORIALS**

**Bernd C. Kellner**

*Mathematisches Institut, Universität Göttingen, Bunsenstr. 3-5, 37073 Göttingen,  
Germany*

bk@bernoulli.org

*Received: 9/24/07, Accepted: 2/21/09*

**Abstract**

We discuss the asymptotic expansions of certain products of Bernoulli numbers and factorials, e.g.,

$$\prod_{\nu=1}^n |B_{2\nu}| \quad \text{and} \quad \prod_{\nu=1}^n (k\nu)!^{\nu^r} \quad \text{as } n \rightarrow \infty$$

for integers  $k \geq 1$  and  $r \geq 0$ . Our main interest is to determine exact expressions, in terms of known constants, for the asymptotic constants of these expansions and to show some relations among them.

**1. Introduction**

Let  $B_n$  be the  $n$ th Bernoulli number. These numbers are defined by

$$\frac{z}{e^z - 1} = \sum_{n=0}^{\infty} B_n \frac{z^n}{n!}, \quad |z| < 2\pi$$

where  $B_n = 0$  for odd  $n > 1$ . The Riemann zeta function  $\zeta(s)$  is defined by

$$\zeta(s) = \sum_{\nu=1}^{\infty} \nu^{-s} = \prod_p (1 - p^{-s})^{-1}, \quad s \in \mathbb{C}, \operatorname{Re} s > 1. \quad (1)$$

By Euler's formula we have for even positive integers  $n$  that

$$\zeta(n) = -\frac{1}{2} \frac{(2\pi i)^n}{n!} B_n. \quad (2)$$

Products of Bernoulli numbers occur in certain contexts in number theory. For example, the Minkowski–Siegel mass formula states that, for positive integers  $n$  with  $8 \mid n$ ,

$$M(n) = \frac{|B_k|}{2k} \prod_{\nu=1}^{k-1} \frac{|B_{2\nu}|}{4\nu}, \quad n = 2k,$$

which describes the mass of the genus of even unimodular positive definite  $n \times n$  matrices (for details see [12, p. 252]). We introduce the following constants which we shall need further on.

**Lemma 1.** *There exist the constants*

$$\begin{aligned} \mathcal{C}_1 &= \prod_{\nu=2}^{\infty} \zeta(\nu) = 2.2948565916\dots, \\ \mathcal{C}_2 &= \prod_{\nu=1}^{\infty} \zeta(2\nu) = 1.8210174514\dots, \\ \mathcal{C}_3 &= \prod_{\nu=1}^{\infty} \zeta(2\nu + 1) = 1.2602057107\dots. \end{aligned}$$

*Proof.* We have  $\log(1 + x) < x$  for real  $x > 0$ . Then

$$\log \prod_{\nu=1}^{\infty} \zeta(2\nu) = \sum_{\nu=1}^{\infty} \log \zeta(2\nu) < \sum_{\nu=1}^{\infty} (\zeta(2\nu) - 1) = \frac{3}{4}. \tag{3}$$

The last sum of (3) is well known and follows by rearranging in geometric series, since we have absolute convergence. We then obtain that  $\pi^2/6 < \mathcal{C}_2 < e^{3/4}$ ,  $\zeta(3) < \mathcal{C}_3 < \mathcal{C}_2$ , and  $\mathcal{C}_1 = \mathcal{C}_2\mathcal{C}_3$ .  $\square$

To compute the infinite products above within a given precision, one can use the following arguments. A standard estimate for the partial sum of  $\zeta(s)$  is given by

$$\zeta(s) - \sum_{\nu=1}^N \nu^{-s} < \frac{N^{1-s}}{s-1}, \quad s \in \mathbb{R}, \quad s > 1.$$

This follows by comparing the sum of  $\nu^{-s}$  and the integral of  $x^{-s}$  in the interval  $(N, \infty)$ . Now, one can estimate the number  $N$  depending on  $s$  and the needed precision. However, we use a computer algebra system, that computes  $\zeta(s)$  to a given precision with already accelerated built-in algorithms. Since  $\zeta(s) \rightarrow 1$  monotonically as  $s \rightarrow \infty$ , we next have to determine a finite product that suffices the precision. From above, we obtain

$$\zeta(s) - 1 < 2^{-s} \left( 1 + \frac{2}{s-1} \right), \quad s \in \mathbb{R}, \quad s > 1. \tag{4}$$

According to (3) and (4), we then get an estimate for the remainder of the infinite product by

$$\log \prod_{\nu > N'} \zeta(\nu) < 2^{-N'+\varepsilon}$$

where we can take  $\varepsilon = 3/N'$ ; the choice of  $\varepsilon$  follows by  $2^x \geq 1 + x \log 2$  and (4).

We give the following example where the constant  $\mathcal{C}_1$  plays an important role; see Finch [8]. Let  $a(n)$  be the number of non-isomorphic abelian groups of order  $n$ . The constant  $\mathcal{C}_1$  equals the average of the numbers  $a(n)$  by taking the limit. Thus, we have

$$\mathcal{C}_1 = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N a(n).$$

By definition the constant  $\mathcal{C}_2$  is connected with values of the Riemann zeta function on the positive real axis. Moreover, this constant is also connected with values of the Dedekind eta function

$$\eta(\tau) = e^{\pi i \tau / 12} \prod_{\nu=1}^{\infty} (1 - e^{2\pi i \nu \tau}), \quad \tau \in \mathbb{C}, \operatorname{Im} \tau > 0$$

on the upper imaginary axis.

**Lemma 2.** *The constant  $\mathcal{C}_2$  is given by*

$$1/\mathcal{C}_2 = \prod_p p^{\frac{1}{12}} \eta\left(i \frac{\log p}{\pi}\right)$$

where the product runs over all primes.

*Proof.* By Lemma 1 and the Euler product (1) of  $\zeta(s)$ , we obtain

$$\mathcal{C}_2 = \prod_{\nu=1}^{\infty} \prod_p (1 - p^{-2\nu})^{-1} = \prod_p \prod_{\nu=1}^{\infty} (1 - p^{-2\nu})^{-1}$$

where we can change the order of the products because of absolute convergence. Rewriting  $p^{-2\nu} = e^{2\pi i \nu \tau}$  with  $\tau = i \log p / \pi$  yields the result.  $\square$

We used MATHEMATICA [17] to compute all numerical values in this paper. The values were checked again by increasing the needed precision to 10 more digits.

## 2. Preliminaries

We use the notation  $f \sim g$  for real-valued functions when  $\lim_{x \rightarrow \infty} f(x)/g(x) = 1$ . As usual,  $O(\cdot)$  denotes Landau's symbol. We write  $\log f$  for  $\log(f(x))$ .

**Definition 3.** Define the linear function spaces

$$\Omega_n = \operatorname{span} \{x^\nu, x^\nu \log x\}, \quad n \geq 0$$

over  $\mathbb{R}$  where  $f \in \Omega_n$  is a function  $f : \mathbb{R}^+ \rightarrow \mathbb{R}$ . Let

$$\Omega_\infty = \bigcup_{n \geq 0} \Omega_n.$$

Define the linear map  $\psi : \Omega_\infty \rightarrow \mathbb{R}$  which gives the constant term of any  $f \in \Omega_\infty$ . For the class of functions

$$F(x) = f(x) + O(x^{-\delta}), \quad f \in \Omega_n, \quad n \geq 0, \quad \delta > 0 \tag{5}$$

define the linear operator  $[\ ] : C(\mathbb{R}^+; \mathbb{R}) \rightarrow \Omega_\infty$  such that  $[F] = f$  and  $[F] \in \Omega_n$ . Then  $\psi([F])$  is defined to be the asymptotic constant of  $F$ .

We shall examine functions  $h : \mathbb{N} \rightarrow \mathbb{R}$  which grow exponentially; in particular these functions are represented by certain products. Our problem is to find an asymptotic function  $\tilde{h} : \mathbb{R}^+ \rightarrow \mathbb{R}$  where  $h \sim \tilde{h}$ . If  $F = \log \tilde{h}$  satisfies (5), then we have  $[\log \tilde{h}] \in \Omega_n$  for a suitable  $n$  and we identify  $[\log \tilde{h}] = [\log h] \in \Omega_n$  in that case.

**Lemma 4.** *Let  $f \in \Omega_n$  where*

$$f(x) = \sum_{\nu=0}^n (\alpha_\nu x^\nu + \beta_\nu x^\nu \log x)$$

*with coefficients  $\alpha_\nu, \beta_\nu \in \mathbb{R}$ . Let  $g(x) = f(\lambda x)$  with a fixed  $\lambda \in \mathbb{R}^+$ . Then  $g \in \Omega_n$  and  $\psi(g) = \psi(f) + \beta_0 \log \lambda$ .*

*Proof.* Since  $g(x) = f(\lambda x)$  we obtain

$$g(x) = \sum_{\nu=0}^n (\alpha_\nu (\lambda x)^\nu + \beta_\nu (\lambda x)^\nu (\log \lambda + \log x)).$$

This shows that  $g \in \Omega_n$ . The constant terms are  $\alpha_0$  and  $\beta_0 \log \lambda$ , and thus  $\psi(g) = \psi(f) + \beta_0 \log \lambda$ . □

**Definition 5.** For a function  $f : \mathbb{R}^+ \rightarrow \mathbb{R}$  we introduce the notation

$$f(x) = \sum'_{\nu \geq 1} f_\nu(x)$$

with functions  $f_\nu : \mathbb{R}^+ \rightarrow \mathbb{R}$  in case  $f$  has a divergent series expansion such that

$$f(x) = \sum_{\nu=1}^{m-1} f_\nu(x) + \theta_m(x) f_m(x), \quad \theta_m(x) \in (0, 1), \quad m \geq N_f,$$

where  $N_f$  is a suitable constant depending on  $f$ .

Next we need some well-known facts which we state without proof (see [10]).

**Proposition 6.** *Let*

$$H_0 = 0, \quad H_n = \sum_{\nu=1}^n \frac{1}{\nu}, \quad n \geq 1$$

*be the  $n$ th harmonic number. These numbers satisfy  $H_n = \gamma + \log n + O(n^{-1})$ ,  $n \geq 1$ , where  $\gamma = 0.5772156649\dots$  is Euler's constant.*

**Proposition 7.** (Stirling's series) *The Gamma function  $\Gamma(x)$  has the divergent series expansion*

$$\log \Gamma(x + 1) = \frac{1}{2} \log(2\pi) + \left(x + \frac{1}{2}\right) \log x - x + \sum'_{\nu \geq 1} \frac{B_{2\nu}}{2\nu(2\nu - 1)} x^{-(2\nu-1)}, \quad x > 0.$$

**Remark 8.** When evaluating the divergent series given above, we have to choose a suitable index  $m$  such that

$$\sum'_{\nu \geq 1} \frac{B_{2\nu}}{2\nu(2\nu - 1)} x^{-(2\nu-1)} = \sum_{\nu=1}^{m-1} \frac{B_{2\nu}}{2\nu(2\nu - 1)} x^{-(2\nu-1)} + \theta_m(x)R_m(x)$$

and the remainder  $|\theta_m(x)R_m(x)|$  is as small as possible. Since  $\theta_m(x) \in (0, 1)$  is not effectively computable in general, we have to use  $|R_m(x)|$  instead as an error bound. Schäfke and Finsterer [15], among others, showed that the so-called Lindelöf error bound  $L = 1$  for the estimate  $L \geq \theta_m(x)$  is best possible for positive real  $x$ .

**Proposition 9.** *If  $\alpha \in \mathbb{R}$  with  $0 \leq \alpha < 1$ , then*

$$\prod_{\nu=1}^n (\nu - \alpha) = \frac{\Gamma(n + 1 - \alpha)}{\Gamma(1 - \alpha)} \sim \frac{\sqrt{2\pi}}{\Gamma(1 - \alpha)} \left(\frac{n}{e}\right)^n n^{\frac{1}{2} - \alpha} \quad \text{as } n \rightarrow \infty.$$

Euler's formula for the Gamma function states the following.

**Proposition 10.** (Euler) *Let  $\Gamma(x)$  be the Gamma function. Then*

$$\prod_{\nu=1}^{n-1} \Gamma\left(\frac{\nu}{n}\right) = \frac{(2\pi)^{\frac{n-1}{2}}}{\sqrt{n}}.$$

**Proposition 11.** (Glaisher [9], Kinkelin [11]) *As  $n \rightarrow \infty$ ,*

$$\prod_{\nu=1}^n \nu^\nu \sim \mathcal{A} n^{\frac{1}{2}n(n+1) + \frac{1}{12}} e^{-\frac{n^2}{4}},$$

*where  $\mathcal{A} = 1.2824271291\dots$  is the Glaisher–Kinkelin constant, which is given by*

$$\log \mathcal{A} = \frac{1}{12} - \zeta'(-1) = \frac{\gamma}{12} + \frac{1}{12} \log(2\pi) - \frac{\zeta'(2)}{2\pi^2}.$$

Numerous digits of the decimal expansion of the Glaisher–Kinkelin constant  $\mathcal{A}$  are recorded as sequence A074962 in OEIS [16].

### 3. Products of Factorials

In this section we consider products of factorials and determine their asymptotic expansions and constants. For these asymptotic constants we derive a divergent series representation as well as a closed formula.

**Theorem 12.** *Let  $k$  be a positive integer. Asymptotically, we have*

$$\prod_{\nu=1}^n (k\nu)! \sim \mathcal{F}_k \mathcal{A}^k (2\pi)^{\frac{1}{4}} \left(\frac{kn}{e^{3/2}}\right)^{\frac{k}{2}n(n+1)} \left(2\pi k e^{k/2-1} n\right)^{\frac{n}{2}} n^{\frac{1}{4} + \frac{k}{12} + \frac{1}{12k}} \quad \text{as } n \rightarrow \infty$$

with certain constants  $\mathcal{F}_k$  which satisfy

$$\log \mathcal{F}_k = \frac{\gamma}{12k} + \sum_{j \geq 2}' \frac{B_{2j} \zeta(2j-1)}{2j(2j-1)k^{2j-1}}.$$

Moreover, the constants have the asymptotic behavior that

$$\lim_{k \rightarrow \infty} \mathcal{F}_k = 1, \quad \lim_{k \rightarrow \infty} \mathcal{F}_k^k = e^{\gamma/12}, \quad \text{and} \quad \prod_{k=1}^n \mathcal{F}_k \sim \mathcal{F}_\infty n^{\gamma/12} \quad \text{as } n \rightarrow \infty$$

with

$$\log \mathcal{F}_\infty = \frac{\gamma^2}{12} + \sum_{j \geq 2}' \frac{B_{2j} \zeta(2j-1)^2}{2j(2j-1)}.$$

**Theorem 13.** *If  $k$  is a positive integer, then*

$$\log \mathcal{F}_k = -\left(k + \frac{1}{k}\right) \log \mathcal{A} + \frac{1}{12k} - \frac{1}{12k} \log k + \frac{k}{4} \log(2\pi) - \sum_{\nu=1}^{k-1} \frac{\nu}{k} \log \Gamma\left(\frac{\nu}{k}\right).$$

We will prove Theorem 13 later, since we shall need several preliminaries.

*Proof of Theorem 12.* Let  $k \geq 1$  be fixed. By Stirling’s approximation, see Proposition 7, we have

$$\log(k\nu)! = \frac{1}{2} \log(2\pi) + \left(k\nu + \frac{1}{2}\right) \log(k\nu) - k\nu + f(k\nu) \tag{6}$$

where we can write the remaining divergent sum as

$$f(k\nu) = \frac{1}{12k\nu} + \sum_{j \geq 2}' \frac{B_{2j}}{2j(2j-1)(k\nu)^{2j-1}}.$$

Define  $S(n) = 1 + \dots + n = n(n+1)/2$ . By summation we obtain

$$\begin{aligned} \sum_{\nu=1}^n \log(k\nu)! &= \frac{n}{2} \log(2\pi k) + \frac{1}{2} \log n! - kS(n) + kS(n) \log k \\ &\quad + k \sum_{\nu=1}^n \nu \log \nu + \sum_{\nu=1}^n f(k\nu). \end{aligned}$$

The term  $\frac{1}{2} \log n!$  is evaluated again by (6). Proposition 11 provides that

$$k \sum_{\nu=1}^n \nu \log \nu = k \log \mathcal{A} + kS(n) \log n + \frac{k}{12} \log n - \frac{k}{2} \left( S(n) - \frac{n}{2} \right) + O(n^{-\delta})$$

with some  $\delta > 0$ . Since  $\lim_{n \rightarrow \infty} H_n - \log n = \gamma$ , we asymptotically obtain for the remaining sum that

$$\lim_{n \rightarrow \infty} \left( \sum_{\nu=1}^n f(k\nu) - \frac{1}{12k} \log n \right) = \frac{\gamma}{12k} + \sum_{j \geq 2} \frac{B_{2j} \zeta(2j-1)}{2j(2j-1) k^{2j-1}} =: \log \mathcal{F}_k. \quad (7)$$

Here we have used the following arguments. We choose a fixed index  $m > 2$  for the remainder of the divergent sum. Then

$$\lim_{n \rightarrow \infty} \sum_{\nu=1}^n \theta_m(k\nu) \frac{B_{2m}}{2m(2m-1) (k\nu)^{2m-1}} = \eta_m \frac{B_{2m} \zeta(2m-1)}{2m(2m-1) k^{2m-1}} \quad (8)$$

with some  $\eta_m \in (0, 1)$ , since  $\theta_m(k\nu) \in (0, 1)$  for all  $\nu \geq 1$ . Thus, we can write (7) as an asymptotic series again. Collecting all terms, we finally get the asymptotic formula

$$\begin{aligned} \sum_{\nu=1}^n \log(k\nu)! &= \log \mathcal{F}_k + k \log \mathcal{A} + \frac{1}{4} \log(2\pi) + kS(n) \left( -\frac{3}{2} + \log(kn) \right) \\ &\quad + \frac{n}{2} \left( \log(2\pi k) + \frac{k}{2} - 1 + \log n \right) \\ &\quad + \left( \frac{1}{4} + \frac{k}{12} + \frac{1}{12k} \right) \log n + O(n^{-\delta'}) \end{aligned}$$

with some  $\delta' > 0$ . Note that the exact value of  $\delta'$  does not play a role here. Now, let  $k$  be an arbitrary positive integer. From (7) we deduce that

$$\log \mathcal{F}_k = \frac{\gamma}{12k} + O(k^{-3}) \quad \text{and} \quad k \log \mathcal{F}_k = \frac{\gamma}{12} + O(k^{-2}). \quad (9)$$

The summation of (7) yields

$$\sum_{k=1}^n \log \mathcal{F}_k = \frac{\gamma}{12} H_n + \sum_{k=1}^n \sum_{j \geq 2} \frac{B_{2j} \zeta(2j-1)}{2j(2j-1) k^{2j-1}}. \quad (10)$$

Similar to (7) and (8), we can write again:

$$\lim_{n \rightarrow \infty} \left( \sum_{k=1}^n \log \mathcal{F}_k - \frac{\gamma}{12} \log n \right) = \frac{\gamma^2}{12} + \sum_{j \geq 2} \frac{B_{2j} \zeta(2j-1)^2}{2j(2j-1)} =: \log \mathcal{F}_\infty. \quad (11)$$

□

The case  $k = 1$  of Theorem 12 is related to the so-called Barnes  $G$ -function (see [2]). Now we shall determine exact expressions for the constants  $\mathcal{F}_k$ . For  $k \geq 2$  this is more complicated.

**Lemma 14.** *We have  $\mathcal{F}_1 = (2\pi)^{\frac{1}{4}} e^{\frac{1}{12}} / \mathcal{A}^2$ .*

*Proof.* Writing down the product of  $n!$  repeatedly in  $n + 1$  rows, one observes by counting in rows and columns that

$$n!^{n+1} = \prod_{\nu=1}^n \nu! \prod_{\nu=1}^n \nu^\nu. \quad (12)$$

From Proposition 7 we have

$$(n + 1) \log n! = \frac{n + 1}{2} \log(2\pi) - n(n + 1) + (n + 1) \left( n + \frac{1}{2} \right) \log n + \frac{1}{12} + O(n^{-1}).$$

Comparing the asymptotic constants of both sides of (12) when  $n \rightarrow \infty$ , we obtain

$$(2\pi)^{\frac{1}{2}} e^{\frac{1}{12}} = \mathcal{F}_1 \mathcal{A} (2\pi)^{\frac{1}{4}} \cdot \mathcal{A}$$

where the right side follows by Theorem 12 and Proposition 11. □

**Proposition 15.** *Let  $k, l$  be integers with  $k \geq 1$ . Define*

$$F_{k,l}(n) := \prod_{\nu=1}^n (k\nu - l)! \quad \text{for } 0 \leq l < k.$$

*Then  $[\log F_{k,l}] \in \Omega_2$  and  $F_{k,0}(n) \cdots F_{k,k-1}(n) = F_{1,0}(kn)$ . Moreover*

$$F_{k,l}(n)/F_{k,l+1}(n) = k^n \prod_{\nu=1}^n \left( \nu - \frac{l}{k} \right) \quad \text{for } 0 \leq l < k - 1$$

*and  $[\log(F_{k,l}/F_{k,l+1})] = [\log F_{k,l}] - [\log F_{k,l+1}] \in \Omega_1$ .*

*Proof.* We deduce the proposed products from  $(k\nu - l)!/(k\nu - (l + 1))! = k\nu - l$  and

$$\prod_{\nu=1}^n (k\nu)! (k\nu - 1)! \cdots (k\nu - (k - 1))! = \prod_{\nu=1}^{kn} \nu!. \quad (13)$$





our equations to  $\mathbb{R}$  by applying the linear map  $\psi$ , since we are only interested in the asymptotic constants. We obtain the linear system of equations

$$M_k \cdot x = b, \quad x, b \in \mathbb{R}^k$$

where

$$x = (\psi([\log F_{k,0}]), \dots, \psi([\log F_{k,k-1}]))^T$$

and  $b = (b_1, \dots, b_k)^T$  with

$$b_{l+1} = \psi([\log(F_{k,l}/F_{k,l+1})]) = \frac{1}{2} \log(2\pi) - \log \Gamma\left(1 - \frac{l}{k}\right) \quad \text{for } l = 0, \dots, k-2$$

using Proposition 9. The last element  $b_k$  is given by Theorem 12, Lemma 14, and Lemma 4:

$$\begin{aligned} b_k &= \psi([\log(F_{1,0}(kn))]) = \frac{1}{4} \log(2\pi) + \log \mathcal{F}_1 + \log \mathcal{A} + \frac{5}{12} \log k \\ &= \frac{1}{2} \log(2\pi) - \log \mathcal{A} + \frac{1}{12} + \frac{5}{12} \log k. \end{aligned}$$

By Lemma 16 we can solve the linear system directly with

$$x = \frac{1}{k} \widetilde{M}_k \cdot b.$$

The first row yields

$$x_1 = \frac{1}{k} b_k + \frac{1}{k} \sum_{\nu=1}^{k-1} (k - \nu) b_\nu.$$

On the other side, we have

$$x_1 = \psi([\log F_{k,0}]) = \log \mathcal{F}_k + \frac{1}{4} \log(2\pi) + k \log \mathcal{A}.$$

This provides

$$\begin{aligned} \log \mathcal{F}_k &= - \left(k + \frac{1}{k}\right) \log \mathcal{A} + \left(\frac{k}{4} + \frac{1}{2k} - \frac{1}{2}\right) \log(2\pi) \\ &\quad + \frac{5}{12k} \log k + \frac{1}{12k} - \sum_{\nu=2}^{k-1} \frac{\nu-1}{k} \log \Gamma\left(\frac{\nu}{k}\right) \end{aligned} \tag{15}$$

after some rearranging of terms. By Euler's formula, see Proposition 10, we have

$$\frac{1}{k} \sum_{\nu=1}^{k-1} \log \Gamma\left(\frac{\nu}{k}\right) = \left(\frac{1}{2} - \frac{1}{2k}\right) \log(2\pi) - \frac{1}{2k} \log k. \tag{16}$$

Finally, substituting (16) into (15) yields the result. □

**Remark 17.** Although the formula for  $\mathcal{F}_k$  has an elegant short form, one might also use (15) instead, since this formula omits the value  $\Gamma(1/k)$ . Thus we easily obtain the value of  $\mathcal{F}_2$  from (15) at once:  $\mathcal{F}_2 = (2\pi)^{\frac{1}{4}} 2^{\frac{5}{24}} e^{\frac{1}{24}} / \mathcal{A}^{\frac{5}{2}}$ .

**Corollary 18.** *Asymptotically, we have*

$$\prod_{\nu=1}^{n-1} \Gamma\left(\frac{\nu}{n}\right)^\nu \sim \frac{e^{\frac{1-\gamma}{12}}}{\mathcal{A}} \left(\frac{(2\pi)^{\frac{1}{4}}}{\mathcal{A}}\right)^{n^2} / n^{\frac{1}{12}} \quad \text{as } n \rightarrow \infty$$

with the constants  $e^{\frac{1-\gamma}{12}} / \mathcal{A} = 0.8077340270\dots$  and  $(2\pi)^{\frac{1}{4}} / \mathcal{A} = 1.2345601953\dots$

*Proof.* On the one hand, we have by (9) that

$$n \log \mathcal{F}_n = \frac{\gamma}{12} + O(n^{-2}).$$

On the other hand, Theorem 13 provides that

$$n \log \mathcal{F}_n = -(n^2 + 1) \log \mathcal{A} + \frac{1}{12} - \frac{1}{12} \log n + \frac{n^2}{4} \log(2\pi) - \sum_{\nu=1}^{n-1} \nu \log \Gamma\left(\frac{\nu}{n}\right).$$

Combining both formulas easily gives the result. □

Since we have derived exact expressions for the constants  $\mathcal{F}_k$ , we can improve the calculation of  $\mathcal{F}_\infty$ . The divergent sum of  $\mathcal{F}_\infty$ , given in Theorem 12, is not suitable to determine a value within a given precision, but we can use this sum in a modified way. Note that we cannot use the limit formula

$$\log \mathcal{F}_\infty = \lim_{n \rightarrow \infty} \left( \sum_{k=1}^n \log \mathcal{F}_k - \frac{\gamma}{12} \log n \right)$$

without a very extensive calculation, because the sequence  $\gamma_n = H_n - \log n$  converges too slowly. Moreover, the computation of  $\mathcal{F}_k$  involves the computation of the values  $\Gamma(\nu/k)$ . This becomes more difficult for larger  $k$ .

**Proposition 19.** *Let  $m, n$  be positive integers. Assume that  $m > 2$  and the constants  $\mathcal{F}_k$  are given by exact expressions for  $k = 1, \dots, n$ . Define the computable values  $\eta_k \in (0, 1)$  implicitly by*

$$\log \mathcal{F}_k = \frac{\gamma}{12k} + \sum_{j=2}^{m-1} \frac{B_{2j} \zeta(2j-1)}{2j(2j-1) k^{2j-1}} + \eta_k \frac{B_{2m} \zeta(2m-1)}{2m(2m-1) k^{2m-1}}.$$

Then

$$\log \mathcal{F}_\infty = \frac{\gamma^2}{12} + \sum_{j=2}^{m-1} \frac{B_{2j} \zeta(2j-1)^2}{2j(2j-1)} + \theta_{n,m} \frac{B_{2m} \zeta(2m-1)^2}{2m(2m-1)}$$

with  $\theta_{n,m} \in (\theta_{n,m}^{\min}, \theta_{n,m}^{\max}) \subset (0, 1)$  where

$$\theta_{n,m}^{\min} = \zeta(2m-1)^{-1} \sum_{k=1}^n \frac{\eta_k}{k^{2m-1}}, \quad \theta_{n,m}^{\max} = 1 + \zeta(2m-1)^{-1} \sum_{k=1}^n \frac{\eta_k - 1}{k^{2m-1}}.$$

The error bound for the remainder of the divergent sum of  $\log \mathcal{F}_\infty$  is given by

$$\theta_{n,m}^{\text{err}} = \left( 1 - \zeta(2m-1)^{-1} \sum_{k=1}^n \frac{1}{k^{2m-1}} \right) \frac{|B_{2m}| \zeta(2m-1)^2}{2m(2m-1)}.$$

*Proof.* Let  $n \geq 1$  and  $m > 2$  be fixed integers. The divergent sums for  $\log \mathcal{F}_k$  and  $\log \mathcal{F}_\infty$  are given by Theorem 12. Since we require exact expressions for  $\mathcal{F}_k$ , we can compute the values  $\eta_k$  for  $k = 1, \dots, n$ . We define

$$\eta_{m,k} = \eta'_{m,k} = \eta_k \quad \text{for } k = 1, \dots, n$$

and

$$\eta_{m,k} = 0, \quad \eta'_{m,k} = 1 \quad \text{for } k > n.$$

We use (10) and (11) to derive the bounds:

$$\theta_{n,m}^{\min} = \zeta(2m-1)^{-1} \sum_{k=1}^{\infty} \frac{\eta_{m,k}}{k^{2m-1}} < \theta_{n,m} < \zeta(2m-1)^{-1} \sum_{k=1}^{\infty} \frac{\eta'_{m,k}}{k^{2m-1}} = \theta_{n,m}^{\max}.$$

We obtain the suggested formulas for  $\theta_{n,m}^{\min}$  and  $\theta_{n,m}^{\max}$  by evaluating the sums with  $\eta_{m,k} = 0$ , resp.  $\eta'_{m,k} = 1$ , for  $k > n$ . The error bound is given by the difference of the absolute values of the minimal and maximal remainder. Therefore

$$\theta_{n,m}^{\text{err}} = (\theta_{n,m}^{\max} - \theta_{n,m}^{\min})R = \left( 1 - \zeta(2m-1)^{-1} \sum_{k=1}^n \frac{1}{k^{2m-1}} \right) R$$

with  $R = |B_{2m}| \zeta(2m-1)^2 / 2m(2m-1)$ . □

**Result 20.** Exact expressions for  $\mathcal{F}_k$ :

$$\begin{aligned} \mathcal{F}_1 &= (2\pi)^{\frac{1}{4}} e^{\frac{1}{12}} / \mathcal{A}^2, & \mathcal{F}_2 &= (2\pi)^{\frac{1}{4}} 2^{\frac{5}{24}} e^{\frac{1}{24}} / \mathcal{A}^{\frac{5}{2}}, \\ \mathcal{F}_3 &= (2\pi)^{\frac{5}{12}} 3^{\frac{5}{36}} e^{\frac{1}{36}} / \mathcal{A}^{\frac{10}{3}} \Gamma\left(\frac{2}{3}\right)^{\frac{1}{3}}, & \mathcal{F}_4 &= (2\pi)^{\frac{1}{2}} 2^{\frac{1}{3}} e^{\frac{1}{48}} / \mathcal{A}^{\frac{17}{4}} \Gamma\left(\frac{3}{4}\right)^{\frac{1}{2}}. \end{aligned}$$

We have computed the constants  $\mathcal{F}_k$  by their exact expression. Moreover, we have determined the index  $m$  of the smallest remainder of their asymptotic divergent series and the resulting error bound given by Theorem 12.

Constant	Value	$m$	Error bound
$\mathcal{F}_1$	1.04633506677050318098...	4	$6.000 \cdot 10^{-4}$
$\mathcal{F}_2$	1.02393741163711840157...	7	$7.826 \cdot 10^{-7}$
$\mathcal{F}_3$	1.01604053706462099128...	10	$1.198 \cdot 10^{-9}$
$\mathcal{F}_4$	1.01204589802394464624...	13	$1.948 \cdot 10^{-12}$
$\mathcal{F}_5$	1.00963997283647705086...	16	$3.272 \cdot 10^{-15}$
$\mathcal{F}_6$	1.00803362724207326544...	20	$5.552 \cdot 10^{-18}$

The weak interval of  $\mathcal{F}_\infty$  is given by Theorem 12. The second value is derived by Proposition 19 with parameters  $m = 17$  and  $n = 7$ . Thus, exact expressions of  $\mathcal{F}_1, \dots, \mathcal{F}_7$  are needed to compute  $\mathcal{F}_\infty$  within the given precision.

Constant	Value / Interval	$m$	Error bound
$\mathcal{F}_\infty$	(1.02428, 1.02491)	4	$6.050 \cdot 10^{-4}$
$\mathcal{F}_\infty$	1.0246068826559721480...	17	$6.321 \cdot 10^{-22}$

#### 4. Products of Bernoulli Numbers

Using results of the previous sections, we are now able to consider several products of Bernoulli numbers and to derive their asymptotic expansions and constants.

**Theorem 21.** *Asymptotically, we have*

$$\prod_{\nu=1}^n |B_{2\nu}| \sim \mathcal{B}_1 \left( \frac{n}{\pi e^{3/2}} \right)^{n(n+1)} (16\pi n)^{\frac{n}{2}} n^{\frac{11}{24}} \text{ as } n \rightarrow \infty,$$

$$\prod_{\nu=1}^n \frac{|B_{2\nu}|}{2\nu} \sim \mathcal{B}_2 \left( \frac{n}{\pi e^{3/2}} \right)^{n^2} \left( \frac{4n}{\pi e} \right)^{\frac{n}{2}} / n^{\frac{1}{24}} \text{ as } n \rightarrow \infty$$

with the constants

$$\mathcal{B}_1 = \mathcal{C}_2 \mathcal{F}_2 \mathcal{A}^2 (2\pi)^{\frac{1}{4}} = \mathcal{C}_2 (2\pi)^{\frac{1}{2}} 2^{\frac{5}{24}} e^{\frac{1}{24}} / \mathcal{A}^{\frac{1}{2}},$$

$$\mathcal{B}_2 = \mathcal{C}_2 \mathcal{F}_2 \mathcal{A}^2 / (2\pi)^{\frac{1}{4}} = \mathcal{C}_2 2^{\frac{5}{24}} e^{\frac{1}{24}} / \mathcal{A}^{\frac{1}{2}}.$$

*Proof.* By Euler’s formula (2) for  $\zeta(2\nu)$  and Lemma 1 we obtain

$$\prod_{\nu=1}^n |B_{2\nu}| \sim \mathcal{C}_2 \prod_{\nu=1}^n \frac{2 \cdot (2\nu)!}{(2\pi)^{2\nu}} \sim \mathcal{C}_2 2^n (2\pi)^{-n(n+1)} \prod_{\nu=1}^n (2\nu)! \text{ as } n \rightarrow \infty.$$

Theorem 12 states for  $k = 2$  that

$$\prod_{\nu=1}^n (2\nu)! \sim \mathcal{F}_2 \mathcal{A}^2 (2\pi)^{\frac{1}{4}} \left( \frac{2n}{e^{3/2}} \right)^{n(n+1)} (4\pi n)^{\frac{n}{2}} n^{\frac{11}{24}} \text{ as } n \rightarrow \infty.$$

The expression for  $\mathcal{F}_2$  is given in Remark 17. Combining both asymptotic formulas above gives the first suggested formula. It remains to evaluate the following product:

$$\prod_{\nu=1}^n (2\nu) = 2^n n! \sim (2\pi)^{\frac{1}{2}} \left(\frac{2n}{e}\right)^n n^{\frac{1}{2}} \quad \text{as } n \rightarrow \infty.$$

After some rearranging of terms we then obtain the second suggested formula.  $\square$

**Remark 22.** Milnor and Husemoller [14, pp. 49–50] give the following asymptotic formula without proof:

$$\prod_{\nu=1}^n |B_{2\nu}| \sim \mathcal{B}' n! 2^{n+1} F(2n+1) \quad \text{as } n \rightarrow \infty \tag{17}$$

where

$$F(n) = \left(\frac{n}{2\pi e^{3/2}}\right)^{\frac{n^2}{4}} \left(\frac{8\pi e}{n}\right)^{\frac{n}{4}} / n^{\frac{1}{24}} \tag{18}$$

and  $\mathcal{B}' \approx 0.705$  is a certain constant. This constant is related to the constant  $\mathcal{B}_2$ .

**Proposition 23.** *The constant  $\mathcal{B}'$  is given by*

$$\mathcal{B}' = 2^{\frac{1}{24}} 2^{-\frac{3}{2}} \mathcal{B}_2 = \mathcal{C}_2 e^{\frac{1}{24}} / 2^{\frac{5}{4}} \mathcal{A}^{\frac{1}{2}} = 0.7048648734\dots$$

*Proof.* By Theorem 21 we have

$$\prod_{\nu=1}^n \frac{|B_{2\nu}|}{2\nu} \sim \mathcal{B}_2 G(n) \quad \text{as } n \rightarrow \infty \tag{19}$$

with

$$G(n) = \left(\frac{n}{\pi e^{3/2}}\right)^{n^2} \left(\frac{4n}{\pi e}\right)^{\frac{n^2}{2}} / n^{\frac{1}{24}}.$$

We observe that (17) and (19) are equivalent so that

$$2 \mathcal{B}' F(2n+1) \sim \mathcal{B}_2 G(n) \quad \text{as } n \rightarrow \infty.$$

We rewrite (18) in the suitable form

$$F(2n+1) = \left(\frac{n+\frac{1}{2}}{\pi e^{3/2}}\right)^{n^2+n+\frac{1}{4}} \left(\frac{4\pi e}{n+\frac{1}{2}}\right)^{\frac{n^2}{2}+\frac{1}{4}} / 2^{\frac{1}{24}} \left(n+\frac{1}{2}\right)^{\frac{1}{24}}.$$

Hence, we easily deduce that

$$G(n)/F(2n+1) = \left(1+\frac{1}{2n}\right)^{-n^2-\frac{n}{2}+\frac{1}{24}} e^{\frac{n}{2}} 2^{\frac{1}{24}} \left(\frac{e^{1/2}}{4}\right)^{\frac{1}{4}}.$$

It is well known that

$$\lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n = e^x \quad \text{and} \quad \lim_{n \rightarrow \infty} e^{-xn} \left(1 + \frac{x}{n}\right)^{n^2} = e^{-\frac{x^2}{2}}.$$

Evaluating the asymptotic terms, we get

$$2\mathcal{B}'/\mathcal{B}_2 \sim G(n)/F(2n+1) \sim e^{\frac{1}{8}} e^{-\frac{1}{4}} 2^{\frac{1}{24}} e^{\frac{1}{8}} 2^{-\frac{1}{2}} \quad \text{as } n \rightarrow \infty,$$

which finally yields  $\mathcal{B}' = 2^{\frac{1}{24}} 2^{-\frac{3}{2}} \mathcal{B}_2$ . □

**Theorem 24.** *The Minkowski–Siegel mass formula asymptotically states for positive integers  $n$  with  $4 \mid n$  that*

$$M(2n) = \frac{|B_n|}{2n} \prod_{\nu=1}^{n-1} \frac{|B_{2\nu}|}{4\nu} \sim \mathcal{B}_3 \left(\frac{n}{\pi e^{3/2}}\right)^{n^2} / \left(\frac{4n}{\pi e}\right)^{\frac{n}{2}} n^{\frac{1}{24}} \quad \text{as } n \rightarrow \infty$$

with  $\mathcal{B}_3 = \sqrt{2} \mathcal{B}_2$ .

*Proof.* Let  $n$  always be even. By Proposition 7 and (2) we have

$$2^{-n} \left| \frac{B_n/n}{B_{2n}/2n} \right| = 2 \frac{\zeta(n)}{\zeta(2n)} \pi^n \frac{n!}{(2n)!} \sim \sqrt{2} \left(\frac{4n}{\pi e}\right)^{-n} \quad \text{as } n \rightarrow \infty,$$

since  $\zeta(n)/\zeta(2n) \sim 1$  and

$$\log \left(\frac{n!}{(2n)!}\right) \sim n - n \log n - \left(2n + \frac{1}{2}\right) \log 2 \quad \text{as } n \rightarrow \infty.$$

We finally use Theorem 21 and (19) to obtain

$$M(2n) = 2^{-n} \left| \frac{B_n/n}{B_{2n}/2n} \right| \prod_{\nu=1}^n \frac{|B_{2\nu}|}{2\nu} \sim \sqrt{2} \mathcal{B}_2 \left(\frac{4n}{\pi e}\right)^{-n} G(n) \quad \text{as } n \rightarrow \infty,$$

which gives the result. □

**Result 25.** The constants  $\mathcal{B}'$ ,  $\mathcal{B}_\nu$  ( $\nu = 1, 2, 3$ ) mainly depend on the constant  $\mathcal{C}_2$  and the Glaisher–Kinkelin constant  $\mathcal{A}$ .

Constant	Expression	Value
$\mathcal{A}$		1.28242712910062263687...
$\mathcal{C}_2$		1.82101745149929239040...
$\mathcal{B}_1$	$\mathcal{C}_2 (2\pi)^{\frac{1}{2}} 2^{\frac{5}{24}} e^{\frac{1}{24}} / \mathcal{A}^{\frac{1}{2}}$	4.85509664652226751252...
$\mathcal{B}_2$	$\mathcal{C}_2 2^{\frac{5}{24}} e^{\frac{1}{24}} / \mathcal{A}^{\frac{1}{2}}$	1.93690332773294192068...
$\mathcal{B}_3$	$\mathcal{C}_2 2^{\frac{17}{24}} e^{\frac{1}{24}} / \mathcal{A}^{\frac{1}{2}}$	2.73919495508550621998...
$\mathcal{B}'$	$\mathcal{C}_2 e^{\frac{1}{24}} / 2^{\frac{3}{4}} \mathcal{A}^{\frac{1}{2}}$	0.70486487346802031057...

**5. Generalizations**

In this section we derive a generalization of Theorem 12. The results show the structure of the constants  $\mathcal{F}_k$  and the generalized constants  $\mathcal{F}_{r,k}$ , which we shall define later, in a wider context. For simplification we introduce the following definitions which arise from the Euler-Maclaurin summation formula.

The sum of consecutive integer powers is given by the well-known formula

$$\sum_{\nu=0}^{n-1} \nu^r = \frac{B_{r+1}(n) - B_{r+1}}{r + 1} = \sum_{j=0}^r \binom{r}{j} B_{r-j} \frac{n^{j+1}}{j + 1}, \quad r \geq 0$$

where  $B_m(x)$  is the  $m$ th Bernoulli polynomial. Now, the Bernoulli number  $B_1 = -\frac{1}{2}$  is responsible for omitting the last power  $n^r$  in the summation above. Because we further need the summation up to  $n^r$ , we change the sign of  $B_1$  in the sum as follows:

$$S_r(n) = \sum_{\nu=1}^n \nu^r = \sum_{j=0}^r \binom{r}{j} (-1)^{r-j} B_{r-j} \frac{n^{j+1}}{j + 1}, \quad r \geq 0.$$

This modification also coincides with

$$\zeta(-n) = (-1)^{n+1} \frac{B_{n+1}}{n + 1}$$

for nonnegative integers  $n$ . We define the extended sum

$$S_r(n; f(\diamond)) = \sum_{j=0}^r \binom{r}{j} (-1)^{r-j} B_{r-j} \frac{n^{j+1} f(j + 1)}{j + 1}, \quad r \geq 0$$

where the symbol  $\diamond$  is replaced by the index  $j + 1$  in the sum. Note that  $S_r$  is linear in the second parameter, i.e.,

$$S_r(n; \alpha + \beta f(\diamond)) = \alpha S_r(n) + \beta S_r(n; f(\diamond)).$$

Finally we define

$$D_k(x) = \sum_{j \geq 1} \widehat{B}_{2j,k} x^{-(2j-1)} \quad \text{where} \quad \widehat{B}_{m,k} = \frac{B_m}{m(m-1)k^{m-1}}.$$

**Theorem 26.** *Let  $r$  be a nonnegative integer. Then*

$$\prod_{\nu=1}^n \nu^{\nu^r} \sim \mathcal{A}_r Q_r(n) \quad \text{as} \quad n \rightarrow \infty,$$



where  $\mathcal{A}_r$  is the generalized Glaisher–Kinkelin constant defined by

$$\log \mathcal{A}_r = -\zeta(-r) H_r - \zeta'(-r).$$

Moreover,  $\log Q_r \in \Omega_{r+1}$  with

$$\log Q_r(n) = (S_r(n) - \zeta(-r)) \log n + S_r(n; H_r - H_\diamond).$$

*Proof.* This formula and the constants easily follow from a more general formula for real  $r > -1$  given in [10, 9.28, p. 595] and after some rearranging of terms.  $\square$

**Remark 27.** The case  $r = 0$  reduces to Stirling’s approximation of  $n!$  with  $\mathcal{A}_0 = \sqrt{2\pi}$ . The case  $r = 1$  gives the usual Glaisher–Kinkelin constant  $\mathcal{A}_1 = \mathcal{A}$ . The expression  $S_r(n; H_r - H_\diamond)$  does not depend on the definition of  $B_1$ , since the term with  $B_1$  is cancelled in the sum. Graham, Knuth, and Patashnik [10, 9.28, p. 595] notice that the constant  $-\zeta'(-r)$  has been determined in a book of de Bruijn [7, §3.7] in 1970. The theorem above has a long history. In 1894 Alexeiewsky [3] gave the identity

$$\prod_{\nu=1}^n \nu^{\nu^r} = \exp(\zeta'(-r, n+1) - \zeta'(-r))$$

where  $\zeta'(s, a)$  is the partial derivative of the Hurwitz zeta function with respect to the first variable. Between 1903 and 1913, Ramanujan recorded in his notebooks [5, Entry 27, pp. 273–276] (the first part was published and edited by Berndt [5] in 1985) an asymptotic expansion for real  $r > -1$  and an analytic expression for the constant  $C_r = -\zeta'(-r)$ . However, Ramanujan only derived closed expressions for  $C_0$  and  $C_{2r}$  ( $r \geq 1$ ) in terms of  $\zeta(2r+1)$ ; see (28) below. In 1933 Bendersky [4] showed that there exist certain constants  $\mathcal{A}_r$ . Since 1980, several others have investigate the asymptotic formula, including MacLeod [13], Choudhury [6], and Adamchik [1, 2].

**Theorem 28.** Let  $k, r$  be integers with  $k \geq 1$  and  $r \geq 0$ , then

$$\prod_{\nu=1}^n (k\nu)!^{\nu^r} \sim \mathcal{F}_{r,k} \mathcal{A}_r^{\frac{1}{2}} \mathcal{A}_{r+1}^k P_{r,k}(n) Q_r(n)^{\frac{1}{2}} Q_{r+1}(n)^k \quad \text{as } n \rightarrow \infty.$$

The constants  $\mathcal{F}_{r,k}$  and functions  $P_{r,k}$  satisfy that  $\lim_{k \rightarrow \infty} \mathcal{F}_{r,k} = 1$  and  $\log P_{r,k} \in \Omega_{r+2}$  where

$$\begin{aligned} \log P_{r,k}(n) = & \frac{1}{2} S_r(n) \log(2\pi k) + k S_{r+1}(n) \log(k/e) \\ & + \widehat{B}_{r+2,k} \log n + \sum_{j=1}^{\lfloor \frac{r+1}{2} \rfloor} \widehat{B}_{2j,k} S_{r+1-2j}(n). \end{aligned}$$

The constants  $\mathcal{A}_r$  and functions  $Q_r$  are defined as in Theorem 26.

The determination of exact expressions for the constants  $\mathcal{F}_{r,k}$  seems to be a very complicated and extensive task in the case  $r > 0$ . The next theorem gives a partial result for  $k = 1$  and  $r \geq 0$ .

**Theorem 29.** *Let  $r$  be a nonnegative integer, then*

$$\log \mathcal{F}_{r,1} = \frac{1}{2} \log \mathcal{A}_r - \log \mathcal{A}_{r+1} + S_r(1; \widehat{B}_{1+\diamond,1} - \log \mathcal{A}_\diamond).$$

Case  $r = 0$ :

$$\log \mathcal{F}_{r,1} = \frac{1}{12} + \frac{1}{2} \log \mathcal{A}_0 - 2 \log \mathcal{A}_1.$$

Case  $r > 0$ :

$$\log \mathcal{F}_{r,1} = \alpha_{r,0} + \sum_{j=1}^{r+1} \alpha_{r,j} \log \mathcal{A}_j$$

where

$$\alpha_{r,j} = \begin{cases} \frac{B_{r+1}}{2r(r+1)}, & r \not\equiv j \pmod{2}, \quad j = 0; \\ \sum_{j=0}^r \binom{r}{j} \frac{B_{r-j} B_{j+2}}{(j+1)^2(j+2)}, & r \equiv j \pmod{2}, \quad j = 0; \\ -\delta_{r+1,j} - \binom{r+1}{j} \frac{B_{r+1-j}}{r+1}, & r \not\equiv j \pmod{2}, \quad j > 0; \\ 0, & r \equiv j \pmod{2}, \quad j > 0 \end{cases}$$

and  $\delta_{i,j}$  is Kronecker's delta.

*Proof of Theorem 28.* Let  $k$  and  $r$  be fixed. We extend the proof of Theorem 12. From (6) we have

$$\log(k\nu)! = \frac{1}{2} \log(2\pi k) + k\nu \log\left(\frac{k}{e}\right) + \left(k\nu + \frac{1}{2}\right) \log \nu + D_k(\nu). \tag{20}$$

The summation yields

$$\sum_{\nu=1}^n \nu^r \log(k\nu)! = F_1(n) + F_2(n) + F_3(n)$$

where

$$F_1(n) = \frac{1}{2} S_r(n) \log(2\pi k) + k S_{r+1}(n) \log(k/e),$$

$$F_2(n) = k \sum_{\nu=1}^n \nu^{r+1} \log \nu + \frac{1}{2} \sum_{\nu=1}^n \nu^r \log \nu,$$

$$F_3(n) = \sum_{\nu=1}^n \nu^r D_k(\nu).$$

Theorem 26 provides

$$F_2(n) = k(\log \mathcal{A}_{r+1} + \log Q_{r+1}(n)) + \frac{1}{2}(\log \mathcal{A}_r + \log Q_r(n)) + O(n^{-\delta})$$

with some  $\delta > 0$ . Let  $R = \lfloor \frac{r+1}{2} \rfloor$ . By definition we have

$$x^r D_k(x) = \sum_{j=1}^R \widehat{B}_{2j,k} x^{r+1-2j} + \sum'_{j>R} \widehat{B}_{2j,k} x^{r+1-2j} =: E_1(x) + E_2(x).$$

Therewith we obtain that

$$F_3(n) = \sum_{j=1}^R \widehat{B}_{2j,k} S_{r+1-2j}(n) + \sum_{\nu=1}^n E_2(\nu).$$

For the second sum above we consider two cases. We use similar arguments which we have applied to (7) and (8). If  $r$  is odd, then

$$\lim_{n \rightarrow \infty} \sum_{\nu=1}^n E_2(\nu) = \sum'_{j>R} \widehat{B}_{2j,k} \zeta(2j - (r + 1)). \tag{21}$$

Note that  $\widehat{B}_{r+2,k} = 0$  in that case. If  $r$  is even, then we have to take care of the term  $\nu^{-1}$ . This gives

$$\lim_{n \rightarrow \infty} \left( \sum_{\nu=1}^n E_2(\nu) - \widehat{B}_{r+2,k} \log n \right) = \gamma \widehat{B}_{r+2,k} + \sum'_{j>R+1} \widehat{B}_{2j,k} \zeta(2j - (r + 1)). \tag{22}$$

The right hand side of (21), resp. (22), defines the constant  $\log \mathcal{F}_{r,k}$ . Finally we have to collect all results for  $F_1$ ,  $F_2$ , and  $F_3$ . This gives the constants and the function  $P_{r,k}$ . It remains to show that  $\lim_{k \rightarrow \infty} \log \mathcal{F}_{r,k} = 0$ . This follows by  $\widehat{B}_{2j,k} \rightarrow 0$  as  $k \rightarrow \infty$ .  $\square$

The following lemma gives a generalization of Equation (12) in Lemma 14. After that we can give a proof of Theorem 29.

**Lemma 30.** *Let  $n, r$  be integers with  $n \geq 1$  and  $r \geq 0$ . Then*

$$n!^{S_r(n)} \prod_{\nu=1}^n \nu^{\nu^r} = \prod_{\nu=1}^n \nu!^{\nu^r} \prod_{\nu=1}^n \nu^{S_r(\nu)}. \tag{23}$$

*Proof.* We regard the following enumeration scheme which can be easily extended to  $n$  rows and  $n$  columns:

$$\begin{array}{ccc} 1^{1^r} & \boxed{2^{1^r}} & \boxed{3^{1^r}} \\ 1^{2^r} & 2^{2^r} & \boxed{3^{2^r}} \\ 1^{3^r} & 2^{3^r} & 3^{3^r} \end{array}$$

The product of all elements, resp. non-framed elements, in the  $\nu$ th row equals  $n!^{\nu^r}$ , resp.  $\nu!^{\nu^r}$ . The product of the framed elements in the  $\nu$ th column equals  $\nu^{S_r(\nu-1)}$ . Thus

$$n!^{S_r(n)} = \prod_{\nu=1}^n \nu!^{\nu^r} \prod_{\nu=1}^n \nu^{S_r(\nu)-\nu^r}.$$

□

*Proof of Theorem 29.* Let  $r \geq 0$ . We take the logarithm of (23) to obtain

$$F_1(n) + F_2(n) = F_3(n) + F_4(n) \tag{24}$$

where

$$\begin{aligned} F_1(n) &= S_r(n) \log n!, & F_2(n) &= \sum_{\nu=1}^n \nu^r \log \nu, \\ F_3(n) &= \sum_{\nu=1}^n \nu^r \log \nu!, & F_4(n) &= \sum_{\nu=1}^n S_r(\nu) \log \nu. \end{aligned}$$

Next we consider the asymptotic expansions  $\tilde{F}_j$  of the functions  $F_j$  ( $j = 1, \dots, 4$ ) when  $n \rightarrow \infty$ . We further reduce the functions  $\tilde{F}_j$  via the maps

$$C(\mathbb{R}^+; \mathbb{R}) \xrightarrow{[\cdot]} \Omega_\infty \xrightarrow{\psi} \mathbb{R}$$

to the constant terms which are the asymptotic constants of  $[\tilde{F}_j]$  in  $\Omega_\infty$ . Consequently (24) turns into

$$\psi([\tilde{F}_1]) + \psi([\tilde{F}_2]) = \psi([\tilde{F}_3]) + \psi([\tilde{F}_4]). \tag{25}$$

We know from Theorem 26 and Theorem 28 that

$$\psi([\tilde{F}_2]) = \log \mathcal{A}_r \quad \text{and} \quad \psi([\tilde{F}_3]) = \log \mathcal{F}_{r,1} + \frac{1}{2} \log \mathcal{A}_r + \log \mathcal{A}_{r+1}.$$

For  $\tilde{F}_4$  we derive the expression

$$\psi([\tilde{F}_4]) = S_r(1; \log \mathcal{A}_\circ), \tag{26}$$

since each term  $s_j \nu^j$  in  $S_r(\nu)$  produces the term  $s_j \log \mathcal{A}_j$ . It remains to evaluate  $\tilde{F}_1$ .

According to (20) we have

$$\log n! = \frac{1}{2} \log(2\pi) - n + \left(n + \frac{1}{2}\right) \log n + D_1(n) =: E(n) + D_1(n).$$

Thus

$$\tilde{F}_1(x) = S_r(x)E(x) + S_r(x)D_1(x).$$

Since  $S_r E \in \Omega_\infty$  has no constant term, we deduce that

$$\psi([\tilde{F}_1]) = \psi([S_r D_1]) = S_r(1; \widehat{B}_{1+\diamond,1}).$$

The latter equation is derived as was (26), except that we regard the constant terms of the product of the polynomial  $S_r$  and the Laurent series  $D_1$ . From (25) we finally obtain

$$\log \mathcal{F}_{r,1} = \frac{1}{2} \log \mathcal{A}_r - \log \mathcal{A}_{r+1} + S_r(1; \widehat{B}_{1+\diamond,1} - \log \mathcal{A}_\diamond).$$

Now, we shall evaluate the expression above. For  $r = 0$  we get

$$\log \mathcal{F}_{0,1} = \frac{1}{12} + \frac{1}{2} \log \mathcal{A}_0 - 2 \log \mathcal{A}_1,$$

since

$$S_0(1; \widehat{B}_{1+\diamond,1} - \log \mathcal{A}_\diamond) = \widehat{B}_{2,1} - \log \mathcal{A}_1 = \frac{1}{12} - \log \mathcal{A}_1.$$

For now, let  $r > 0$ . We may represent  $\log \mathcal{F}_{r,1}$  in terms of  $\log \mathcal{A}_j$  as follows:

$$\log \mathcal{F}_{r,1} = \alpha_{r,0} + \sum_{j=1}^{r+1} \alpha_{r,j} \log \mathcal{A}_j.$$

The term  $\alpha_{r,0}$  is given by

$$\alpha_{r,0} = S_r(1; \widehat{B}_{1+\diamond,1}) = \sum_{j=0}^r \binom{r}{j} (-1)^{r-j} B_{r-j} \frac{\widehat{B}_{j+2,1}}{j+1}$$

where the sum runs over even  $j$ , since  $\widehat{B}_{j+2,1} = 0$  for odd  $j$ . If  $r$  is odd, then the sum simplifies to the term  $B_{r+1}/2r(r+1)$ . Otherwise we derive for even  $r$  that

$$\alpha_{r,0} = \sum_{j=0}^r \binom{r}{j} \frac{B_{r-j} B_{j+2}}{(j+1)^2(j+2)}.$$

It remains to determine the coefficients  $\alpha_{r,j}$  for  $r+1 \geq j \geq 1$ . Since  $\frac{1}{2}x^r - x^{r+1} - S_r(x)$  is an odd, resp. even, polynomial for even, resp. odd,  $r > 0$ , this property

transfers in a similar way to  $\frac{1}{2} \log \mathcal{A}_r - \log \mathcal{A}_{r+1} - S_r(1; \log \mathcal{A}_\circ)$ , such that  $\alpha_{r,j} = 0$  when  $2 \mid r - j$ . Otherwise we get

$$\alpha_{r,j} = -\binom{r}{j-1} \frac{B_{r-(j-1)}}{j} - \delta_{r+1,j} = -\binom{r+1}{j} \frac{B_{r+1-j}}{r+1} - \delta_{r+1,j} \quad (27)$$

for  $2 \nmid r - j$ , where the term  $-\log \mathcal{A}_{r+1}$  is represented by  $-\delta_{r+1,j}$ . □

**Corollary 31.** *Let  $r$  be an odd positive integer, then*

$$\begin{aligned} \log \mathcal{F}_{r,1} &= -\frac{r!}{(2\pi i)^{r+1}} \left( \frac{\zeta(r+1)}{r} + \sum_{j=1}^{\frac{r-1}{2}} \zeta(r+1-2j)\zeta(2j+1) - \frac{(r+2)\zeta(r+2)}{2} \right) \\ &= (-1)^{\frac{r-1}{2}} \frac{r!}{2} \left( \frac{|B_{r+1}|}{r(r+1)!} + \sum_{j=1}^{\frac{r-1}{2}} \frac{|B_{r+1-2j}| \zeta(2j+1)}{(r+1-2j)! (2\pi)^{2j}} - \frac{(r+2)\zeta(r+2)}{(2\pi)^{r+1}} \right). \end{aligned}$$

*Proof.* As a consequence of the functional equation of  $\zeta(s)$  and its derivative, we have for even positive integers  $n$  that

$$\log \mathcal{A}_n = -\zeta'(-n) = -\frac{1}{2} \frac{n!}{(2\pi i)^n} \zeta(n+1) \quad (28)$$

where the left hand side of (28) follows by definition [5, p. 276]. Theorem 29 provides

$$\log \mathcal{F}_{r,1} = \frac{B_{r+1}}{2r(r+1)} + \sum_{j=1}^{\frac{r+1}{2}} \alpha_{r,2j} \log \mathcal{A}_{2j}.$$

Combining (27) and (28) gives the second equation above. By Euler’s formula (2) we finally derive the first equation. □

**Remark 32.** For the sake of completeness, we give an analogue of (28) for odd integers. From the logarithmic derivatives of  $\Gamma(s)$  and the functional equation of  $\zeta(s)$ , see [5, pp. 183, 276], it follows for even positive integers  $n$ , that

$$\log \mathcal{A}_{n-1} = \frac{B_n}{n} H_{n-1} - \zeta'(1-n) = \frac{B_n}{n} (\gamma + \log(2\pi)) + 2 \frac{(n-1)!}{(2\pi i)^n} \zeta'(n)$$

where

$$\zeta'(n) = -\sum_{\nu=2}^{\infty} \log(\nu) \nu^{-n}.$$

However, MATHEMATICA is able to compute values of  $\zeta'$  for positive and negative argument values to any given precision.

**Result 33.** Exact expressions for  $\mathcal{F}_{r,1}$  in terms of  $\mathcal{A}_j$ :

Constant	Expression	Value
$\mathcal{F}_{0,1}$	$e^{\frac{1}{12}} \mathcal{A}_0^{\frac{1}{2}} \mathcal{A}_1^{-2}$	1.04633506677050318098...
$\mathcal{F}_{1,1}$	$e^{\frac{1}{24}} \mathcal{A}_2^{-\frac{3}{2}}$	0.99600199446870605433...
$\mathcal{F}_{2,1}$	$e^{\frac{7}{540}} \mathcal{A}_1^{-\frac{1}{6}} \mathcal{A}_3^{-\frac{4}{3}}$	0.99904614418135586848...
$\mathcal{F}_{3,1}$	$e^{-\frac{1}{720}} \mathcal{A}_2^{-\frac{1}{4}} \mathcal{A}_4^{-\frac{5}{4}}$	1.00097924030236153773...
$\mathcal{F}_{4,1}$	$e^{-\frac{67}{18900}} \mathcal{A}_1^{\frac{1}{30}} \mathcal{A}_3^{-\frac{1}{3}} \mathcal{A}_5^{-\frac{6}{5}}$	1.00007169725554110099...
$\mathcal{F}_{5,1}$	$e^{\frac{1}{2520}} \mathcal{A}_2^{\frac{1}{12}} \mathcal{A}_4^{-\frac{5}{12}} \mathcal{A}_6^{-\frac{7}{6}}$	0.99937792615674804266...

Exact expressions for  $\mathcal{F}_{r,1}$  in terms of  $\zeta(2j + 1)$ :

$$\mathcal{F}_{1,1} = \exp\left(\frac{1}{24} - \frac{3\zeta(3)}{8\pi^2}\right),$$

$$\mathcal{F}_{3,1} = \exp\left(-\frac{1}{720} - \frac{\zeta(3)}{16\pi^2} + \frac{15\zeta(5)}{16\pi^4}\right),$$

$$\mathcal{F}_{5,1} = \exp\left(\frac{1}{2520} + \frac{\zeta(3)}{48\pi^2} + \frac{5\zeta(5)}{16\pi^4} - \frac{105\zeta(7)}{16\pi^6}\right).$$

For the first 15 constants  $\mathcal{F}_{r,1}$  ( $r = 0, \dots, 14$ ) we find that

$$\max_{0 \leq r \leq 14} |\mathcal{F}_{r,1} - 1| < 0.05,$$

but, e.g.,  $\mathcal{F}_{19,1} \approx 371.61$  and  $\mathcal{F}_{20,1} \approx 1.16 \cdot 10^{-7}$ .

**Acknowledgements.** The author would like to thank Steven Finch for informing us about the formula of Milnor–Husemoller and the problem of finding suitable constants; also for giving several references. The author is also grateful to the referee for several remarks and suggestions.

**References**

[1] V. S. Adamchik, *Polygamma functions of negative order*, J. Comput. Appl. Math. **100** (1998), 191–199.

[2] V. S. Adamchik, *On the Barnes function*, Proc. 2001 Int. Symp. Symbolic and Algebraic Computation, Academic Press, 2001, 15–20.

[3] W. Alexeiewsky, *Ueber eine Classe von Functionen, die der Gammafunction analog sind*, Sitz.ber. Sächs. Akad. Wiss. Leipzig Math.-Nat.wiss. Kl. **46** (1894), 268–275.

[4] L. Bendersky, *Sur la fonction gamma généralisée*, Acta Math. **61** (1933), 263–322.

- [5] B. C. Berndt, *Ramanujan's Notebooks, Part I*, Springer-Verlag, 1985.
- [6] B. K. Choudhury, *The Riemann zeta-function and its derivatives*, Proc. Royal Soc. London A **450** (1995), 477–499.
- [7] N. G. de Bruijn, *Asymptotic Methods in Analysis*, third edition 1970. Reprinted by Dover, 1981.
- [8] S. R. Finch, *Abelian group enumeration constants, Mathematical Constants*, Cambridge Univ. Press, 2003, 273-276.
- [9] J. W. L. Glaisher, *On the product  $1^1 2^2 3^3 \cdots n^n$* , Messenger of Math. **7** (1878), 43-47.
- [10] R. L. Graham, D. E. Knuth, O. Patashnik, *Concrete Mathematics*, Addison-Wesley, Reading, MA, USA, 1994.
- [11] H. Kinkelin, *Ueber eine mit der Gammafunction verwandte Transcendente und deren Anwendung auf die Integralrechnung*, J. Reine Angew. Math. **57** (1860), 122-158.
- [12] M. Koecher, A. Krieg, *Elliptische Funktionen und Modulformen*, Springer-Verlag, 1998.
- [13] R. A. MacLeod, *Fractional part sums and divisor functions*, J. Number Theory **14** (1982), 185–227.
- [14] J. Milnor, D. Husemoller, *Symmetric bilinear forms*, Springer-Verlag, 1973.
- [15] F. W. Schäfke, A. Finsterer, *On Lindelöf's error bound for Stirling's series*, J. Reine Angew. Math. **404** (1990), 135–139.
- [16] N. J. A. Sloane, *Online Encyclopedia of Integer Sequences (OEIS)*, electronically published at: <http://www.research.att.com/~njas/sequences>.
- [17] Wolfram Research Inc., MATHEMATICA, Wolfram Research Inc., Champaign, IL.