

Knuth relations for the hyperoctahedral groups

Thomas Pietraho

Received: 27 March 2008 / Accepted: 22 July 2008 / Published online: 22 August 2008
© Springer Science+Business Media, LLC 2008

Abstract C. Bonnafé, M. Geck, L. Iancu, and T. Lam have conjectured a description of Kazhdan-Lusztig cells in unequal parameter Hecke algebras of type B which is based on domino tableaux of arbitrary rank. In the integer case, this generalizes the work of D. Garfinkle. We adapt her methods and construct a family of operators which generate the equivalence classes on pairs of arbitrary rank domino tableaux described in the above conjecture.

Keywords Unequal parameter Iwahori-Hecke algebra · Domino tableaux · Robinson-Schensted algorithm

1 Introduction

In [6], D. Garfinkle classified the primitive spectrum of the universal enveloping algebra for a complex semisimple Lie algebra in types B and C . By using annihilators of highest weight modules, this problem is reduced to studying equivalence classes in the corresponding Weyl group W_n . The existence of a Robinson-Schensted bijection between elements of W_n and same shape pairs of standard domino tableaux with n dominos [4] turns this into an essentially combinatorial problem. In fact, Garfinkle's classification shows that two elements in W_n are equivalent iff their left domino tableaux are related by moving through a set of open cycles, a certain combinatorial operation. The key step of this classification was achieved by studying the action of the wall-crossing operators arising from the general τ -invariant, as defined in [21], which were shown to be generators for both equivalences.

When interpreted in terms of cells of the equal parameter Hecke algebra of type B , the above work takes on a new meaning.

T. Pietraho (✉)
Bowdoin College, Brunswick, ME 04011, USA
e-mail: tpietrah@bowdoin.edu

In the 1980s, G. Lusztig extended the cell theory to Hecke algebras with unequal parameters; see [12] for an exposition. For Weyl groups of type B , cell theory then depends on an additional parameter s , which reduces to the equal parameter case when $s = 1$; Garfinkle's work classified cells in exactly this latter setting. As observed in [20], Garfinkle's bijection also admits an extension

$$G_r : W_n \rightarrow SDT_r(n) \times SDT_r(n)$$

to a bijection between W_n and same shape pairs of standard domino tableaux of rank r . It is reasonable to hope that the above two parameters can be linked and that a similar classification of cells is possible in this more general case. In fact:

Conjecture ([3]) *When s is a positive integer, two elements of W_n lie in the same left cell if and only if their left domino tableaux of rank $s - 1$ are related by moving through a set of open cycles.*

This is of course true when $s = 1$ and has been verified in the asymptotic case $s \geq n$, when the bijection G_{s-1} degenerates to the generalized Robinson-Schensted correspondence of [18], see [2]. Among finite Coxeter groups, cells in unequal parameter Hecke algebras have been classified in the dihedral groups and type F_4 by Lusztig [12] and Geck [7]. Only the problem of their classification in type B remains.

We will say that two elements of the hyperoctahedral group W_n are in the same irreducible combinatorial left cell of rank r if they share the same left domino tableau under the Robinson-Schensted map G_r , and in the same reducible combinatorial left cell of rank r if their rank r left domino tableaux are related by moving through an open cycle. The previous conjecture can be restated as:

Conjecture ([3]) *When s is a positive integer, left cells in W_n coincide with reducible combinatorial left cells of rank $s - 1$.*

Inspired by Garfinkle's approach in the equal parameter case, the main goal of this paper is to construct a set of generators for the reducible combinatorial left cells in arbitrary rank which draws on the notion of the generalized τ -invariant used in the equal parameter case. Such a set Λ^{r+1} is constructed in Section 3. The main theorem then can be stated as:

Theorem 3.9 *The family of operators Λ^{r+1} generates the reducible combinatorial left cells of rank r . More precisely, given w and $v \in W_n$ whose rank r left domino tableaux differ only by moving through a set of non-core open cycles, there is a sequence of operators in Λ^{r+1} sending w to v .*

This result falls into a family of similar theorems on generating sets for equivalence classes of standard tableaux, Garfinkle's work notwithstanding. In type A , a family of operators generating a similar equivalence for the symmetric group on standard Young tableaux appears in [11] and is known as the set of Knuth relations. Domino tableaux whose shapes are also partitions of nilpotent orbits in types B , C , or D correspond to the so-called orbital varieties. Their classification has been carried

out by W. M. McGovern in [13] by relying on a similar set of generators found in [9], see also [14]. The work of C. Bonnafé and L. Iancu in the asymptotic parameter case relies on finding a generating set for cells defined in terms of standard bitableaux. Finally, very recently M. Taskin has independently found another set of generators in the arbitrary rank case, see [19].

Combinatorial cells in the unequal parameter Hecke algebras in type B have also appeared in the work of I. Gordon and M. Martino, where it is shown that nilpotent points of the Calogero-Moser space correspond to the partitions arising from arbitrary rank domino tableaux [8].

This paper is organized as follows. In Section 2, we define the necessary objects and catalogue basic results. Section 3 defines the family of operators Λ^{r+1} and Section 4, in addition to showing that reducible combinatorial cells are stable under their action, describes this action on pairs of domino tableaux. In Section 5, we verify the main result, and leave the proof of a few crucial lemmas to Sections 6 and 7.

2 Definitions and preliminaries

2.1 Robinson-Schensted algorithms

The hyperoctahedral group W_n of rank n is the group of permutations of the set $\{\pm 1, \pm 2, \dots, \pm n\}$ which commute with the involution $i \mapsto -i$. It is the Weyl group of type B_n . We will write $w \in W_n$ in one-line notation as

$$w = (w(1) w(2) \dots w(n)).$$

A Young diagram is a finite left-justified array of squares arranged with non-increasing row lengths. We will denote the square in row i and column j of the diagram by $S_{i,j}$ so that $S_{1,1}$ is the uppermost left square in the Young diagram below:



Definition 2.1 Let $r \in \mathbb{N}$ and λ be a partition of a positive integer m . A domino tableau of rank r and shape λ is a Young diagram of shape λ whose squares are labeled by integers in such a way that 0 labels $S_{i,j}$ iff $i + j < r + 2$, each element of some set M labels exactly two adjacent squares, and all labels increase weakly along both rows and columns. A domino tableau is standard iff $M = \{1, \dots, n\}$ for some n . We will write $SDT_r(n)$ for the set of standard domino tableaux with n dominos. The set of squares labeled by 0 will be called the core of T .

Following [4] and [20], we describe the Robinson-Schensted bijections

$$G_r : W_n \rightarrow SDT_r(n) \times SDT_r(n)$$

between elements of W_n and same-shape pairs of rank r standard domino tableaux. The algorithm is based on an insertion map α which, given an entry $w(j)$ in the one-line notation for $w \in W_n$, inserts a domino with label $w(j)$ into a domino tableau.

This insertion map is similar to the usual Robinson-Schensted insertion map and is precisely defined in [4](1.2.5). To construct the left tableau, start with $T_1(0)$, the only tableau in $SDT_r(0)$. Define $T_1(1) = \alpha(w(1), T_1(0))$ and continue inductively by letting

$$T_1(k + 1) = \alpha(w(k + 1), T_1(k)).$$

The left domino tableau $T_1(n)$ will be standard and of rank r . The right tableau keeps track of the sequence of shapes of the left tableaux; we define $T_2(n)$ to be the unique tableau so that $T_2(k) \in SDT_r(k)$ has the same shape as $T_1(k)$ for all $k \leq n$. The Robinson-Schensted map is then defined by $G_r(w) = (T_1(n), T_2(n))$. We will also often use the notation $(T_1(n - 1), T_2(n - 1)) = (T_1(n), T_2(n))'$.

2.2 Cycles and extended cycles

We briefly recall the definition of a cycle in a domino tableau as well as a number of related notions which we will use later.

For a standard domino tableau T of arbitrary rank r , we will say the square S_{ij} is fixed when $i + j$ has the opposite parity as r , otherwise, we will call it variable. If S_{ij} is variable and i is odd, we will say S_{ij} is of type X ; if i is even, we will say S_{ij} is of type W . We will write $D(k, T)$ for the domino labeled by the positive integer k in T and $supp D(k, T)$ will denote its underlying squares. Write $label S_{ij}$ for the label of the square S_{ij} in T . We extend this notion slightly by letting $label S_{ij} = 0$ if either i or j is less than or equal to zero, and $label S_{ij} = \infty$ if i and j are positive but S_{ij} is not a square in T .

Definition 2.2 Suppose that $supp D(k, T) = \{S_{ij}, S_{i+1,j}\}$ or $\{S_{i,j-1}, S_{ij}\}$ and the square S_{ij} is fixed. Define $D'(k)$ to be a domino labeled by the integer k with $supp D'(k, T)$ equal to

1. $\{S_{ij}, S_{i-1,j}\}$ if $k < label S_{i-1,j+1}$
2. $\{S_{ij}, S_{i,j+1}\}$ if $k > label S_{i-1,j+1}$

Alternately, suppose that $supp D(k, T) = \{S_{ij}, S_{i-1,j}\}$ or $\{S_{i,j+1}, S_{ij}\}$ and the square S_{ij} is fixed. Define $supp D'(k, T)$ to be

1. $\{S_{ij}, S_{i,j-1}\}$ if $k < label S_{i+1,j-1}$
2. $\{S_{ij}, S_{i+1,j}\}$ if $k > label S_{i+1,j-1}$

Definition 2.3 The cycle $c = c(k, T)$ through k in a standard domino tableau T is a union of labels of T defined by the condition that $l \in c$ if either

1. $l = k$,
2. $supp D(l, T) \cap supp D'(m, T) \neq \emptyset$ for some $m \in c$, or
3. $supp D'(l, T) \cap supp D(m, T) \neq \emptyset$ for some $m \in c$.

We will identify the labels contained in a cycle with their underlying dominos.

Starting with a standard domino tableau T of rank r containing a cycle c , it is possible to define a new domino tableau $MT(T, c)$ by replacing every domino

$D(l, T) \in c$ by the shifted domino $D'(l, T)$ defined above. This operation changes the labels of the variable squares in c while preserving the labels of all of the fixed squares of T . In fact, if we pick a label l of a square in T , the definition of a cycle together with [4](1.5.27) imply that moving through $c(l, T)$ is in some sense the minimal transformation of T which changes the label of the variable square of $D(l, T)$, maintains the labels of all of the fixed squares of T , and results in a standard domino tableau.

The shape of $MT(T, c)$ either equals the shape of T , or one square will be removed (or added to its core) and one will be added. In the former case, the cycle c is called closed; otherwise, it is called open. If moving through c adds a square to the core, we will call c a core open cycle; the other open cycles will be called non-core. For an open cycle c of a tableau T , we will write $S_b(c)$ for the square that has been removed or added to the core by moving through c . Similarly, we will write $S_f(c)$ for the square that is added to the shape of T . Note that $S_b(c)$ and $S_f(c)$ are always variable squares.

Definition 2.4 A variable square S_{ij} satisfying the conditions that

1. neither $S_{i,j+1}$ nor $S_{i+1,j}$ lie in T , and
2. either
 - a) both $S_{i-1,j}$ and $S_{i,j-1}$ lie in T , or
 - b) either $S_{i-1,j}$ lies in T and $j = 1$ or $S_{i,j-1}$ lies in T and $i = 1$,

will be called a hole if it is of type W and a corner if it is of type X . It will be called full if $S_{ij} \in T$ and empty otherwise.

Let U be a set of cycles in T . Because the order in which one moves through a set of cycles is immaterial by [4](1.5.29), we can unambiguously write $MT(T, U)$ for the tableau obtained by moving through all of the cycles in the set U .

Moving through a cycle in a pair of same-shape tableaux is somewhat problematic, as it may result in a pair of tableaux which is not same-shape. We require the following definition.

Definition 2.5 Consider (T_1, T_2) a pair of same-shape domino tableaux, k a label in T_1 , and c the cycle in T_1 through k . The extended cycle \tilde{c} of k in T_1 relative to T_2 is a union of cycles in T_1 which contains c . Further, the union of two cycles $c_1 \cup c_2$ lies in \tilde{c} if either one is contained in \tilde{c} and, for some cycle d in T_2 , $S_b(d)$ coincides with a square of c_1 and $S_f(d)$ coincides with a square of $MT(T_1, c_2)$. The symmetric notion of an extended cycle in T_2 relative to T_1 is defined in the natural way.

For an extended cycle \tilde{c} in T_2 relative to T_1 , write $\tilde{c} = c_1 \cup \dots \cup c_m$ and let d_1, \dots, d_m be cycles in T_1 such that $S_b(c_i) = S_b(d_i)$ for all i , $S_f(d_m) = S_f(c_1)$, and $S_f(d_i) = S_f(c_{i+1})$ for $1 \leq i < m$. The union $\tilde{d} = d_1 \cup \dots \cup d_m$ is an extended cycle in T_1 relative to T_2 called the extended cycle corresponding to \tilde{c} . Symmetrically, \tilde{c} is the extended cycle corresponding to \tilde{d} .

We can now define a moving through operation for a pair of same-shape domino tableaux. Write b for an ordered pair (\tilde{c}, \tilde{d}) of extended cycles in (T_1, T_2) that corre-

spend to each other. Define $MT((T_1, T_2), b)$ to equal

$$(MT(T_1, \tilde{c}), MT(T_2, \tilde{d})).$$

It is clear that this operation produces another pair of same-shape domino tableaux. Speaking loosely, we will often refer to this operation as moving through either of the extended cycles \tilde{c} or \tilde{d} . Note that if the cycle c is closed, then $\tilde{c} = c$ and moving through a pair of tableaux boils down to the operation $(MT(T_1, c), T_2)$.

Definition 2.6 We will say that a set of squares in a domino tableau is *boxed* iff it is entirely contained in a set of squares of the form

$$\{S_{ij}, S_{i+1,j}, S_{i,j+1}, S_{i+1,j+1}\}$$

where S_{ij} must be of type X . A set will be called *unboxed* if it is not boxed.

Boxing is well-behaved with respect to cycles. If c is a cycle in T , then all of its underlying dominos are either boxed or unboxed. Furthermore, moving through a domino changes its boxing, and consequently, the boxing of all the dominos in the cycle containing it. The same holds for all of the dominos in an extended cycle of a same-shape tableau pair.

We will say that two sets of squares are adjacent in a tableau T if there are two squares, one in each set, which share a common side. One set of squares in T will be said to be below another if all of its squares lie in rows strictly below the rows of the squares of the other set. The notion of above is defined similarly. We will often call the squares underlying a domino its position, and say that a position is extremal if its removal from a Young diagram results in another Young diagram corresponding to a partition of the same rank.

2.3 Combinatorial cells

Definition 2.7 Consider $w, v \in W_n$ of type B_n and fix a non-negative integer r letting $G_r(w) = (T_1, T_2)$ and $G_r(v) = (\tilde{T}_1, \tilde{T}_2)$. We will say that w and v are

1. in the same irreducible combinatorial left cell of rank r if $T_2 = \tilde{T}_2$, and
2. in the same reducible combinatorial left cell of rank r if there is a set U of non-core open cycles in T_2 such that $\tilde{T}_2 = MT(T_2, U)$.

We will say that w and v are in the same irreducible and reducible combinatorial right cells iff their inverses lie in the same irreducible and reducible combinatorial left cells, respectively.

When $r \geq n - 1$, the situation is somewhat simpler; there are no non-core open cycles implying that combinatorial left cells are determined simply by right tableaux. Furthermore, by the main result of [15], for these values of r all combinatorial cells are actually independent of r .

Reducible combinatorial left cells of rank r can also be described in terms of right tableaux of ranks r and $r + 1$. A similar characterization holds for combinatorial right cells.

Theorem 2.8 ([16]) *Reducible combinatorial left cells of rank r in the Weyl group of type B_n are generated by the equivalence relations of having the same right tableau in either rank r or rank $r + 1$.*

In what follows we will focus on reducible combinatorial right cells. Since our generalizations of the Robinson-Schensted algorithm G_r behave well with respect to inverses by merely changing the order of the two tableaux, all the statements can be easily modified to apply to reducible combinatorial left cells as well.

3 Knuth relations

The purpose of this section is to define the operators on the hyperoctahedral group W_n which generate the reducible combinatorial right cells of rank r . We first recount the situation in type A for the symmetric group S_n .

3.1 Type A

Writing the elements of S_n in one-line notation, the Knuth relations on S_n are the transformations

$$(w(1) w(2) \dots w(j - 1) w(j + 1) w(j) \dots w(n))$$

which transpose the j th and $(j + 1)$ st entries of $w \in S_n$ whenever

- (i) $j \geq 2$ and $w(j - 1)$ lies between $w(j)$ and $w(j + 1)$, or
- (ii) $j < n - 1$ and $w(j + 2)$ lies between $w(j)$ and $w(j + 1)$.

Every Knuth relation preserves the Robinson-Schensted left tableau $T_1(w)$ of w . However, even more can be said:

Theorem 3.1 ([11]) *Knuth relations generate the combinatorial left cells in S_n . More precisely, given w and v with $T_1(w) = T_1(v)$, there is a sequence of Knuth relations sending w to v .*

Aiming to adapt this theorem to the hyperoctahedral groups, we begin by rephrasing the Knuth relations first in terms of the length function on S_n , and then again in terms of the τ -invariant.

The group S_n is a Weyl group of a complex semisimple Lie algebra \mathfrak{g} of type A_{n-1} with Cartan subalgebra \mathfrak{h} . Let $\Pi_n = \{\alpha_1, \alpha_2, \dots, \alpha_{n-1}\}$ be a set of simple roots for a choice of positive roots in the root system $\Delta(\mathfrak{g}, \mathfrak{h})$ and write s_α for the simple reflection corresponding to $\alpha \in \Pi_n$. We view S_n as the group generated by the above simple reflections and let $\ell : S_n \rightarrow \mathbb{Z}$ be the corresponding length function on S_n . If we identify the reflection s_{α_j} with the transposition interchanging the j th and $(j + 1)$ st entries of the permutation corresponding to $w \in S_n$, then the Knuth relations on S_n are the transformations taking w to ws_{α_j} . Noting that $\ell(ws_{\alpha_j}) < \ell(w)$ exactly when $w(j) > w(j + 1)$, the domain for the Knuth relations is the set of $w \in S_n$ satisfying

- (i) $\ell(ws_{\alpha_i}) < \ell(w) < \ell(ws_{\alpha_j}) < \ell(ws_{\alpha_j}s_{\alpha_i})$, or
- (ii) $\ell(ws_{\alpha_i}) > \ell(w) > \ell(ws_{\alpha_j}) > \ell(ws_{\alpha_j}s_{\alpha_i})$

for some $\alpha_i \in \Pi_n$. This condition is satisfied only when α_i and α_j are adjacent simple roots, that is, when $i = j \pm 1$.

Armed with this restatement of the Knuth conditions, following [21] we define the τ -invariant for S_n and a related family of operators.

Definition 3.2 Write $W = S_n$. For $w \in W$, let $\tau(w) = \{\alpha \in \Pi_n \mid \ell(ws_{\alpha}) < \ell(w)\}$. Given simple roots α and β in Π_n , let

$$D_{\alpha\beta}(W) = \{w \in W \mid \alpha \notin \tau(w) \text{ and } \beta \in \tau(w)\}.$$

When α and β are adjacent roots, define $T_{\alpha\beta} : D_{\alpha\beta}(W) \rightarrow D_{\beta\alpha}(W)$ by

$$T_{\alpha\beta}(w) = \{ws_{\alpha}, ws_{\beta}\} \cap D_{\beta\alpha}(W).$$

When defined, the operators $T_{\alpha\beta}$ are single-valued and preserve the Robinson-Schensted left tableau $T_1(w)$ of w . These so-called wall-crossing operators also appear as the “star operator” in [10]. The following is a direct consequence of the result on Knuth relations:

Corollary 3.3 *The family of operators $T_{\alpha\beta} : D_{\alpha\beta}(W) \rightarrow D_{\beta\alpha}(W)$ generates the combinatorial left cells in S_n . More precisely, given w and v with $T_1(w) = T_1(v)$, there is a sequence of $T_{\alpha\beta}$ operators sending w to v .*

That combinatorial left cells for S_n coincide with Kazhdan-Lusztig left cells is shown in [10] using A. Joseph and D. Vogan’s results on the primitive spectrum of semisimple Lie algebras. A direct proof of this fact using entirely combinatorial methods has subsequently appeared, see [1].

3.2 Type B

In order to define similar relations for the hyperoctahedral groups, we mimic the final construction in type A. The group W_n is a Weyl group of a complex semisimple Lie algebra \mathfrak{g} of type B_n with Cartan subalgebra \mathfrak{h} . Let $\{\epsilon_1, \dots, \epsilon_n\}$ be a basis for \mathfrak{h}^* such that if we define $\alpha'_1 = \epsilon_1$ and $\alpha_i = \epsilon_i - \epsilon_{i-1}$, then

$$\Pi_n = \{\alpha'_1, \alpha_2, \dots, \alpha_n\}$$

is the set of simple roots for some choice of positive roots $\Delta^+(\mathfrak{g}, \mathfrak{h})$. While this choice of simple roots is not standard, we adopt it to obtain somewhat cleaner statements and reconcile our work with [6].

We modify Π_n slightly to include certain non-simple roots. For $i \leq n$, let $\alpha'_i = \alpha'_1 + \alpha_2 + \dots + \alpha_i$ and when k is a non-negative integer write

$$\Pi_n^k = \{\alpha'_1, \alpha'_2, \dots, \alpha'_{\min(n,k)}, \alpha_2, \dots, \alpha_n\}.$$

Further, write s_i for the simple reflection $s_{\alpha_{i+1}}$ and t_i for the reflection $s_{\alpha'_i}$. We realize W_n as a set of signed permutations on n letters by identifying s_i with the transposition $(1\ 2 \dots i + 1\ i \dots n)$ and t_i with the element $(1\ 2 \dots -i \dots n)$.

The generating set for reducible combinatorial right cells will be drawn out of the following three types of operators:

Definition 3.4 For $w \in W_n$, and a non-negative integer k , let

$$\tau^k(w) = \{\alpha \in \Pi_n^k \mid \ell(ws_\alpha) < \ell(w)\}.$$

Given roots α and β in Π_n^k define $D_{\alpha\beta}^k(W_n) = \{w \mid \alpha \notin \tau^k(w) \text{ and } \beta \in \tau^k(w)\}$. The operator $T_{\alpha\beta}^k : D_{\alpha\beta}^k(W_n) \rightarrow D_{\beta\alpha}^k(W_n)$ is defined by

$$T_{\alpha\beta}^k(w) = \{ws_\alpha, ws_\beta\} \cap D_{\beta\alpha}^k(W_n).$$

The above definitions add the parameter k to the standard notions of root system Π_n , τ -invariant $\tau(w)$, domains $D_{\alpha\beta}$, and the wall-crossing operators $T_{\alpha\beta}$ as defined explicitly, say in [5]. Our definitions coincide with these standard ones when the value of this parameter is $k = 1$.

When the set $\{\alpha, \beta\}$ contains roots of unequal length, the operator $T_{\alpha\beta}^k$ may be two-valued. The next definitions are an attempt to remedy this by splitting this operator into two new ones: T_{IN}^k and T_{SC}^k . Henceforth, we will reserve the notation $T_{\alpha\beta}$ to mean $T_{\alpha\beta}^0$ and $D_{\alpha\beta}$ to mean $D_{\alpha\beta}^0$, in which case the operators are single-valued.

Definition 3.5 We will say $w \in D_{IN}^k(W_n)$ if for some $\{\delta, \beta\} = \{\alpha_{k+1}, \alpha'_k\}$, we have

$$w \in D_{\delta\beta}^k(W_n) \text{ and } ws_k \in D_{\beta\delta}^k(W_n).$$

Define T_{IN}^k for $w \in D_{IN}^k(W_n)$ by $T_{IN}^k(w) = ws_k$.

Definition 3.6 We will say $w \in D_{SC}^k(W_n)$ if for some choice of $\{\delta_i, \beta_i\} = \{\alpha_{i+1}, \alpha'_i\}$ for every $i \leq k$, we have

$$w \in D_{\delta_i\beta_i}^k(W_n) \text{ and } wt_i \in D_{\beta_i\delta_i}^k(W_n) \quad \forall i \leq k.$$

Define T_{SC}^k for $w \in D_{SC}^k(W_n)$ by $T_{SC}^k(w) = wt_1$.

To phrase the above in more digestible terms, we note that in W_n the length function satisfies $\ell(ws_j) < \ell(w)$ iff $w(j) > w(j + 1)$, and $\ell(wt_j) < \ell(w)$ iff $w(j) < 0$. Armed with this characterization, the domains and actions for the three types of operators which we will be interested in can be described more succinctly.

$T_{\alpha\beta}$: w lies in $D_{\alpha\beta}(W)$ for $\{\alpha, \beta\} = \{\alpha_{j+1}, \alpha_{j+2}\}$ whenever $w(j + 1)$ is either greater than or smaller than both $w(j)$ and $w(j + 2)$; $T_{\alpha\beta}$ then interchanges the smallest and largest entries among $w(j), w(j + 1)$, and $w(j + 2)$.

T_{IN}^k : $w \in D_{IN}^k(W)$ iff $w(k)$ and $w(k + 1)$ are of opposite sign. The operator T_{IN}^k interchanges the k th and $k + 1$ st entries of w .

T_{SC}^k : $w \in D_{SC}^k(W)$ iff $|w(1)| > |w(2)| > \dots > |w(k + 1)|$. The operator T_{SC}^k changes the sign of $w(1)$.

Definition 3.7 For an integer k , we define a set of operators

$$\Lambda^k = \{T_{\alpha\beta} \mid \alpha, \beta \in \Pi_n^0\} \cup \{T_{IN}^i \mid i \leq k\} \cup \{T_{SC}^k\}.$$

This is the sought-after set of generators for reducible combinatorial right cells of arbitrary rank. The following are our two main results:

Theorem 3.8 *The operators in Λ^{r+1} preserve reducible combinatorial left cells of rank r . If $S \in \Lambda^{r+1}$, and $w \in W_n$ is in the domain of S , then the Robinson-Schensted left domino tableaux $T_1(w)$ and $T_1(Sw)$ of rank r differ only by moving through a set of non-core open cycles in $T_1(w)$.*

Theorem 3.9 *The family of operators Λ^{r+1} generates the reducible combinatorial left cells of rank r . More precisely, given w and $v \in W_n$ whose Robinson-Schensted left domino tableaux $T_1(w)$ and $T_1(v)$ of rank r differ only by moving through a set of non-core open cycles, there is a sequence of operators in Λ^{r+1} sending w to v .*

These results have previously been obtained in two special cases. In her work on the primitive spectrum of the universal enveloping algebras in types B and C , Devra Garfinkle constructed a generating set for the reducible combinatorial cells of rank zero [6]. Additionally, C. Bonnafé and L. Iancu obtained a generating set for combinatorial left cells in the so-called asymptotic case when $r \geq n - 1$. The generating set Λ^{r+1} proposed above for domino tableaux of arbitrary rank generalizes both of these results.

Theorem 3.10 (D. Garfinkle) *The reducible combinatorial left cells of rank zero are generated by the family of operators $T_{\alpha\beta}^1$ where α and β are adjacent simple roots in Π_n .*

At first glance, the set Λ^1 is not exactly Garfinkle’s generating set. The latter contains multi-valued operators $T_{\alpha'_1\alpha_2}^1$ and $T_{\alpha_2\alpha'_1}^1$. However, we note that the combined action and domains of $T_{\alpha'_1\alpha_2}^1$ and $T_{\alpha_2\alpha'_1}^1$ agree precisely with those of T_{SC}^1 and $T_{IN}^1 \in \Lambda^1$. The operators T_{SC}^1 and T_{IN}^1 merely split up Garfinkle’s original multi-valued operators.

Theorem 3.11 (C. Bonnafé and L. Iancu) *When the rank $r \geq n - 1$, the reducible and irreducible combinatorial left cells coincide. They are generated by the family of operators $\Lambda^{r+1} = \Lambda^n$.*

We note that in the asymptotic case, the operator T_{SC}^{r+1} is not defined, so that Λ^{r+1} consists entirely of the $T_{\alpha\beta}$ and T_{IN}^k operators, as exhibited in [2].

It was observed in [17] that combinatorial right cells are in general not well-behaved with increasing rank. This phenomenon is readily explained by examining the composition of Λ^{r+1} more closely. While increasing r expands the family of T_{IN} operators contained in Λ^{r+1} , it also diminishes the domain of T_{SC} . The complicated behavior of right cells as r increases is a manifestation of this interplay.

Recent work of M. Taskin finds a family of relations on W_n which generates the irreducible combinatorial left cells of rank r . Using the results of [16] which show that reducible combinatorial left cells are common refinements of irreducible ones, it is then possible to describe a set of generators of the reducible combinatorial left cells of rank r based on Taskin’s work. This set, however, does not coincide with Λ_{r+1} , but can be shown to be equivalent. It would be desirable to use the methods of the current paper to produce results on irreducible cells; however, it seems to the author that the moving-through operation which changes the shape of the underlying tableau is very intricately intertwined with the method of proof and the necessary unentanglement does not appear to be an easy task.

4 Stability under Λ^{r+1}

The goal of this section is to prove Theorem 3.8, showing that reducible right cells in rank r are invariant under the operators in Λ^{r+1} . We will rely on a decomposition of W_n , which we describe presently. Let W_m be the subgroup of W_n generated by the reflections $\{s_1, \dots, s_{m-1}, t_1\}$ and define X_m^n to be the set of $x \in W_n$ which satisfy

$$0 < x(1) < x(2) < \dots < x(m).$$

Then X_m^n is a cross-section of W_n/W_m and we can write every $w \in W_n$ as a product $w = xw'$ with $x \in X_m^n$ and $w' \in W_m$. The following appears in [3].

Proposition 4.1 *If $x \in X_m^n$ and $w', v' \in W_m$ are in the same irreducible combinatorial left cell of rank r , then xw' and $xv' \in W_n$ are in the same irreducible combinatorial left cell of rank r .*

Corollary 4.2 *If $x \in X_m^n$ and $w', v' \in W_m$ are in the same reducible combinatorial left cell of rank r , then xw' and $xv' \in W_n$ are in the same reducible combinatorial left cell of rank r .*

Proof The main result of [16] shows that reducible combinatorial left cells of rank r are the least common refinements of irreducible combinatorial left cells of ranks r and $r + 1$. Hence the corollary follows directly from the above proposition. □

4.1 The operator T_{SC}

We first examine the family of operators T_{SC}^k which, under appropriate circumstances, change the sign of the first entry of w .

Proposition 4.3 *The operator T_{SC}^{r+1} preserves the irreducible and, consequently, the reducible combinatorial right cells of rank r in W_{r+2} .*


Proof Consider $w \in D_{SC}^{r+1}(W_{r+2})$ so that

$$|w(1)| > |w(2)| > \dots |w(r + 2)|.$$


Write $v = w^{-1}$ and $u = (T_{SC}^{r+1}(w))^{-1}$. We will show that their right tableaux $T_2(v)$ and $T_2(u)$ agree. Note that the condition on w implies $|v(1)| > |v(2)| > \dots |v(r + 2)|$ and the fact that T_{SC}^{r+1} only changes the sign of $w(1)$ forces

1. $v(i) = u(i)$ for $i \leq r + 1$, and
2. $u(r + 2) = -v(r + 2) \in \{\pm 1\}$.

If we write the positive entries in v as $a_1 > \dots > a_p$ and the negative ones as $-b_1 < \dots < -b_q$, then the left tableau of v before the insertion of $v(r + 2)$ must have the form:



This is also the left tableau of u before the insertion of $u(r + 2)$ and we point out that the corresponding right tableaux for both u and v are the same. If $a_1 > b_1$, then the insertion of $v(r + 2)$ and $u(r + 2)$ into the above tableau yields either:




In either case, the same domino is added to their shared right tableau and $T_2(v) = T_2(u)$. The proof is similar when $a_1 < b_1$. □

Corollary 4.4 *When $n \geq r + 2$, the operator T_{SC}^{r+1} preserves the irreducible and, consequently, the reducible, combinatorial left cells of rank r in W_n .*

Proof For an element w of the set $D_{SC}^{r+1}(W_n)$, write $w = xw' \in X_n^{r+2}W_{r+2}$. Then $w' \in D_{SC}^{r+1}(W_{r+2})$ and by Proposition 4.3, $T_{SC}^{r+1}(w')$ and w' share the same irreducible combinatorial left cell of rank r . Since $T_{SC}^{r+1}(w) = xT_{SC}^{r+1}(w') \in X_n^{r+2}W_{r+2}$, Proposition 4.1 and Corollary 4.2 imply the same is true of $T_{SC}^{r+1}(w)$ and w . □

Our final aim is to describe the action of T_{SC}^{r+1} on the right tableau of w . This is more or less captured in the proof of Proposition 4.3, but we will attempt to be more precise. Consider $w \in D_{SC}^{r+1}(W_n)$, write T_2 for the right tableau $T_2(w)$ of rank r , and

define $T_2(r + 2)$ to be the subtableau of T_2 which contains only dominos whose labels are less than or equal to $r + 2$. Because $w \in D_{SC}^{r+1}(W_n)$, $T_2(r + 2)$ contains exactly the dominos adjacent to the core of T_2 as well as the unique domino of T_2 which shares its long edge with the long edge of one of the dominos adjacent to the core. Furthermore, $T_2(r + 2)$ must have the form



where $\alpha_1 = r + 2$, noting that p may equal zero. By the proof of Proposition 4.3, T_{SC}^{r+1} acts on $T_2(r + 2)$ by interchanging one of its possible two forms of the same shape with the other. It is easy to see that the remaining dominos of the tableaux T_2 and the right tableau $T_2(T_{SC}^{r+1}(w))$ are the same, since both keep track of the subsequent insertions into the left tableau $T_1(w)(r + 2) = T_1(T_{SC}^{k+1}(w))(r + 2)$.

Example 4.5 Consider $w = (4, -3, -2, 1)$ which lies in $D_{SC}^3(W_4)$. Then $T_{SC}^3(w) = (-4, -3, -2, 1)$ and

$$T_2(w) = \begin{array}{|c|c|c|} \hline & & 1 \\ \hline & & 4 \\ \hline & 3 & \\ \hline 2 & & \\ \hline \end{array} \quad \text{and} \quad T_2(T_{SC}^3(w)) = \begin{array}{|c|c|c|} \hline & & 3 \quad 4 \\ \hline & 2 & \\ \hline 1 & & \\ \hline \end{array}$$

On a final note, we observe that this description of the action of T_{SC}^{r+1} reproduces the action of a portion of the multi-valued $T_{\alpha'_1\alpha_2}$ operator detailed in [5](2.3.4) for the rank zero case.

4.2 The operators $T_{\alpha\beta}$

Given adjacent simple roots $\{\alpha, \beta\} = \{\alpha_{j+1}, \alpha_{j+2}\}$, an operator $T_{\alpha\beta}$ interchanges the smallest and largest entries among $w(j)$, $w(j + 1)$, and $w(j + 2)$. The following is a consequence of Proposition 3.10 in [3].

Proposition 4.6 *The operators $T_{\alpha\beta}$ preserve the irreducible and, consequently, the reducible combinatorial right cells of rank r in W_n .*

What is still required is a description of their action on right tableaux. We begin with an explicit description of their domains. Recall that $\alpha_{j+1} \in \tau(w)$ iff $\ell(ws_{\alpha_{j+1}}) < \ell(w)$ which occurs iff $w(j) > w(j + 1)$. This condition is easily read off from the right tableau $T_2(w)$. We will say that k lies below l in a domino tableau iff every row containing a square of the domino labeled k lies below every row of the domino

labeled l . Then $\alpha_{j+1} \in \tau(w)$ iff $j + 1$ lies below j in $T_2(w)$. Unraveling the definition of $D_{\alpha\beta}(W)$, we find that when $\{\alpha, \beta\} = \{\alpha_{j+1}, \alpha_{j+2}\}$, w lies in the domain of $T_{\alpha\beta}$ iff in the tableau $T_R(w)$ either:

1. $j + 1$ lies below j and $j + 2$ does not lie below $j + 1$, or
2. $j + 1$ does not lie below j and $j + 2$ lies below $j + 1$.

Two cases are necessary to describe the action of $T_{\alpha\beta}$.

Case 1. In the following, let $k = j + 1$ and $l = j + 2$. If either one of the following four configurations of dominos appears in $T_2(w)$:

$$\begin{aligned}
 F_1(j) &= \begin{array}{|c|c|} \hline j & k \\ \hline & l \\ \hline \end{array} & \text{and} & \tilde{F}_1(j) &= \begin{array}{|c|c|} \hline j & l \\ \hline k & \\ \hline \end{array} \\
 F_2(j) &= \begin{array}{|c|c|} \hline j & \\ \hline k & l \\ \hline \end{array} & \text{and} & \tilde{F}_2(j) &= \begin{array}{|c|c|} \hline j & k \\ \hline & l \\ \hline \end{array}
 \end{aligned}$$

then the action $T_{\alpha\beta}$ on the right tableau of w swaps $F_i(j)$ and $\tilde{F}_i(j)$ within $T_R(w)$.

Case 2. If none of the above four configurations appear in $T_2(w)$ and w lies in the domain $D_{\alpha\beta}(W_n)$, then $T_{\alpha\beta}$ acts by swapping either the dominos labeled j and $j + 1$ or $j + 1$ and $j + 2$.

The proof that this description of $T_{\alpha\beta}$ on domino tableau accurately depicts the action of $T_{\alpha\beta}$ defined on W_n is not difficult. It appears as [5](2.1.19) in rank zero, and follows almost identically for higher rank tableaux.

4.3 The operators T_{IN}^k

The operator T_{IN}^k interchanges the entries $w(k)$ and $w(k + 1)$ in w whenever they are of opposite signs. The following is again a consequence of Proposition 3.10 in [3].

Proposition 4.7 *When $k \leq r$, the operator T_{IN}^k preserves the irreducible and, consequently, the reducible combinatorial left cells of rank r in W_n .*

Slightly more is true:

Proposition 4.8 *The operator T_{IN}^{r+1} preserves the reducible combinatorial left cells of rank r in W_n .*

Proof By Proposition 4.7, the operator T_{IN}^{r+1} preserves the irreducible combinatorial left cells of rank $r + 1$ in W_n . Since reducible combinatorial left cells of rank r are the least common refinements of irreducible cells in ranks r and $r + 1$, the result follows. □

Next, we describe the action of T_{IN}^k on the left and right tableaux $T_1(w)$ and $T_2(w)$. Recall that $\alpha'_k \in \tau(w)$ iff $\ell(wt_k) < \ell(w)$ which occurs iff $w(k) < 0$. As long as $k \leq r + 1$, this occurs iff the domino with label k in $T_2(w)$ is vertical. Recalling that $\alpha_{k+1} \in \tau(w)$ iff $k + 1$ lies below k in $T_2(w)$, we can describe the domain of T_{IN}^k as follows.

Definition 4.9 We will call a tableau T of rank r *sparse* if there is a square of the form $S_{m,r+3-m}$ which is empty in T .

Case 1. If the tableau $T_2(w)(r + 2)$ is sparse, then $w \in S_{IN}^k$ for all $k \leq r + 1$ iff the dominos with labels k and $k + 1$ in $T_2(w)$ are of opposite orientations. In this setting, since T_{IN}^k swaps $w(k)$ and $w(k + 1)$ in w whenever they are of opposite sign, its action merely reverses the order in which the k th and $k + 1$ st dominos are inserted into the left tableau. Since $T_2(w)(r + 2)$ is sparse, these insertions do not interact with each other. Thus on the shape-tracking tableau $T_2(w)$, T_{IN}^k merely swaps the dominos with labels k and $k + 1$, while acting trivially on $T_1(w)$.

Case 2. If the tableau $T_2(w)(r + 2)$ is not sparse, then $k = r + 1$ and the insertions of $w(k)$ and $w(k + 1)$ into the left tableau interact with each other.

It is easy to see that there are four possible configurations of dominos with labels $k = r + 1$ and $l = r + 2$ within $T_2(w)$ when $w \in S_{IN}^{r+1}(W)$:

$$\begin{aligned}
 E_0(j) &= \begin{array}{|c|c|} \hline k & l \\ \hline \end{array} & \text{and} & \tilde{E}_0(j) &= \begin{array}{|c|c|} \hline k & l \\ \hline \end{array} \\
 E_1(j) &= \begin{array}{|c|} \hline k \\ \hline l \\ \hline \end{array} & \text{and} & \tilde{E}_1(j) &= \begin{array}{|c|} \hline k \\ \hline l \\ \hline \end{array}
 \end{aligned}$$

Proposition 4.10 Consider $w \in D_{IN}^{r+1}(W)$ and let $v = T_{IN}^{r+1}(w)$. If $G_r(w) = (T_1, T_2)$, then the tableau pair $G_r(v) = (\tilde{T}_1, \tilde{T}_2)$ admits the following description:

1. If $T_2(r + 2)$ is sparse, then $T_1 = \tilde{T}_1$ and \tilde{T}_2 is obtained by swapping the dominos with labels $r + 1$ and $r + 2$ in the tableau T_2 .
2. If $T_2(r + 2)$ is not sparse, then the description of the action depends on the exact configuration of the dominos with labels $r + 1$ and $r + 2$ in T_2 :
 - (a) If the configuration $E_0(r + 1)$ or $E_1(r + 1)$ appears in T_2 , write it as $E_i(r + 1)$ and let $\tilde{T}_2 = (T_2 \setminus E_i(r + 1)) \cup E_{1-i}(r + 1)$. Let c be the extended cycle through $r + 2$ in \tilde{T}_2 relative to T_1 . Then

$$(\tilde{T}_1, \tilde{T}_2) = MT((T_1, \tilde{T}_2), c).$$

- (b) If $\tilde{E}_0(r + 1)$ or $\tilde{E}_1(r + 1)$ appears in T_2 , let c be the extended cycle through $r + 2$ in T_2 relative to T_1 and define $(\tilde{T}_1, \tilde{T}_2) = MT((T_1, T_2), c)$. Note that \tilde{T}_2 must contain one of the configurations $E_0(k)$ or $E_1(k)$, which we label $E_i(r + 1)$. Then

$$(\tilde{T}_1, \tilde{T}_2) = (\tilde{T}_1, (\tilde{T}_2 \setminus E_i(r + 1)) \cup E_{1-i}(r + 1)).$$

Proof The first case has already been considered above. In the special situation when $n = r + 2$, the second case follows by inspection. We mimic the proof of [5](2.3.8) to verify the second case in general. So consider $w \in D_{IN}^{r+1}(W)$ assuming that $T_2(r + 2)$ is not sparse. By symmetry and the fact that T_{IN}^{r+1} is an involution, it is sufficient to consider only w for which also $|w(r + 1)| < |w(r + 2)|$, $w(r + 1) > 0$ and $w(r + 2) < 0$. For such a w , let $\bar{w} = vt_{r+1}$. Write $G_r(\bar{w}) = (\bar{T}_1, \bar{T}_2)$. Then

1. $\tilde{E}_1(r + 1) \subset T_2, E_0(r + 1) \subset \tilde{T}_2,$ and $E_1(r + 1) \subset \tilde{T}_2,$
2. $\tilde{T}_2 = (\tilde{T}_2 \setminus E_0(r + 1)) \cup E_1(r + 1)$ and $\tilde{T}_1 = \tilde{T}_1,$
3. $(\tilde{T}_1, \tilde{T}_2) = MT((T_1, T_2), c)$ where c is the extended cycle through $r + 2$ in T_2 relative to $T_1.$

Once these are verified, the proposition follows. The last two parts imply that $\tilde{T}_1 = \tilde{T}_1,$ and the latter equals $MT(T_1, d)$ where d is the extended cycle in T_1 corresponding to $c,$ as desired. Meanwhile, the second part then implies that \tilde{T}_2 will be as specified by the proposition.

Statements (1) and (2) follow easily from the definition of domino insertion, while the proof of (3) is identical to the rank zero case. That the extended cycle through $r + 2$ in T_2 consists only of non-core cycles follows from the remark at the end of this section, and then, for non-core cycles, the description detailing the relationship between moving through and domino insertion of [5](2.3.2) carries without modification to the arbitrary rank case. □

Example 4.11 Let $w = (4, -3, -2, 1) \in D_{IN}^3(W_4).$ Then $T_{IN}^3(w) = (4, -3, 1, -2)$ and the corresponding tableau pairs are:

$$G_2(w) = \left(\begin{array}{|c|c|c|} \hline & & 1 \\ \hline & 3 & 4 \\ \hline 2 & & \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline & & 1 \\ \hline & 3 & 4 \\ \hline 2 & & \\ \hline \end{array} \right) \text{ and}$$

$$G_2(T_{IN}^3(w)) = \left(\begin{array}{|c|c|c|} \hline & & 1 \\ \hline & 3 & 4 \\ \hline 2 & & \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline & & 1 \\ \hline & 3 & \\ \hline 2 & 4 & \\ \hline \end{array} \right)$$

The extended cycle through $r + 2 = 4$ in $T_2(w)$ consists of the open cycle $\{4\}$ in $T_2(w)$ and the corresponding open cycle $\{4\}$ in $T_1(w).$

Remark 4.12 In the case when $T_2(r + 2),$ is not sparse it is not immediately clear that the operation on tableaux described in Proposition 4.10 actually produces a domino tableau of rank $r.$ That it does follows from the fact that the extended cycle through which the tableaux are being moved through does not contain any core open cycles. We verify this presently. Suppose that $w \in T_{IN}^{r+1},$ so that $w(r + 1)$ and $w(r + 2)$ are of opposite sign.

Case 1. Suppose either $\tilde{E}_0(r + 1)$ or $\tilde{E}_1(r + 1)$ appears in $T_2,$ and without loss of generality, assume that it is $\tilde{E}_0(r + 1).$ The domino with label $r + 2$ is adjacent to $r + 1$ as well as to another domino adjacent to the core of $T_2,$ which we label $l.$ For $r + 2$ to be in an extended cycle containing a core open cycle, the extended cycle must also contain either $r + 1$ or $l.$ We show that this is impossible. First, $r + 1$ and $r + 2$ cannot be in the same extended cycle; one of them is boxed but not the other. Since $w(r + 1)$ and $w(r + 2)$ are of opposite sign, it is easy to check that *within* $T_2(r + 2),$ $r + 2$ and l are in different extended cycles. But Lemmas 3.6 and 3.7 of [15] imply that this remains the case within the full tableau $T_2,$ see also [5](2.3.3).

Case 2. Suppose either $E_0(r + 1)$ or $E_1(r + 1)$ appears in T_2 , and without loss of generality, assume that it is $E_0(r + 1)$. The first operation prescribed by Proposition 4.10 is to swap $E_0(r + 1)$ with $E_1(r + 1)$ in T_2 . If we now define l as before, then $r + 2$ and l must be in different extended cycles as one of them is boxed but not the other. Since $w(r + 1)$ and $w(r + 2)$ are of opposite sign, it is again easy to check that within $(T_2(r + 2) \setminus E_0(r + 1)) \cup E_1(r + 1)$, $r + 2$ and $r + 1$ are in different extended cycles, and the rest of the proof follows as above.

5 Generators of combinatorial cells

The goal of this section is to verify that the set of operators in Λ^{r+1} generates the reducible combinatorial right cells of rank r in the hyperoctahedral group W_n . We state Theorem 3.9 more precisely as:

Theorem 5.1 *Suppose that $w, v \in W_n$ with $G_r(w) = (T_1, T_2)$ and $G_r(v) = (\tilde{T}_1, \tilde{T}_2)$. If $\tilde{T}_1 = MT(T_1, U)$ for some set of non-core open cycles U in T_1 , then there is a sequence of operators Σ in the set Λ^{r+1} such that $\Sigma(w) = v$.*

Its proof follows directly from two auxiliary facts. However, we need a definition first.

Definition 5.2 We will say that a domino tableau is *somewhat special* if all of its non-core open cycles are boxed.

A tableau is somewhat special iff all of its corners are empty, thus this is really a property of the underlying partition. Since moving through a non-core open cycle changes it from boxed to unboxed and vice-versa, there is a unique somewhat special tableau in the family of tableaux obtained from each other by moving through non-core open cycles. We will write $S(T)$ for the somewhat special tableau corresponding to T .

Proposition 5.3 *Given $w \in W_n$, there is a sequence of operators Σ in Λ^{r+1} such that the left tableau of $\Sigma(w)$ is somewhat special.*

Proposition 5.4 *Suppose that $w, v \in W_n$ with $G_r(w) = (T_1, T_2)$ and $G_r(v) = (\tilde{T}_1, \tilde{T}_2)$. If $T_1 = \tilde{T}_1$ and both are somewhat special, then there is a sequence of operators Σ in the set Λ^{r+1} such that $\Sigma(w) = v$.*

The verification of these two propositions will occupy the remaining sections of this paper. Our main approach to their proof is inductive; to check Propositions 5.3 and 5.4, we will assume that the proposition itself as well as Theorem 3.9 is true for smaller n . In this section, we will show how Theorem 3.9 follows from Proposition 5.3 and Proposition 5.4 once they are known to hold, as well as check the base case for the induction. Section 6 contains the proof of Proposition 5.4 while Section 7 contains the proof of Proposition 5.3 and an auxiliary lemma which also relies on a similar induction.

We now show how the main theorem follows from the two propositions. We will write Σ for a sequence of operators, Σ^{-1} for the same sequence taken in the opposite order, and $\Sigma_2 \Sigma_1$ for the sequence of operators obtained from sequences Σ_1 and Σ_2 by applying the operators in Σ_1 first followed by the ones from Σ_2 .

Proof Since $\tilde{T}_1 = MT(T_1, U)$ for some set of non-core open cycles U , the definition of somewhat special implies that $S(T_1) = S(\tilde{T}_1)$. According to Proposition 5.3, there are sequences of operators Σ_1 and Σ_2 in Λ^{r+1} such that

1. $\Sigma_1(w)$ has left tableau $S(T_1)$, and
2. $\Sigma_2(v)$ has left tableau $S(\tilde{T}_1)$

Since $S(T_1) = S(\tilde{T}_1)$, Proposition 5.4 implies that there is another sequence Σ_3 in Λ^{r+1} such that $\Sigma_3(\Sigma_1(w)) = \Sigma_2(v)$. In other words,

$$\Sigma_2^{-1} \Sigma_3 \Sigma_1(w) = v. \quad \square$$

The following lemma in a sense serves as the base case for our inductive approach to Propositions 5.3 and 5.4. We show that family of operators Λ^{r+1} is enough to generate combinatorial left cells in two special cases when n is small when compared to r . In what follows, we will identify group elements with their corresponding tableau pairs and apply the operators in Λ^{r+1} directly to them via the action described in Section 3.2.

Lemma 5.5 *Let $\tilde{\Lambda}^{r+1}$ consist of the operators $\Lambda^{r+1} \setminus \{T_{SC}^{r+1}\}$. Then*

1. $\tilde{\Lambda}^{r+1}$ generates the irreducible, and consequently the reducible combinatorial left cells of rank r in W_n when $n \leq r + 1$, and
2. Λ^{r+1} generates the reducible combinatorial left cells of rank r in W_{r+2} .

Proof The first part is just a restatement of Theorem 3.11, proved by C. Bonnafé and L. Iancu. So consider w, v in W_{r+2} and let

$$G_r(w) = (T_1, T_2) \text{ and } G_r(v) = (\tilde{T}_1, \tilde{T}_2)$$

so that $\tilde{T}_1 = MT(T_1, U)$ for some set U of non-core open cycles in T_1 . We would like to define a sequence Σ of operators in Λ^{r+1} so that $\Sigma(w) = v$, or identifying group elements with tableau pairs, $\Sigma(T_1, T_2) = (\tilde{T}_1, \tilde{T}_2)$.

Suppose first that $T_1 = \tilde{T}_1$ and the domino with label $r + 2$ appears in the same position in T_2 and \tilde{T}_2 . Then by the first part of the lemma, there is a sequence $\Sigma_1 \subset \Lambda^{r+1}$ so that $\Sigma_1(T_1, T_2)' = (\tilde{T}_1, \tilde{T}_2)'$. But because $T_1 = \tilde{T}_1$, this implies $\Sigma_1(T_1, T_2) = (\tilde{T}_1, \tilde{T}_2)$.

Next, suppose that $T_1 = \tilde{T}_1$ and the domino with label $r + 2$ appears in different positions in T_2 and \tilde{T}_2 . We will say that a tableau of rank r is full if all of its squares of the form $S_{i,r+3-i}$ are occupied. First suppose that T_2 is not full. We will show that there are sequences Σ'_2 and Σ''_2 so that $\Sigma'_2(T_1, T_2)$ and $\Sigma''_2(\tilde{T}_1, \tilde{T}_2)$ have the $r + 2$ domino in the same position; because of what we have already proved, this will be sufficient. Because T_2 and \tilde{T}_2 are not full, either their top rows or first columns must

have two dominos. Without loss of generality, assume that it is the top rows, and furthermore, assume that $r + 2$ is not in the top row of T_2 , for otherwise we could simply swap (T_1, T_2) and $(\tilde{T}_1, \tilde{T}_2)$. By the first part of the lemma, there is a sequence Σ'_2 so that $\Sigma'_2(T_1, T_2)'$ has the dominos r and $r + 1$ in the top row. However, note that $T_{\alpha_{r+1}\alpha_{r+2}}$ now can swap the $r + 1$ and $r + 2$ dominos in the right tableau of $\Sigma'_2(T_1, T_2)$. If $r + 2$ is in the top row of \tilde{T}_2 , then we are done. If not, we can repeat the above procedure with $(\tilde{T}_1, \tilde{T}_2)$ to form a sequence Σ''_2 .

Next, suppose that $T_1 = \tilde{T}_1$, the domino with label $r + 2$ appears in different positions in T_2 and \tilde{T}_2 , and T_2 is full. Let Σ'_3 be the sequence of operators in Λ^{r+1} which arranges the entries of w in decreasing order of absolute values; because T_2 is full, the sequences of positive and negative entries in w are both decreasing in absolute value and so such a Σ'_3 always exists. By construction, $\Sigma'_3(T_1, T_2)$ lies in the domain of T_{SC} . If T_{IN}^{r+1} is not an element of the sequence Σ'_3 , then the left tableau of $\Sigma'_3(T_1, T_2)$ is T_1 and the sought-after sequence of operators is $T_{SC}\Sigma'_3$. If T_{IN}^{r+1} is in Σ'_3 , then the left tableau of $\Sigma'_3(T_1, T_2)$ differs from T_1 by moving through the cycle $\{r + 2\}$. Then, it is always possible to find a sequence $\Sigma''_3 \subset \Lambda^r$ so that $T_{SC}\Sigma''_3(T_1, T_2)$ is in the domain of T_{IN}^{r+1} , perhaps after interchanging the roles of w and v . The desired sequence is then $T_{IN}^{r+1}\Sigma''_3T_{SC}\Sigma'_3$.

Finally, suppose $T_1 \neq \tilde{T}_1$. The only possible non-core cycle in T_1 is $\{r + 2\}$ so that $D(r + 2, \tilde{T}_1) = D'(r + 2, T_1)$. Because of what we have already proved, it will be enough to find a sequence Σ_4 so that $\Sigma_4(T_1, T_2)$ and $(\tilde{T}_1, \tilde{T}_2)$ are of the same shape. Note that because T_1 and \tilde{T}_1 are of different shapes, either w or v must contain some entries with opposite signs. Without loss of generality, assume that it is w ; then it is possible to find a sequence $\Sigma'_4 \subset \Lambda^r$ so that $\Sigma'_4(T_1, T_2)$ is in the domain of T_{IN}^{r+1} . The desired sequence is then $\Sigma_4 = T_{IN}^{r+1}\Sigma'_4$. □

6 Somewhat special cells

The goal of this section is to verify Proposition 5.4 and prove a few auxiliary facts which we will need for Proposition 5.3, whose proof we defer to the next section. We will say that a sequence Σ in Λ^{r+1} is inductive if it does not contain the operators $T_{\alpha_{n-1}\alpha_n}$ and $T_{\alpha_n\alpha_{n-1}}$, or the operators T_{SC}^{n-1} and T_{IN}^{n-1} . The first lemma describes what happens to the maximal domino in the right tableau of $\Sigma(w)$ for an inductive sequence Σ in Λ^{r+1} and is essentially [6] (3.2.7). While the statement of the result is the same, minor adaptations in the proof are necessary in this more general setting. We relate the entire proof for completeness.

Lemma 6.1 *Consider an inductive sequence Σ of operators in Λ^{r+1} , and let $G_r(w) = (T_1, T_2)$ and $G_r(\Sigma(w)) = (\tilde{T}_1, \tilde{T}_2)$. Then:*

1. $D(n, \tilde{T}_2)$ is either $D(n, T_2)$ or its image under moving through.
2. If $\tilde{T}_2(n - 1)$ and $T_2(n - 1)$ have the same shape, then $D(n, \tilde{T}_2) = D(n, T_2)$.
3. If $D(n, T_2)$ is boxed and $\tilde{T}_2(n - 1)$ is somewhat special, then so is \tilde{T}_2 .

Proof The proof of part one proceeds by induction on the size of Σ . If Σ consists of one operator, then because Σ is inductive the operators $T_{\alpha\beta}$, T_{SC}^{r+1} and T_{IN}^k for $k \leq r$ (when inductive) leave $D(n, T_2)$ untouched. The operator T_{IN}^{r+1} , when inductive, changes $D(n, T_2)$ into $D'(n, T_2)$ iff n is in the extended cycle affected by its action. The inductive step is an immediate consequence of the fact that n is the maximal entry in T_2 and that moving through is an involution.

The other two parts rely on the following consequence of the condition $D(n, \tilde{T}_2) \neq D(n, T_2)$. If this is the case, then either

1. for some $k < n$ the positions of the dominos $D(k, \tilde{T}_2)$ and $D(n, T_2)$ intersect, so that k is in the same cycle as n , or
2. for some $k < n$ there are non-core open cycles c, d in \tilde{T}_2 and a cycle e in \tilde{T}_1 with $S_b(e) = S_b(d) \in D(k, \tilde{T}_2)$ and $S_f(e) = S_f(c) \in D(n, T_2)$, implying that k is in the same extended cycle as n .

Both possibilities imply that the shapes of $\tilde{T}_2(n - 1)$ and $T_2(n - 1)$ are not the same, which proves the second part of the lemma. For the third part, we will show that $D(n, \tilde{T}_2) = D(n, T_2)$, which will imply that \tilde{T}_2 is somewhat special. We will argue by contradiction, so assume that $D(n, \tilde{T}_2) = D'(n, T_2)$ instead. If (1) holds, then the boxing condition implies that $D(k, \tilde{T}_2) \cap D(n, T_2)$ is a filled corner in $\tilde{T}_2(n - 1)$, contradicting the hypothesis that $\tilde{T}_2(n - 1)$ is somewhat special. On the other hand, if (2) holds then $D(n, \tilde{T}_2)$ is unboxed as is its entire extended cycle in \tilde{T}_2 . Then $S_b(d) \in D(k, \tilde{T}_2)$ is a filled corner $\tilde{T}_2(n - 1)$, contradicting the hypothesis that $T_2'(n - 1)$ is somewhat special. □

The next lemma shows that inductive sequences behave well with respect to domino insertion.

Lemma 6.2 *Let Σ in A^{r+1} be an inductive sequence and let $(T_1, T_2)'$ be the tableau pair obtained by deleting the highest-numbered domino from T_2 and reversing one step of the insertion procedure for T_1 . Then*

$$\Sigma((T_1, T_2)') = (\Sigma(T_1, T_2))'$$

Proof It is enough to verify this statement when Σ consists of just one operator. When the operator is of the form $T_{\alpha\beta}$, T_{IN}^k for $k \leq r$, and T_{SC}^{r+1} for $r + 2 < n$ the result is clear. When $r + 2 > n$, T_{SC}^{r+1} is not defined, and when $r + 2 = n$, it is excluded. It remains to check the lemma for the operator T_{IN}^{r+1} under the assumption that $r + 2 < n$.

If the action of the operator T_{IN}^{r+1} merely swaps two not adjacent dominos in T_2 , then the result is again clear. If not, then its action on T_2 either swaps two domino configurations and then moves through the extended cycle through $r + 2$, or vice versa. Write T for the tableau to which the moving through will be applied. By Remark 4.12, the extended cycle through $r + 2$ in T with respect to T_1 contains only non-core cycles. Thus to show the lemma, it is only necessary to verify the following:

Lemma 6.3 *Consider an extended cycle c in T_2 relative to T_1 which contains only non-core cycles and excludes the cycle $\{n\}$, if one exists. Write $(\tilde{T}_1, \tilde{T}_2)$ for the pair*

$MT((T_1, T_2), c)$, and form the set c' by deleting the label n from any cycle in c . Write $(\tilde{T}'_1, \tilde{T}'_2) = (\tilde{T}_1, \tilde{T}_2)'$. Then

1. c' is an extended cycle in T'_2 relative to T'_1 , and
2. $(\tilde{T}'_1, \tilde{T}'_2) = MT((T_1, T_2)', c')$.

Because c contains only non-core cycles, [5](2.2.9) holds, and the proof of the lemma is identical to that of [5](2.3.3). □

We will need one more lemma to prove Proposition 5.4, which we state presently and whose proof we defer until the next section. For a somewhat special tableau, it constructs a sequence of operators whose action moves the domino with label n to any other pre-prescribed extremal position.

Lemma 6.4 *Consider (T_1, T_2) of rank r with T_2 somewhat special. If P' is a removable pair of adjoining squares in T_2 , then there is a sequence of operators $\Sigma \subset \Lambda_n$ such that the n domino of the right tableau of $\Sigma(T_1, T_2)$ is in position P' and the left tableau of $\Sigma(T_1, T_2)$ remains equal to T_1 .*

Armed with this, we are ready to prove Proposition 5.4, which we restate for the reader’s convenience.

Proposition 5.4 *Suppose that $w, v \in W_n$ with $G_r(w) = (T_1, T_2)$ and $G_r(v) = (\tilde{T}_1, \tilde{T}_2)$. If $T_1 = \tilde{T}_1$ and both are somewhat special, then there is a sequence of operators Σ in the set Λ^{r+1} such that $\Sigma(w) = v$.*

Proof We argue by induction and assume that Theorem 3.9 holds for numbers smaller than n . Let P' be the position of the n domino in \tilde{T}_2 . By Lemma 6.4, there is a sequence Σ_1 of operators so that the right tableau of $\Sigma_1(T_1, T_2) = (T_1^1, T_2^1)$ has its n domino in position P' and $T_1 = T_1^1$. Note that the left tableaux of $(T_1^1, T_2^1)'$ and $(\tilde{T}_1, \tilde{T}_2)'$ must be the same since \tilde{T}_2 and T_2^1 have their n dominos in the same position, and by hypothesis, $T_1^1 = \tilde{T}_1$. By induction, we can construct an inductive sequence Σ_2 such that $\Sigma_2(T_1^1, T_2^1)' = (\tilde{T}_1, \tilde{T}_2)'$, which, by Lemma 6.2, also equals $(\Sigma_2(T_1^1, T_2^1))'$.

We would like to show that $\Sigma_2(T_1^1, T_2^1) = (\tilde{T}_1, \tilde{T}_2)$, which will complete the proof. Since $(T_1^1, T_2^1)'$ and $(\tilde{T}_1, \tilde{T}_2)'$ must have the same shape, Lemma 6.1 implies that the n dominos of the right tableau of $\Sigma_2(T_1^1, T_2^1)$ and T_2^1 must be in the same position, which also happens to be the position of the n domino in \tilde{T}_2 . But since $\Sigma_2(T_1^1, T_2^1)' = (\tilde{T}_1, \tilde{T}_2)'$, the right tableaux of $\Sigma_2(T_1^1, T_2^1)$ and $(\tilde{T}_1, \tilde{T}_2)$ must agree. Hence their left tableaux must be of the same shape, and because $T_1^1 = \tilde{T}_1$, the rest follows. □

7 Technical lemmas

In this section, we verify Proposition 5.3 and Lemma 6.4. They are both adaptations of lemmas from [5] and follow from case by case analyses of relative domino positions.

Proposition 5.3 *Given $w \in W_n$, there is a sequence of operators Σ in Λ^{r+1} such that the left tableau of $\Sigma(w)$ is somewhat special.*

Proof Consider $w \in W_n$ and let $G_r(w) = (T_1, T_2)$. Our goal is to find a sequence $\Sigma \in \Lambda^{r+1}$ so that $\Sigma(w)$ has a somewhat special left tableau. We will superficially follow the outline of the proof of [6] (3.2.4), but the new situations in the general rank case will require a slightly different approach and our cases are somewhat different. We will use induction on n , assuming both, that Proposition 5.3 and Theorem 3.9 are true for values smaller than n .

Let $(\bar{T}_1, \bar{T}_2) = (T_1, T_2)'$. By induction, we know that there is an inductive sequence Σ_1 so that the left tableau of $\Sigma_1((T_1, T_2)') = (\bar{T}_1^1, \bar{T}_2^1)$ is somewhat special. Let $\Sigma_1(T_1, T_2) = (T_1^1, T_2^1)$. Then Lemma 6.2 implies $(T_1^1, T_2^1)' = (\bar{T}_1^1, \bar{T}_2^1)$. There are two cases:

1. If T_2^1 is somewhat special, then so is T_1^1 and we are done.
2. If T_2^1 is not somewhat special, then one of its non-core open cycles is not boxed.

But since \bar{T}_2^1 itself is somewhat special, the only possibility is that $\{n\}$ is an unboxed extended open cycle in T_2^1 relative to T_1^1 .

We assume the latter is true and without loss of generality, take the domino $D(n, T_2^1)$ to be horizontal writing $\{S_{ij}, S_{ij+1}\}$ for its underlying squares. Because $\{n\}$ is a cycle, $D(n, T_2^1)$ is unboxed, and T_2^1 is not somewhat special, S_{ij+1} must be a filled corner and $S_{i+1,j}$ must be empty hole in T_1^1 implying that $S_{i,j+1}$ is a square of type X . We will examine a number of cases in order to find a sequence of operators which will take T_2^1 to a somewhat special tableau.

Our general goal will be to find two pairs of adjacent squares P_1 and P_2 in T_2^1 , so that P_1 is extremal in $T_2^1(n - 1)$ and P_2 is extremal in $T_2^1(n - 1) \setminus P_1$. If this is possible, then induction on Theorem 3.9 provides a sequence Σ_2 so that $(\bar{T}_1^2, \bar{T}_2^2) = \Sigma_2(\bar{T}_1^1, \bar{T}_2^1)$ with $\bar{T}_1^2 = \bar{T}_1^1$ and the $n - 1$ domino of \bar{T}_2^2 is P_1 while its $n - 2$ domino is P_2 . Let $(T_1^2, T_2^2) = \Sigma_2(T_1^1, T_2^1)$. If (T_1^2, T_2^2) is in the domain of either $T_{\alpha\beta} = T_{\alpha_{n-1}\alpha_n}$ or $T_{\alpha_n\alpha_{n-1}}$, then let $(T_1^3, T_2^3) = T_{\alpha\beta}(T_1^2, T_2^2)$. Again by induction, it is then possible to find a sequence Σ_4 so that $(\bar{T}_1^4, \bar{T}_2^4) = \Sigma_4(T_1^3, T_2^3)'$ is somewhat special. Finally, let

$$(T_1^4, T_2^4) = \Sigma_4(T_1^3, T_2^3).$$

This turns out to be somewhat special, and $\Sigma = \Sigma_4(T_{\alpha\beta})\Sigma_2\Sigma_1$ is the promised sequence of operators.

If pairs of adjacent squares P_1 and P_2 cannot be suitably chosen, we will find that T_2^1 satisfies the hypotheses of Lemma 5.5, which provides the desired sequence of operators. To simplify the details, we will let a number of our cases overlap.

Case 1. Assume $i > 1$, which, because of our choice of boxing, implies that $i \geq 3$. Let s and t be the lengths of the $i - 1$ st and $i - 2$ nd rows of T_2^1 , and further assume that $t - s$ and $s - j$ are both greater than 1. Define $P_1 = \{S_{i-2,t-1}, S_{i-2,t}\}$ and $P_2 = \{S_{i-1,s-1}, S_{i-1,s}\}$. Then (T_1^2, T_2^2) lies in the domain of $T_{\alpha_{n-1}\alpha_n}$, which acts on T_2^2 by interchanging the n and $n - 1$ dominos. That (T_1^4, T_2^4) is somewhat special follows from Lemma 6.1 since \bar{T}_2^4 is somewhat special and P_1 is boxed in \bar{T}_2^1 which is also somewhat special.

Case 2. Assume the same situation as in Case 1, but suppose $t = s > j + 1$. The proof is identical to Case 1 if we let $P_1 = \{S_{i-2,t}, S_{i-1,t}\}$ and $P_2 = \{S_{i-2,t-1}, S_{i-1,t-1}\}$.

Case 3. Assume the same situation as in Case 1, but suppose $t = s = j + 1$. Define $P_1 = \{S_{i-2,t}, S_{i-1,t}\}$ and $P_2 = \{S_{i-2,t-1}, S_{i-1,t-1}\}$. The action of $T_{\alpha_{n-1}\alpha_n}$ on T_2^2 this time is not just an interchange and is described by case 1 of 4.2. That T_1^4 is somewhat special follows because \bar{T}_2^4 is somewhat special and $D(n, T_2^4)$ occupies the squares $\{S_{i-2,j+1}, S_{i-1,j+1}\}$ and therefore is boxed.

Case 4. Assume that $S_{i+2,j-1}$ is in T_2^1 . Because \bar{T}_2^1 is somewhat special, this implies $S_{i+3,j-1}$ is also in T_2^1 , otherwise \bar{T}_2^1 would have a filled corner. Let l be the length the of $j - 1$ st column of T_2^1 and define $P_1 = \{S_{l-1,j-1}, S_{l,j-1}\}$ $P_2 = \{S_{l-3,j-1}, S_{l-2,j-1}\}$. The rest of the proof follows as before, except the operator $T_{\alpha_n\alpha_{n-1}}$ must be used.

Case 5. Assume $S_{i+2,j-1}$ is not in T_2^1 , but that $S_{i+2,j-2}$ is. Let P_2 be the pair of squares $\{S_{i,j-1}, S_{i+1,j-1}\}$. The definition of P_1 is more involved. Let m be the smallest positive number so that either $\{S_{i+m+1,j-m-1}\}$ is not a square in T_2^1 or $\{S_{i+m+1,j-m}\}$ is a square in T_2^1 . We examine the former as case (a), the latter as case (b), and if neither occurs as case (c).

- (a) Define $P_1 = \{S_{i+m,j-m-1}, S_{i+m,j-m}\}$,
- (b) Let p be the length of the $(j - m)$ th column of T_2^1 and define P_1 to be the pair of squares $\{S_{p-1,j-m-1}, S_{p,j-m}\}$.

In either of these two cases, the proof follows as before using the operator $T_{\alpha_n\alpha_{n-1}}$.

- (c) In this case, no adequate extremal position for P_1 exists below $D(n, T_2^1)$. If $i = 1$, then T_2^1 in fact has to satisfy the hypotheses of Lemma 5.5, so we can assume (as described in Case (1)) that $i > 2$. We may further assume that T_2^1 does not satisfy the hypotheses of any of the previous cases, so that we can also assume $s = j + 1$ and $t = j + 2$. Define m to be the smallest positive number so that either $\{S_{i-m-1,j+m+1}\}$ is not a square in T_2^1 or $\{S_{i-m,j+m+1}\}$ is a square in T_2^1 . As before, define P_1 to be respectively $\{S_{i-m-1,j+m}, S_{i-m,j+m}\}$ or $\{S_{i-m,q-1}, S_{i-m,q}\}$ where q is the length of the $i - m$ th row of T_2^1 . In either of these two cases, the proof follows as before using the operator $T_{\alpha_{n-1}\alpha_n}$. If no such m exists however, then T_2^1 again has to satisfy the hypotheses of Lemma 5.5, which provides the necessary sequence of operators.

Case 6. Assume $j > 2$ and $S_{i+2,j-2} \notin T_2^1$. Note that since $S_{i+2,j-2} \notin T_2^1$, the square $S_{i,j-1}$ cannot lie in the core of T_2^1 . Thus let $P_1 = \{S_{i+1,j-2}, S_{i+1,j-1}\}$ and $P_2 = \{S_{i,j-2}, S_{i,j-1}\}$. The proof proceeds as before, with $D(n, T_2^4) = D'(n, T_2^3) = \{S_{i+1,j-1}, S_{i+1,j}\}$.

Case 7. Assume $j = 2$ and $S_{i+2,j-1} \notin T_2^1$. If we assume that T_2^1 satisfies the hypotheses of none of the previous cases, then let $P_2 = \{S_{i,j-1}, S_{i+1,j-1}\}$ and the proof proceeds exactly as in Case 5(c).

Case 8. Assume $j = 1$, so that i cannot equal 1, implying $i > 2$. If we assume that T_2^1 satisfies the hypotheses of none of the other cases above, then we can take $s = 2$ and $t = 3$. Let $P_2 = \{S_{i-1,1}, S_{i-1,2}\}$. Again, the proof proceeds exactly as in

Case 5(c). It is a quick check to see that our cases cover all possibilities, and so the proof of the lemma is complete. \square

The following lemma is a general rank version of [6] (3.2.9). We adapt its proof to the general rank case. There are two complications: the existence of a nontrivial core and the fact that the tableau is only somewhat special, so we cannot count on all holes being filled. Almost entirely, cases which do not match those in the rank zero proof will degenerate into the setting of Lemma 5.5, where we proved the main result for tableaux whose rank is large relative to n .

Lemma 6.4 *Consider (T_1, T_2) of rank r with T_2 somewhat special. If P' is a removable pair of adjoining squares in T_2 , then there is a sequence of operators $\Sigma \subset \Lambda_n$ such that the n domino of the right tableau of $\Sigma(T_1, T_2)$ is in position P' and the left tableau of $\Sigma(T_1, T_2)$ remains equal to T_1 .*

Proof The proof is by induction where we assume both, that Lemma 6.4 and Theorem 3.9 are true for values smaller than n . Write P for the set of squares occupied by the n domino in T_2 and assume that $P \neq P'$. Without loss of generality assume that P is horizontal, writing $P = \{S_{ij}, S_{i,j+1}\}$.

Cases 1-4. For the first cases, also assume that $P' = \{S_{kl}, S_{k,l+1}\}$ is horizontal. We first define an auxiliary domino position P_1 .

1. If $k = i - 1$, we must have $l \geq j + 2$ since P' is extremal. If $S_{i-1,j}$ is in the core of T_2 , then $l \geq j + 3$. In either case, let $P_1 = \{S_{i-1,l-2}, S_{i-1,l-1}\}$.
2. If $k < i - 1$, let u be the length of the $(i - 1)$ st row of T_2 . Then $l \geq u + 1$ and we can set $P_1 = \{S_{i-1,u-1}, S_{i-1,u}\}$.
3. If $k = i + 1$, then $l \leq j - 2$. Let $P_1 = \{S_{i,j-2}, S_{i,j-1}\}$. Note that if $S_{i,j-2}$ is in the core of T_2 , then P' could not have been extremal.
4. If $k > i + 1$, then let u be the length of the $(k - 1)$ st row of T_2 . Let $P_1 = \{S_{k-1,u-1}, S_{k-1,u}\}$.

Proceeding by induction as in the proof of Proposition 5.3, we can find an inductive sequence Σ_1 so that $\Sigma_1(T_1, T_2)'$ has the same left tableau as $(T_1, T_2)'$ and whose right tableau has domino $n - 1$ in position P' , and $n - 2$ in position P_1 . By Lemma 6.1, $\Sigma_1(T_1, T_2)$ has left tableau T_1 . The desired sequence is then $T_{\alpha_{n-1}, \alpha_n} \Sigma_1$ in the first two cases and $T_{\alpha_n, \alpha_{n-1}} \Sigma_1$ in the second two.

Cases 5-7. For the rest of the cases assume that $P' = \{S_{kl}, S_{k+1,l}\}$ is vertical. We again define an auxiliary domino position P_1 .

- (5) If $k + 1 = i - 1$, let $P_1 = \{S_{i-1,l-2}, S_{i-1,l-1}\}$ as in Case (1).
- (6) If $k + 1 < i - 1$, let $P_1 = \{S_{i-1,u-1}, S_{i-1,u}\}$ as in Case (2).
- (7) If $k = i + 1$ and $l < j - 1$, let $P_1 = \{S_{i,j-2}, S_{i,j-1}\}$ as in Case (3).

The proof proceeds exactly as in the first four cases, with $T_{\alpha_{n-1}, \alpha_n}$ applied in Cases 5 and 6, and $T_{\alpha_n, \alpha_{n-1}}$ in Case 7.

Case 8. Again assume that $P' = \{S_{kl}, S_{k+1,l}\}$ and suppose that $k > i + 1$.

- i. Suppose that P'' is an extremal domino shape in $T_2 \setminus P \setminus P'$. If P'' is above P' but not above P , let $P_1 = P'$ and proceed as in Case (4). If P'' is below P' and

horizontal, then using cases (1)-(4) we can find a sequence so that the resulting right tableau will have the n domino in position P'' . Then we are in the setting of cases (5)-(7). If P'' is below P' and vertical, by induction it is possible to find a sequence so that the resulting right tableau will have the $n - 1$ domino in position P' and the $n - 2$ domino in position P'' . After applying $T_{\alpha_n, \alpha_{n-1}}$, we are in the setting of the transpose of cases (1)-(4). If P'' is above P and also extremal in T_2 itself, then arguments similar to the above apply.

- ii. Suppose that $P'' = \{S_{k+1, l-2}, S_{k+1, l-1}\}$ is extremal in $T_2 \setminus P'$ and $P''' = \{S_{k, l-2}, S_{k, l-1}\}$ is extremal in $T_2 \setminus P' \setminus P''$. By induction it is possible to find a sequence so that the resulting right tableau will have the $n - 1$ domino in position P' , $n - 2$ domino in position P'' , and the $n - 3$ domino in position P''' . After applying $T_{\alpha_{n-2}, \alpha_{n-1}}$, we are in the setting of cases (1)-(4). A similar argument works if
 - (a) $P'' = \{S_{k, l-1}, S_{k+1, l-1}\}$ is extremal in $T_2 \setminus P'$ and the domino shape $P''' = \{S_{k-1, l-1}, S_{k-1, l}\}$ is extremal in $T_2 \setminus P' \setminus P''$.
 - (b) $P'' = \{S_{i-2, j+1}, S_{i-1, j+1}\}$ is extremal in $T_2 \setminus P$ and the domino shape $P''' = \{S_{i-2, j}, S_{i-1, j}\}$ is extremal in $T_2 \setminus P \setminus P''$.
 - (c) $P'' = \{S_{i-1, j}, S_{i-1, j+1}\}$ is extremal in $T_2 \setminus P$ and the domino shape $P''' = \{S_{i-1, j-1}, S_{i, j-1}\}$ is extremal in $T_2 \setminus P \setminus P''$.

If there is no extremal P'' in $T_2 \setminus P \setminus P'$ above P' but not above P , $S_{i+1, j-2} \in T_2$, but $S_{i+1, j-1} \notin T_2$. It then follows by an easy but tedious inspection that if we are not in any of the above cases, then we must be within the scope of Lemma 5.5.

Case 9. Assume that P and P' are both boxed and that $k = i - 1$, so that P and P' intersect in the square $S_{i, j+1}$. Since they are both boxed, we know that S_{ij+1} must be of type W , and consequently, that i is even.

- i. Assume that $i > 2$, and hence that $i \geq 4$. Let u be the length of the row $i - 2$ of T_2 and let s be the length of row $i - 3$.
 - (a) If $u = j + 1$, let $P_1 = \{S_{i-1, j+1}, S_{i-2, j+1}\}$ and $P_2 = \{S_{i-1, j}, S_{i-2, j}\}$. Note that $S_{i-2, j}$ cannot be in the core of T_2 , since $D(n, T_2)$ occupies the square S_{ij} . By induction, we can find a sequence Σ_1 so that the $n - 1$ domino in the right tableau of $\Sigma_1(T_1, T_2)'$ is P_1 and the $n - 2$ domino is P_2 . Then the n domino of $T_{\alpha_{n-1}, \alpha_n} \Sigma_1(T_1, T_2)$ occupies the squares $\{S_{i, j+1}, S_{i-1, j+1}\}$, as desired.
 - (b) Assume that $s = u > j + 1$. If we let $P'' = \{S_{i-3, u}, S_{i-2, u}\}$, then Case 6 provides a sequence Σ_1 so that the right tableau of $\Sigma_1(T_1, T_2)$ has the n domino in position P'' . The transpose of the first two cases now provides a sequence Σ_2 such that the n domino of $\Sigma_2 \Sigma_1(T_1, T_2)$ is P' , as desired.
 - (c) Assume that $s > u > j + 1$. If there is an extremal domino shape P'' in T_2 strictly above P , we can use the already considered cases (or their transposes), to find a sequence Σ_1 so that $\Sigma_1(T_1, T_2)$ has left tableau T_1 and the n domino is in position P'' in its right tableau. Then, again by the already considered cases (or their transposes), we can find a sequence Σ_2 so that $\Sigma_2 \Sigma_1(T_1, T_2)$ has left tableau T_1 and the n domino is in position P' in its right tableau, as desired. A similar argument works if there is an extremal domino shape P'' in T_2 strictly below P . If neither is the case, then we are in the scope of Lemma 5.5, which provides the necessary sequence.

ii. If $i = 2$ and $j > 2$, then this case is parallel to the previous one. If $i = 2$ and $j = 2$, then we are either in the setting of Lemma 5.5 or Case 8. Finally, if $i = 2$ and $j = 1$, we are in rank zero, which has been considered in [6].

Case 10. Assume that P and P' are both boxed and that $k = i + 1$ and $l = j - 1$. The boxing condition implies that $S_{i,j+1}$ must be of type W , and consequently, that i is even. Let r be the length of the $i - 1$ st row of T_2 . If $u = j + 1$, then let $P'' = \{S_{i-1,j+1}, S_{i,j+1}\}$. By Case 9, we can find a sequence Σ_1 so that $\Sigma_1(T_1, T_2)$ has left tableau T_1 and its right tableau has the n domino in position P'' . Then Case 2 provides a sequence Σ_2 so that $\Sigma_2\Sigma_1(T_1, T_2)$ has left tableau T_1 and its right tableau has the n domino in position P' . If $u > j + 1$, then because T_2 is somewhat special it cannot have filled corners, so that $u \geq j + 3$. But then we can let $P'' = \{S_{i-1,r-1}, S_{i,r}\}$ and the proof is the same as when $r = j + 1$.

Case 11. Assume that P' is boxed and P is not. Our goal is to find a sequence of operators which will move the n domino to a boxed position, reducing our work to the previous cases. Because T_2 is somewhat special, it cannot have filled corners, so that S_{ij} must be of type W and consequently, $i \geq 2$. Let u be the length of the $i - 1$ row of T_2 . If $u = j + 1$, let Σ_1 be the sequence of operators such that the right tableau of $\Sigma_1(T_1, T_2)'$ is somewhat special; it exists by induction and Proposition 5.3. The shape of the tableaux $\Sigma_1(T_1, T_2)'$ is $(\text{shape}(T_2) \setminus S_{i-1,j+1}) \cup S_{ij}$. If T_2^1 is the right tableau of $\Sigma_1(T_1, T_2)$, then Lemma 6.1, implies that $D(n, T_2^1) = D'(n, T_1) = \{S_{i-1,j+1}, S_{i,j+1}\}$, which is boxed, as desired. If $u > j + 1$, then because T_2 is somewhat special and cannot have filled corners, $u \geq j + 3$. Let $P'' = \{S_{i-1,u-1}, S_{i-1,u}\}$. Then from Cases 1 through 4, where a boxing condition was not assumed, we can find a sequence of operators Σ_1 so that the n domino in the right tableau of $\Sigma_1(T_1, T_2)$ is P'' , which, because T_2 is special, must be boxed.

Case 12. Assume that P' is unboxed in T_2 . Since Cases 1 through 8 were verified without any boxing assumptions, we only have to check this case when

- i. $k = i - 1$ so that P and P' intersect in the square $S_{i,j+1}$, or
- ii. $k = i + 1$ and $l = j - 1$.

Furthermore, we can also take P to be boxed, as otherwise the boxing conditions imply in both cases that $S_{i,j+1}$ is a filled corner, contradicting the assumption that T_2 is somewhat special. With this assumption, the transpose of (i) is the last case considered in [6](3.2.9), and we omit its proof. So assume that we are in case (ii). Then the type of S_{ij} is X .

- (a) Assume that there are no extremal domino positions strictly below P' or above P . If $i > 1$ and $S_{i-1,j+2}$ is not a square in T_2 , or if $S_{i+2,j-2} \in T_2$ and $S_{i+3,j-2} \notin T_2$, then we can use case (i) above to find a sequence Σ_1 so that the right tableau of $\Sigma_1(T_1, T_2)$ has the n domino in position $\{S_{i,j+1}, S_{i-1,j+1}\}$. Then $\Sigma_1(T_1, T_2)$ is in the setting of Case 2. If neither of the two possibilities above is true, then we are in the setting of Lemma 5.5.
- (b) If there is an extremal domino position P'' strictly below P' , then using induction we can find a sequence Σ_1 so that the right tableau of $\Sigma_1(T_1, T_2)$ has domino $n - 1$ in position P'' and $n - 2$ in position P' . Then $T_{\alpha_n \alpha_{n-1}} \Sigma_1(T_1, T_2)$ is within scope of the previous cases. If there is an extremal domino position P' strictly

above P , then by the previous cases, there is a sequence Σ_1 so that the n domino of the right tableau of $\Sigma_1(T_1, T_2)$ is P'' . We can use induction to find a sequence Σ_2 which puts the $n - 2$ domino in position P and the $n - 1$ domino in position P' . Then the desired sequence is $T_{\alpha_n \alpha_{n-1}} \Sigma_2 \Sigma_1(T_1, T_2)$. \square

References

1. Ariki, S.: Robinson-Schensted correspondence and left cells. In: Combinatorial methods in representation theory, Kyoto, 1998. Adv. Stud. Pure Math., vol. 28, pp. 1–20. Kinokuniya, Tokyo (2000)
2. Bonnafé, C., Iancu, L.: Left cells in type B_n with unequal parameters. Represent. Theory **7**, 587–609 (2003)
3. Bonnafé, C., Geck, M., Iancu, L., Lam, T.: On domino insertion and Kazhdan–Lusztig cells in type B_n . In: Progress in Math. (Lusztig Birthday Volume). Birkhauser, Boston (to appear)
4. Garfinkle, D.: On the classification of primitive ideals for complex classical Lie algebras (I). Compositio Math. **75**(2), 135–169 (1990)
5. Garfinkle, D.: On the classification of primitive ideals for complex classical Lie algebras (II). Compositio Math. **81**(3), 307–336 (1992)
6. Garfinkle, D.: On the classification of primitive ideals for complex classical Lie algebras (III). Compositio Math. **88**(2), 187–234 (1993)
7. Geck, M.: Computing Kazhdan-Lusztig cells for unequal parameters. J. Algebra **281**(1), 342–365 (2004)
8. Gordon, I.G., Martino, M.: Calogero-Moser Space, reduced rational Cherednik algebras and two-sided cells. arXiv:[math.RT/0703153](https://arxiv.org/abs/math.RT/0703153)
9. Hopkins, B.: Domino tableaux and single-valued wall-crossing operators. Ph.D. dissertation, University of Washington (1997)
10. Kazhdan, D., Lusztig, G.: Representations of Coxeter groups and Hecke algebras. Invent. Math. **53**(2), 165–184 (1979)
11. Knuth, D.E.: The art of computer programming, vol. 3. Addison-Wesley Publishing Co., Reading (1973)
12. Lusztig, G.: Hecke algebras with unequal parameters. CRM Monograph Series, vol. 18. American Mathematical Society, Providence
13. McGovern, W.M.: On the Spaltenstein-Steinberg map for classical Lie algebras. Comm. Algebra **27**(6), 2979–2993 (1999)
14. Pietraho, T.: Components of the Springer fiber and domino tableaux. J. Algebra **272** (2), 711–729 (2004)
15. Pietraho, T.: A relation for domino Robinson-Schensted algorithms. Ann. Comb. (to appear)
16. Pietraho, T.: Equivalence classes in the Weyl groups of type B_n . J. Algebraic Comb. **27**(2), 247–262 (2008)
17. Pietraho, T.: Cells and constructible representations in type B_n . arXiv:[math.CO/0710.3846](https://arxiv.org/abs/math.CO/0710.3846)
18. Stanton, D.W., White, D.E.: A Schensted algorithm for rim hook tableaux. J. Combin. Theory Ser. A **40**(2), 211–247 (1985)
19. Taskin, M.: Plactic relations for r -domino tableaux. arXiv:[math.RT/0803.1148](https://arxiv.org/abs/math.RT/0803.1148)
20. van Leeuwen, M.A.A.: The Robinson-Schensted and Schützenberger algorithms, an elementary approach. Electron. J. Comb. **3**(2) (1996)
21. Vogan, D.: A generalized τ -invariant for the primitive spectrum of a semisimple Lie algebra. Math. Ann. **242**(3), 209–224 (1979)