

A Note on the Homology of Signed Posets

PHIL HANLON*

Department of Mathematics, University of Michigan, Ann Arbor, MI 48109-1003

Received February 7, 1994; Revised May 17, 1995

Abstract. Let S be a signed poset in the sense of Reiner [4]. Fischer [2] defines the homology of S , in terms of a partial ordering $P(S)$ associated to S , to be the homology of a certain subcomplex of the chain complex of $P(S)$.

In this paper we show that if $P(S)$ is Cohen-Macaulay and S has rank n , then the homology of S vanishes for degrees outside the interval $[n/2, n]$.

Keywords: poset, Cohen-Macaulay, signed poset

1. Introduction

Let R be a set of vectors in \mathbb{R}^n . The *positive linear closure* of R , denoted \bar{R} is defined to be the span of all linear combinations of vectors in R with non-negative real coefficients.

For each $i = 1, 2, \dots, n$ let e_i denote the i th unit coordinate vector in \mathbb{R}^n and let e_{-i} denote $-e_i$. Recall that the root system B_n is the set

$$B_n = \{\pm(e_i \pm e_j) : 1 \leq i < j \leq n\} \cup \{\pm e_i : 1 \leq i \leq n\}.$$

Definition 1 A *signed poset* is a subset S of B_n such that

- (a) $S \cap (-S) = \emptyset$.
- (b) $\bar{S} \cap B_n = S$.

Let (P, \leq) be an ordinary poset with $P = \{1, 2, \dots, n\}$. Let S be the collection of all $e_i - e_j$ such that $i < j$. Then S is a subset of the root system A_n which satisfies conditions (a) and (b) of Definition 1 (where B_n is replaced by A_n in condition (b)). Vic Reiner introduced the notion of signed poset [4] to be a B_n -analogue of the notion of poset.

In more recent work Steve Fischer [2] defined a homology theory for signed posets. According to Fischer's definition, the homology of a signed poset S is the homology of a certain simplicial complex $C_*^0(S)$ associated to S . This simplicial complex is analogous to the simplicial complex of chains in a poset. Fischer showed that the Euler characteristic of this homology can be computed via a "2-Mobius function" and that analogues of Weisner's Theorem and Crapo's Complementation Theorem can be used to calculate this 2-Mobius

*Research partially supported by the National Science Foundation and the John Simon Guggenheim Foundation.

function when S is a “signed lattice”. In view of these results on the 2-Mobius function, it would be interesting to know if there are combinatorial labelling conditions which would imply that the simplicial complex associated to S is shellable.

There is an obvious analogue of EL-labelling that can be defined for signed posets, namely we say that S is EL-labellable iff $P(S)$ is EL-labellable. Here $P(S)$ is a poset whose chains are used to define $C_*^0(S)$. An EL-labellable signed poset is pure in the sense that all facets of $C_*^0(S)$ have the same dimension (which we will call the dimension of S). Originally Fischer had hoped to show that if S is EL-labellable then the homology of $C_*^0(S)$ is zero except in the top dimension. But then he constructed two EL-labellable signed posets S_0 and S_1 such that

- (a) the homology of $C_*^0(S_0)$ is nonzero exactly in degree equal to half the top dimension.
- (b) the homology of $C_*^0(S_1)$ is nonzero exactly in degree equal to the top dimension.

He went on to define “signed EL-labelling” to be an EL-labelling that satisfies other conditions and showed that the existence of a signed EL-labelling of S implies that $C_*^0(S)$ is shellable.

The purpose of this note is to prove that the examples S_0 and S_1 above are the extreme cases, i.e., we will prove.

Theorem 1 *Suppose S is an EL-labellable signed poset of dimension n . Then $H_r(S)$ is 0 unless $\lfloor n/2 \rfloor \leq r \leq n$.*

2. Homology of a signed poset

We begin this section by defining the simplicial complex $C_*^0(S)$ that Fischer uses to compute the homology of S . This complex is given in terms of the chains in a certain poset $P(S)$.

Definition 2 Let S be a signed poset in B_n . Define the poset $P(S)$ with vertex set $\{\pm 1, \dots, \pm n\} = V$ as follows. For $u, v \in V$ we say

$$u \leq_{P(S)} v$$

if and only if

- (i) $e_u - e_v \in S$ for $|u| \neq |v|$ or
- (ii) $e_u \in S$ for $v = -u$.

Fischer showed that $P(S)$ is a self dual poset.

Definition 3 An isotropic r -chain in $P(S)$ is an r -chain

$$\alpha_1 < \alpha_2 < \dots < \alpha_r$$

such that α_i is not equal to $-\alpha_j$ for any i, j . Let $\Delta_r^0(S)$ denote the collection of isotropic r -chains in $P(S)$ and let $C_r^0(S)$ denote the \mathbb{C} -span of $\Delta_r^0(S)$ (with $C_0^0(S) = \mathbb{C}$).

Note that $\Delta_r^0(S)$ is a simplicial complex in $2^{P(S)}$. This gives a boundary map $\partial_r: C_r^0(S) \rightarrow C_{r-1}^0(S)$,

$$\partial_r(\alpha_1 < \alpha_2 < \cdots < \alpha_r) = \sum_{i=1}^r (-1)^{i-1} (\alpha_1 < \cdots < \hat{\alpha}_i < \cdots < \alpha_r).$$

Definition 4 Define $H_r^0(S)$ to be the r th homology of the complex $(C_*^0(S), \partial_*)$, i.e.

$$H_r^0(S) = \ker \partial_r / \text{im } \partial_{r+1}.$$

We call $H_*^0(S)$ the *signed poset homology* of S .

We say S is *EL-labellable* if $P(S)$ has an EL-labelling. In [2], Fischer computes $C_*^0(S)$ and $H_*^0(S)$ for a number of signed posets S . In particular he constructs a family of posets $\Gamma_n \subseteq B_n$ such that:

- $\Delta^0(\Gamma_n)$ is pure of dimension n
- Γ_n is EL-labellable
- $\Delta^0(\Gamma_n)$ is homotopic to the $\lfloor n/2 \rfloor$ -dimensional sphere.

This family of signed posets shows that an EL-labelling on S does not imply that $\Delta^0(S)$ is shellable.

3. The main result

Let Q be a finite, ranked, self-dual poset. Let $x \rightarrow x^*$ be a fixed order-reversing involution on Q . Split $Q = Q^L \cup Q^U$ so that Q^L is an order ideal in Q , $(Q^L)^* \cap Q^L = \{x \in Q : x^* = x\}$ and $(Q^U)^* \subseteq Q^L$. For each chain $\gamma = \alpha_1 < \alpha_2 < \cdots < \alpha_r$, define $\omega(\gamma)$ to be the number of pairs (α_i, α_j) with $i < j$ and $\alpha_j = \alpha_i^*$. We say γ is *isotropic* if $\omega(\gamma) = 0$.

Let $C_r(Q)$ denote the span of all r -chains and $C_*^0(Q)$ the span of all isotropic r -chains. The boundary map $\partial_*: C_*(Q) \rightarrow C_{*-1}(Q)$ preserves $C_*^0(Q)$ and so $(C_*^0(Q), \partial_*)$ is a subcomplex of $(C_*(Q), \partial_*)$. Let $H_*^0(Q)$ denote the homology of that subcomplex. The main theorem for this section is:

Theorem 2 *Suppose Q is Cohen-Macaulay of rank n . Then*

$$H_d^0(Q) = 0 \quad \text{unless } \frac{n}{2} \leq d \leq n.$$

Proof: We prove this by induction on $|Q|$. If Q is the empty poset then $H_d^0(Q)$ is 0 unless $d = 0$. This agrees with the statement in Theorem 2 since $n = 0$ in this case.

Consider an arbitrary Q and assume the result is true for all Q' with $|Q'| < |Q|$.

Let γ be a chain in $C_*(Q)$. We assign a non-negative integer $\rho(\gamma)$ to γ as follows:

- 1) If γ is isotropic then $\rho(\gamma) = 0$.
- 2) If γ is not isotropic, write γ as

$$\alpha_1 < \alpha_2 < \cdots < \alpha_r.$$

Then $\rho(\gamma)$ is the rank of α_i where i is maximal subject to the condition that $\alpha_i^* = \alpha_j$ for some $j > i$. We also write $A(\gamma)$ to denote α_i . Note that $A(\gamma) \in Q^L$.

For $r, p \in \mathbb{N}$ let $C_{r,p}(Q)$ denote the span of all r -chains γ with $\rho(\gamma) = p$. Note that the boundary map ∂ satisfies:

$$\partial(C_{r,p}(Q)) \subseteq \bigoplus_{t \leq p} C_{r-1,t}(Q).$$

Thus $(C_*(Q), \partial)$ is filtered by the parameter ρ . Let (E^s, ∂^s) be the associated spectral sequence which abuts to $E^\infty = H_*(Q)$. Background material on spectral sequences can be found in any introductory text in homological algebra (e.g. [1] or [3]).

Our first step will be to compute the E^1 term in this spectral sequence.

E^0 is the associated graded complex. Let γ be an r -chain in E_r^0 . Write γ as

$$\gamma = \alpha_1 < \alpha_2 < \cdots < \alpha_i < \alpha_{i+1} < \cdots < \alpha_{j-1} < \alpha_j = \alpha_i^* < \alpha_{j+1} < \cdots < \alpha_r.$$

Then

$$\partial^0 \gamma = \sum_{s=1, s \neq i, j}^r (-1)^{s-1} (\alpha_1 < \cdots < \hat{\alpha}_s < \cdots < \alpha_r). \quad (1)$$

Let $E_r^0[\alpha]$ denote the span of all r -chains γ with $A(\gamma) = \alpha$ and let $E_r^0[\hat{0}]$ denote $C_r^0(Q)$. Then

- 1) $E_r^0 = C_r^0(Q) \oplus \bigoplus_{\alpha \in Q^L \setminus \{\hat{0}\}} E_r^0[\alpha]$
- 2) $\partial^0(E_r^0[\alpha]) \subseteq E_{r-1}^0[\alpha]$ for all $\alpha \in Q^L \cup \{\hat{0}\}$.

So the complex (E_r^0, ∂^0) splits as a direct sum of the subcomplexes

$$\bigoplus_{\alpha \in Q^L \setminus \{\hat{0}\}} (E_*^0[\alpha], \partial^0).$$

We now analyze the subcomplex $(E_*^0[\alpha], \partial^0)$. Assume $\alpha \in Q^L$ and that $\alpha^* > \alpha$. For a chain γ to have $A(\gamma) = \alpha$, it is necessary and sufficient for γ to consist of any chain up to α , then an isotropic chain α to α^* , and then any chain from α^* upward. So,

$$E_{*+2}^0[\alpha] \cong C_*(I_\alpha) \otimes C_*((\alpha, \alpha^*)) \otimes C_*(I^{\alpha^*}) \quad (2)$$

where I_α denotes the open order ideal generated by α in Q , I^{α^*} denotes the open order filter generated by α^* in Q and (α, α^*) is the open interval from α to α^* in Q . Moreover, (1)




Figure 1.

shows that the tensor product of vector spaces given by (2) extends to a tensor product of complexes.

Let p be the rank of α so the rank of α^* is $n + 1 - p$. Since Q is Cohen-Macaulay we have

$$H_d(I_\alpha) = H_d(I^{\alpha^*}) = 0 \quad \text{unless } d = p - 1.$$

The self-dual poset (α, α^*) is Cohen-Macaulay of rank $(n - p) - p = n - 2p$. By our induction hypothesis

$$H_d^0((\alpha, \alpha^*)) = 0 \quad \text{unless } \frac{n - 2p}{2} \leq d \leq n - 2p.$$

Combining these observations we find:

$$E_d^1[\alpha] = 0 \quad \text{unless } \frac{n}{2} + p \leq d \leq n.$$

At this point we know nothing about

$$E_*^1[\hat{0}] = \text{Homology of } (C_*^0(Q), \partial) = H_*^0(Q).$$

However we can draw a diagram of $E_{r,p}^1$ letting a square box denote values of r, p where $E_{r,p}^1$ might be non-zero. This appears in Figure 1.

The ∂^1 differential on E^1 maps $E_{r,p}^1$ to $E_{r-1,p-1}^1$. More generally, the ∂^s differential on E^s maps $E_{r,p}^s$ to $E_{r-1,p-s}^s$. It follows by induction on s that

$$E_{r,0}^s = E_{r,0}^1 = E_r^1[\hat{0}] = H_r^0(Q)$$

for $0 \leq r < \frac{n}{2}$ and all s . Thus

$$H_r^0(Q) = E_{r,0}^\infty \subseteq H_r(Q) = 0 \quad \text{for } 0 \leq r < \frac{n}{2}.$$

This proves Theorem 2. □

Theorem 1 follows immediately from Theorem 2 by taking $Q = P(S)$.

4. Other problems

The question answered by Theorem 2 has an obvious generalization. Let C be a simplicial complex, pure of dimension n , with vertex set V and let $G \subseteq \text{Sym}(V)$ be a group of automorphisms of C . Let C^0 be the collection of all faces of V which do not contain two elements of V from the same orbit.

Question Suppose C is shellable. What can you say about the dimensions t where $H_t(C^0)$ is nonzero?

References

1. H. Cartan and S. Eilenberg, *Homological Algebra*, Oxford University Press, Oxford, 1956.
2. S. Fischer, "Signed poset homology and q -analog Mobius functions," preprint.
3. P.J. Hilton and U. Stambach, *A Course in Homological Algebra*, Springer Graduate Texts in Mathematics, Springer-Verlag, 1971.
4. V. Reiner, "Signed posets," *JCTA* **62**(2) (1993), 324–360.