

DIAGONAL FIBRATIONS ARE POINTWISE FIBRATIONS

A.M. CEGARRA AND J. REMEDIOS

(communicated by Ronald Brown)

Abstract

On the category of bisimplicial sets there are different Quillen closed model structures associated to various definitions of fibrations. In one of them, which is due to Bousfield and Kan and that consists of seeing a bisimplicial set as a simplicial object in the category of simplicial sets, fibrations are those bisimplicial set maps such that each of the induced simplicial set maps is a Kan fibration, that is, the pointwise fibrations. In another of them, introduced by Moerdijk, a bisimplicial map is a fibration if it induces a Kan fibration of associated diagonal simplicial sets, that is, the diagonal fibrations. In this note, we prove that every diagonal fibration is a pointwise fibration.

1. Introduction and summary

There are several (Quillen) closed model structures on the category of bisimplicial sets, see [3, IV, §3]. This paper concerns two of them, namely, the so-called Bousfield-Kan and Moerdijk structures, that we briefly recall below:

On the one hand, in the closed model structure by Bousfield-Kan, bisimplicial sets are regarded as diagrams of simplicial sets and then fibrations are the pointwise Kan fibrations and weak equivalences are the pointwise weak homotopy equivalences. To be more precise, a bisimplicial set $X : \Delta^{op} \times \Delta^{op} \rightarrow \mathbf{Set}$, $([p], [q]) \mapsto X_{p,q}$, is seen as a “horizontal” simplicial object in the category of “vertical” simplicial sets, $X : \Delta^{op} \rightarrow \mathbf{S}$, $[p] \mapsto X_{p,*}$ and then, a bisimplicial map $f : X \rightarrow Y$ is a fibration (resp. a weak equivalence) if all simplicial maps $f_{p,*} : X_{p,*} \rightarrow Y_{p,*}$, $p \geq 0$, are Kan fibrations (resp. weak homotopy equivalences).

On the other hand, the Moerdijk closed model structure on the bisimplicial set category is transferred from the ordinary model structure on the simplicial set category through the diagonal functor, $X \mapsto \text{diag } X : [n] \mapsto X_{n,n}$. Thus, in this closed model structure, a bisimplicial map $f : X \rightarrow Y$ is a fibration (resp. a weak equivalence) if the induced diagonal simplicial map $\text{diag } f : \text{diag } X \rightarrow \text{diag } Y$ is a Kan fibration (resp. a weak homotopy equivalence).

Supported by DGI of Spain: (MTM2004-01060 and MTM2006-06317), FEDER, and Junta de Andalucía: P06-FQM-1889.

Received August 17, 2006, revised March 21, 2007; published on December 27, 2007.

2000 Mathematics Subject Classification: 55U10, 5P05, 55U35

Key words and phrases: bisimplicial set, closed model structure, fibration, weak homotopy equivalence

© 2007, A.M. Cegarra and J. Remedios. Permission to copy for private use granted.

Several useful relationships between these two different homotopy theories of bisimplicial sets have been established and, perhaps, the best known of them is the following:

Theorem. (Bousfield-Kan) *Let $f : X \rightarrow Y$ be a bisimplicial map such that $f_{p,*} : X_{p,*} \rightarrow Y_{p,*}$ is a weak homotopy equivalence for each $p \geq 0$. Then $\text{diag } f : \text{diag } X \rightarrow \text{diag } Y$ is a weak homotopy equivalence.*

The purpose of this brief note is to state and prove a suitable counterpart to Bousfield-Kan's theorem for fibrations, namely:

Theorem 1. *Let $f : X \rightarrow Y$ be a bisimplicial map such that $\text{diag } f : \text{diag } X \rightarrow \text{diag } Y$ is a Kan fibration. Then $f_{p,*} : X_{p,*} \rightarrow Y_{p,*}$ is a Kan fibration for each $p \geq 0$.*

Note that the converse of Theorem 1 is not true in general. A counterexample is given in the last section of the paper.

Acknowledgements. The authors are much indebted to the referees, whose useful observations greatly improved our exposition. The second author is grateful to the Algebra Department in the University of Granada for the excellent atmosphere and hospitality.

2. Some preliminaries

We use the standard conventions and terminology which can be found in texts on simplicial homotopy theory, e. g. [3] or [6]. For definiteness or emphasis we state the following.

We denote by Δ the category of finite ordered sets of integers $[n] = \{0, 1, \dots, n\}$, $n \geq 0$, with weakly order-preserving maps between them. The category of *simplicial sets* is the category of functors $X : \Delta^{op} \rightarrow \text{Set}$, where Set is the category of sets. If X is a simplicial set and $\alpha : [m] \rightarrow [n]$ is a map in Δ , then we write $X_n = X[n]$ and $\alpha^* = X(\alpha) : X_n \rightarrow X_m$. Recall that all maps in Δ are generated by the injections $\delta_i : [n-1] \rightarrow [n]$ (cofaces), $0 \leq i \leq n$, which miss out the i th element and the surjections $\sigma_i : [n+1] \rightarrow [n]$ (codegeneracies), $0 \leq i \leq n$, which repeat the i th element (see [5, VII, §5, Proposition 2]). Thus, in order to define a simplicial set, it suffices to give the sets of n -simplices X_n , $n \geq 0$, together with maps

$$\begin{aligned} d_i &= \delta_i^* : X_n \rightarrow X_{n-1}, & 0 \leq i \leq n & \quad (\text{the face maps}), \\ s_i &= \sigma_i^* : X_n \rightarrow X_{n+1}, & 0 \leq i \leq n & \quad (\text{the degeneracy maps}), \end{aligned}$$

satisfying the well-known basic *simplicial identities* such as $d_i d_j = d_{j-1} d_i$ if $i < j$, etc. (see [5, p. 175]). In addition, we shall write down a list of other identities between some iterated compositions of face and degeneracy maps, which will be used latter. The proof of these equalities is straightforward and left to the reader.

Lemma 1. *On any simplicial set, the following equalities hold:*

$$d_i d_j^m = d_j^m d_{i+m} \quad \text{if} \quad i \geq j \quad (1)$$

$$d_i^m = d_i^{m-1} d_j \quad \text{if} \quad i \leq j < i + m \quad (2)$$

$$d_i s_j^m = s_j^m d_{i-m} \quad \text{if} \quad i > j + m \quad (3)$$

$$d_i s_j^m = s_j^{m-1} \quad \text{if} \quad j \leq i \leq j + m. \quad (4)$$

Let $f : X \rightarrow Y$ be a simplicial map. A collection of simplices

$$x_i \in X_{n-1}, \quad i \in I,$$

$$y \in Y_n,$$

where $I \subseteq [n]$ is any subset, is said to be *f-compatible* whenever the following equalities hold:

$$d_i x_j = d_{j-1} x_i \quad \text{for all} \quad i, j \in I, i < j,$$

$$d_i y = f x_i \quad \text{for all} \quad i \in I.$$

The map f is said to be a *Kan fibration* whenever for every given collection of *f-compatible* simplices

$$x_0, \dots, x_{k-1}, x_{k+1}, \dots, x_n \in X_{n-1},$$

$$y \in Y_n,$$

there is a simplex $x \in X_n$ such that $d_i x = x_i$ for all $i \neq k$ and $f x = y$. The next lemma (cf. [6, Lemma 7.4]) will be very useful in our development. For I any finite set, $|I|$ denotes its number of elements.

Lemma 2. *Let $f : X \rightarrow Y$ be a Kan fibration. Suppose that there are given a subset $I \subseteq [n]$ such that $1 \leq |I| \leq n$ and an *f-compatible* family of simplices*

$$x_i \in X_{n-1}, \quad i \in I,$$

$$y \in Y_n.$$

Then, there exists $x \in X_n$ such that $d_i x = x_i$ for all $i \in I$ and $f x = y$.

Proof. Suppose $|I| = r$. If $r = n$, the statement is true since f is a Kan fibration. Hence the statement holds for $n = 1$. We now proceed by induction: Assume $n > 1$ and the result holds for $n' < n$ and assume $r < n$ and the result holds for $r' > r$. Taking $k = \max\{i \mid i \in [n], i \notin I\}$, we wish to find a simplex $x_k \in X_{n-1}$, such that the collection of simplices

$$x_i \in X_{n-1}, \quad i \in I \cup \{k\},$$

$$y \in Y_n,$$

be *f-compatible*, since then an application of the induction hypothesis on r gives the claim. To find such an x_k , let $I' \subseteq [n-1]$ be the subset

$$I' = \{i \mid i \in I, i < k\} \cup \{i-1 \mid i \in I, i > k\},$$

and let

$$\begin{aligned} x'_i &\in X_{n-2}, & i &\in I', \\ y' &\in Y_{n-1}, \end{aligned}$$

be the family of simplices defined by $x'_i = d_{k-1}x_i$ for $i \in I, i < k, x'_{i-1} = d_kx_i$ for $i \in I, i > k$, and $y' = d_ky$. This family is f -compatible, whence the induction hypothesis on $n - 1$ gives the required x_k , that is, a $(n - 1)$ -simplex of X satisfying $d_ix_k = x'_i$ for $i \in I'$ and $fx_k = y'$. \square

The category of *bisimplicial sets* is the category of functors $X : \Delta^{op} \times \Delta^{op} \rightarrow \text{Set}$. It is often convenient to see a bisimplicial set X as a (horizontal) simplicial object in the category of (vertical) simplicial sets. If $\alpha : [p] \rightarrow [p']$ and $\beta : [q] \rightarrow [q']$ are any two maps in Δ , then we will write $\alpha^{*h} : X_{p',q} \rightarrow X_{p,q}$ and $\beta^{*v} : X_{p,q'} \rightarrow X_{p,q}$ for the images $X(\alpha, id)$ and $X(id, \beta)$ respectively. In particular, the horizontal and vertical face and degeneracy maps are $d_i^h = (\delta_i)^{*h}, d_i^v = (\delta_i)^{*v}, s_i^h = (\sigma_i)^{*h}$ and $s_i^v = (\sigma_i)^{*v}$.

By composing with the diagonal functor $\Delta \rightarrow \Delta \times \Delta, [n] \mapsto ([n], [n])$, we get the *diagonal functor* from bisimplicial sets to simplicial sets, which provides a simplicial set $\text{diag } X : [n] \mapsto X_{n,n}$, associated to each bisimplicial set X , whose face and degeneracy operators are given in terms of those of X by the formulas $d_i = d_i^h d_i^v$ and $s_i = s_i^h s_i^v$, respectively.

3. Proof of Theorem 1

Let $p \geq 0$ be any fixed integer. Then, in order to prove that the simplicial set map $f_{p,*} : X_{p,*} \rightarrow Y_{p,*}$ is a Kan fibration, suppose that, for some integers $q \geq 1$ and $0 \leq \ell \leq q$, there is given a collection of bisimplices

$$\begin{aligned} x_0, \dots, x_{\ell-1}, x_{\ell+1}, \dots, x_q &\in X_{p,q-1}, \\ y &\in Y_{p,q}, \end{aligned}$$

which is $f_{p,*}$ -compatible, that is, such that $d_i^v x_j = d_{j-1}^v x_i$ for $0 \leq i < j \leq q, i \neq \ell \neq j$, and $d_i^v y = fx_i$ for $0 \leq i \leq q, i \neq \ell$. We must therefore find a bisimplex $x \in X_{p,q}$ such that

$$\begin{aligned} d_i^v x &= x_i & \text{for } 0 \leq i \leq q, i \neq \ell, \\ fx &= y. \end{aligned} \tag{5}$$

To do that, we start by considering the subset

$$I \subseteq [p + q]$$

defined by

$$I = \{i \mid 0 \leq i < \ell\} \cup \{p + i \mid \ell < i \leq q\},$$

and the family of diagonal bisimplices

$$\begin{aligned} \bar{x}_i &\in (\text{diag } X)_{p+q-1}, & i \in I, \\ \bar{y} &\in (\text{diag } Y)_{p+q}, \end{aligned} \tag{6}$$

where

$$\bar{x}_i = (s_0^h)^{\ell-1} (s_p^h)^{q-\ell} (s_{\ell-1}^v)^p x_i \quad \text{for } 0 \leq i < \ell, \tag{7}$$

$$\bar{x}_{p+i} = (s_0^h)^\ell (s_p^h)^{q-\ell-1} (s_\ell^v)^p x_i \quad \text{for } \ell < i \leq q, \tag{8}$$

$$\bar{y} = (s_0^h)^\ell (s_p^h)^{q-\ell} (s_\ell^v)^p y. \tag{9}$$

The next verifications show that this family (6) of diagonal bisimplices is actually *diag f*-compatible:

- for $0 \leq i < j < \ell$,

$$\begin{aligned} d_i \bar{x}_j &= d_i^h d_i^v (s_0^h)^{\ell-1} (s_p^h)^{q-\ell} (s_{\ell-1}^v)^p x_j \\ &= d_i^h (s_0^h)^{\ell-1} (s_p^h)^{q-\ell} d_i^v (s_{\ell-1}^v)^p x_j \\ &= (s_0^h)^{\ell-2} (s_p^h)^{q-\ell} (s_{\ell-2}^v)^p d_i^v x_j && \text{by (4)} \\ &= (s_0^h)^{\ell-2} (s_p^h)^{q-\ell} (s_{\ell-2}^v)^p d_{j-1}^v x_i \\ &= (s_0^h)^{\ell-2} (s_p^h)^{q-\ell} d_{j-1}^v (s_{\ell-1}^v)^p x_i \\ &= d_{j-1}^h (s_0^h)^{\ell-1} (s_p^h)^{q-\ell} d_{j-1}^v (s_{\ell-1}^v)^p x_i && \text{by (4)} \\ &= d_{j-1}^h d_{j-1}^v (s_0^h)^{\ell-1} (s_p^h)^{q-\ell} (s_{\ell-1}^v)^p x_i \\ &= d_{j-1} \bar{x}_i. \end{aligned}$$

- for $0 \leq i < \ell < j \leq q$,

$$\begin{aligned} d_i \bar{x}_{p+j} &= d_i^h d_i^v (s_0^h)^\ell (s_p^h)^{q-\ell-1} (s_\ell^v)^p x_j \\ &= d_i^h (s_0^h)^\ell (s_p^h)^{q-\ell-1} d_i^v (s_\ell^v)^p x_j \\ &= (s_0^h)^{\ell-1} (s_p^h)^{q-\ell-1} (s_{\ell-1}^v)^p d_i^v x_j && \text{by (4)} \\ &= (s_0^h)^{\ell-1} (s_p^h)^{q-\ell-1} (s_{\ell-1}^v)^p d_{j-1}^v x_i \\ &= (s_0^h)^{\ell-1} d_{p+j-\ell}^h (s_p^h)^{q-\ell} (s_{\ell-1}^v)^p d_{j-1}^v x_i && \text{by (4)} \\ &= d_{p+j-1}^h (s_0^h)^{\ell-1} (s_p^h)^{q-\ell} d_{p+j-1}^v (s_{\ell-1}^v)^p x_i && \text{by (3)} \\ &= d_{p+j-1}^h d_{p+j-1}^v (s_0^h)^{\ell-1} (s_p^h)^{q-\ell} (s_{\ell-1}^v)^p x_i \\ &= d_{p+j-1} \bar{x}_i. \end{aligned}$$

- for $\ell < i < j \leq q$,

$$\begin{aligned}
 d_{p+i}\bar{x}_{p+j} &= d_{p+i}^h(s_0^h)^\ell(s_p^h)^{q-\ell-1}d_{p+i}^v(s_\ell^v)^p x_j \\
 &= (s_0^h)^\ell d_{p+i-\ell}^h(s_p^h)^{q-\ell-1}(s_\ell^v)^p d_i^v x_j && \text{by (3)} \\
 &= (s_0^h)^\ell(s_p^h)^{q-\ell-2}(s_\ell^v)^p d_{j-1}^v x_i && \text{by (4)} \\
 &= (s_0^h)^\ell d_{p+j-\ell-1}^h(s_p^h)^{q-\ell-1}(s_\ell^v)^p d_{j-1}^v x_i && \text{by (4)} \\
 &= d_{p+j-1}^h(s_0^h)^\ell(s_p^h)^{q-\ell-1}d_{p+j-1}^v(s_\ell^v)^p x_i && \text{by (3)} \\
 &= d_{p+j-1}^h d_{p+j-1}^v(s_0^h)^\ell(s_p^h)^{q-\ell-1}(s_\ell^v)^p x_i \\
 &= d_{p+j-1}\bar{x}_{p+i}.
 \end{aligned}$$

- for $0 \leq i < \ell$,

$$\begin{aligned}
 d_i\bar{y} &= d_i^h d_i^v(s_0^h)^\ell(s_p^h)^{q-\ell}(s_\ell^v)^p y \\
 &= d_i^h(s_0^h)^\ell(s_p^h)^{q-\ell}d_i^v(s_\ell^v)^p y \\
 &= (s_0^h)^{\ell-1}(s_p^h)^{q-\ell}(s_{\ell-1}^v)^p d_i^v y && \text{by (4)} \\
 &= (s_0^h)^{\ell-1}(s_p^h)^{q-\ell}(s_{\ell-1}^v)^p f x_i \\
 &= f(s_0^h)^{\ell-1}(s_p^h)^{q-\ell}(s_{\ell-1}^v)^p x_i \\
 &= f\bar{x}_i.
 \end{aligned}$$

and, finally,

- for $\ell < i \leq q$,

$$\begin{aligned}
 d_{p+i}\bar{y} &= d_{p+i}^h d_{p+i}^v(s_0^h)^\ell(s_p^h)^{q-\ell}(s_\ell^v)^p y \\
 &= d_{p+i}^h(s_0^h)^\ell(s_p^h)^{q-\ell}d_{p+i}^v(s_\ell^v)^p y \\
 &= (s_0^h)^\ell d_{p+i-\ell}^h(s_p^h)^{q-\ell}(s_\ell^v)^p d_i^v y && \text{by (3)} \\
 &= (s_0^h)^\ell(s_p^h)^{q-\ell-1}(s_\ell^v)^p d_i^v y && \text{by (4)} \\
 &= (s_0^h)^\ell(s_p^h)^{q-\ell-1}(s_\ell^v)^p f x_i \\
 &= f(s_0^h)^\ell(s_p^h)^{q-\ell-1}(s_\ell^v)^p x_i \\
 &= f\bar{x}_{p+i}.
 \end{aligned}$$

Then, since by hypothesis $\text{diag } f : \text{diag } X \rightarrow \text{diag } Y$ is a Kan fibration, from Lemma 2 we get a diagonal bisimplex

$$\bar{x} \in (\text{diag } X)_{p+q}$$

such that $d_i^h d_i^v \bar{x} = \bar{x}_i$ for $0 \leq i < \ell$, $d_{p+i}^h d_{p+i}^v \bar{x} = \bar{x}_{p+i}$ for $\ell < i \leq q$ and $f\bar{x} = \bar{y}$.

Now, using the bisimplex \bar{x} we construct the bisimplex

$$x = (d_{p+1}^h)^{q-\ell} (d_0^h)^\ell (d_\ell^v)^p \bar{x} \in X_{p,q},$$

which, we claim, satisfies Relations (5). Actually:

- for $0 < i < \ell$,

$$\begin{aligned} d_i^v x &= d_i^v (d_{p+1}^h)^{q-\ell} (d_0^h)^\ell (d_\ell^v)^p \bar{x} \\ &= (d_{p+1}^h)^{q-\ell} (d_0^h)^\ell d_i^v (d_\ell^v)^p \bar{x} \\ &= (d_{p+1}^h)^{q-\ell} (d_0^h)^\ell (d_{\ell-1}^v)^p d_i^v \bar{x} \\ &= (d_{p+1}^h)^{q-\ell} (d_0^h)^{\ell-1} d_i^h (d_{\ell-1}^v)^p d_i^v \bar{x} && \text{by (2)} \\ &= (d_{p+1}^h)^{q-\ell} (d_0^h)^{\ell-1} (d_{\ell-1}^v)^p d_i^h d_i^v \bar{x} \\ &= (d_{p+1}^h)^{q-\ell} (d_0^h)^{\ell-1} (d_{\ell-1}^v)^p \bar{x}_i \\ &= (d_{p+1}^h)^{q-\ell} (d_0^h)^{\ell-1} (d_{\ell-1}^v)^p (s_0^h)^{\ell-1} (s_p^h)^{q-\ell} (s_{\ell-1}^v)^p x_i && \text{by (7)} \\ &= (d_{p+1}^h)^{q-\ell} (d_0^h)^{\ell-1} (s_0^h)^{\ell-1} (s_p^h)^{q-\ell} (d_{\ell-1}^v)^p (s_{\ell-1}^v)^p x_i \\ &= x_i. \end{aligned}$$

- for $\ell < i \leq q$,

$$\begin{aligned} d_i^v x &= d_i^v (d_{p+1}^h)^{q-\ell} (d_0^h)^\ell (d_\ell^v)^p \bar{x} \\ &= (d_{p+1}^h)^{q-\ell} (d_0^h)^\ell d_i^v (d_\ell^v)^p \bar{x} \\ &= (d_{p+1}^h)^{q-\ell} (d_0^h)^\ell (d_\ell^v)^p d_{p+i}^v \bar{x} && \text{by (1)} \\ &= (d_{p+1}^h)^{q-\ell-1} d_{p+i-\ell}^h (d_0^h)^\ell (d_\ell^v)^p d_{p+i}^v \bar{x} && \text{by (2)} \\ &= (d_{p+1}^h)^{q-\ell-1} (d_0^h)^\ell d_{p+i}^h (d_\ell^v)^p d_{p+i}^v \bar{x} && \text{by (1)} \\ &= (d_{p+1}^h)^{q-\ell-1} (d_0^h)^\ell (d_\ell^v)^p d_{p+i}^h d_{p+i}^v \bar{x} \\ &= (d_{p+1}^h)^{q-\ell-1} (d_0^h)^\ell (d_\ell^v)^p (s_0^h)^\ell (s_p^h)^{q-\ell-1} (s_\ell^v)^p x_i && \text{by (8)} \\ &= (d_{p+1}^h)^{q-\ell-1} (d_0^h)^\ell (s_0^h)^\ell (s_p^h)^{q-\ell-1} (d_\ell^v)^p (s_\ell^v)^p x_i \\ &= x_i. \end{aligned}$$

and, finally,

$$\begin{aligned}
 fx &= f(d_{p+1}^h)^{q-\ell}(d_0^h)^\ell(d_\ell^v)^p \bar{x} \\
 &= (d_{p+1}^h)^{q-\ell}(d_0^h)^\ell(d_\ell^v)^p f \bar{x} \\
 &= (d_{p+1}^h)^{q-\ell}(d_0^h)^\ell(d_\ell^v)^p \bar{y} \\
 &= (d_{p+1}^h)^{q-\ell}(d_0^h)^\ell(d_\ell^v)^p (s_0^h)^\ell (s_p^h)^{q-\ell} (s_\ell^v)^p y && \text{by (9)} \\
 &= (d_{p+1}^h)^{q-\ell}(d_0^h)^\ell (s_0^h)^\ell (s_p^h)^{q-\ell} (d_\ell^v)^p (s_\ell^v)^p y \\
 &= y.
 \end{aligned}$$

Hence Theorem 1 is proved.

Let us stress that Theorem 1 not only says that, when $\text{diag } f : \text{diag } X \rightarrow \text{diag } Y$ is a Kan fibration, all simplicial maps $f_{p,*} : X_{p,*} \rightarrow Y_{p,*}$ are Kan fibrations, but that the simplicial maps $f_{*,p} : X_{*,p} \rightarrow Y_{*,p}$ are Kan fibrations as well. This is a consequence of the symmetry of the hypothesis: “ $\text{diag } f$ is a Kan fibration”, that is, the fact follows by exchanging the vertical and horizontal directions.

4. The converse of Theorem 1 is false

It is possible that, for a bisimplicial map $f : X \rightarrow Y$, all simplicial maps $f_{p,*}$ and $f_{*,p}$, $p \geq 0$, be Kan fibrations and, however, $\text{diag } f$ be not a Kan fibration: Take X to be the double nerve of an suitable double groupoid (i.e. a groupoid object in the category of groupoids); then, all simplicial sets $X_{p,*}$ and $X_{*,p}$ are Kan complexes (since they are nerves of groupoids) but, as we show below, the diagonal simplicial set $\text{diag } X$ is not necessarily a Kan complex.

We briefly recall some standard terminology about double groupoids; see for example [2] or [4]. A double groupoid consists of objects a, b, \dots , horizontal and vertical morphisms between them

$$a \xrightarrow{f} b, \quad \begin{array}{c} b \\ \uparrow_g \\ a \end{array}, \dots,$$

and squares σ, τ , etc., of the form

$$\begin{array}{ccc} c & \xleftarrow{f} & b \\ g' \uparrow & \sigma & \uparrow g \\ d & \xleftarrow{f'} & a \end{array} .$$

These satisfy axioms such that the objects together with the horizontal morphisms as well as the objects together with the vertical morphisms form groupoids. Furthermore, the squares form a groupoid under both horizontal and vertical juxtaposition,

and in the situation

$$\begin{array}{cccc}
 \cdot < \cdot < \cdot < \cdot < \cdot \\
 \uparrow \sigma & \uparrow \tau & \uparrow & \uparrow \\
 \cdot < \cdot < \cdot < \cdot < \cdot \\
 \uparrow \gamma & \uparrow \delta & \uparrow & \uparrow \\
 \cdot < \cdot < \cdot < \cdot < \cdot
 \end{array}$$

horizontal and vertical composition commute in the sense that

$$(\sigma \cdot_h \tau) \cdot_v (\gamma \cdot_h \delta) = (\sigma \cdot_v \gamma) \cdot_h (\tau \cdot_v \delta).$$

In addition to the vertical identity morphisms, there are horizontal identity squares for each vertical morphism. Its horizontal edges are the identity arrows of the horizontal groupoid of arrows. Similarly, there are vertical identity squares for each horizontal morphism with vertical identity arrows. These identity squares are compatible in the sense that vertical and horizontal identity squares for the vertical and horizontal identity morphisms are the same.

Given a double groupoid \mathbb{G} , one can construct its bisimplicial nerve (or double nerve) NNG . A typical (p, q) simplex of NNG is a subdivision of a square of \mathbb{G} as matrix of $p \times q$ horizontally and vertically composable squares of the form

$$\begin{array}{ccccccc}
 \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot \\
 \uparrow \sigma_{11} & \uparrow \sigma_{21} & \uparrow & \uparrow \sigma_{p1} & \uparrow & & \uparrow \\
 \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot \\
 \uparrow \sigma_{12} & \uparrow \sigma_{22} & \uparrow & \uparrow \sigma_{p2} & \uparrow & & \uparrow \\
 \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot \\
 & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
 \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot \\
 \uparrow \sigma_{1q} & \uparrow \sigma_{2q} & \uparrow & \uparrow \sigma_{pq} & \uparrow & & \uparrow \\
 \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot
 \end{array}$$

The bisimplicial face operators are induced by horizontal and vertical composition of squares, and degeneracy operators by appropriated identity squares. We picture NNG so that the set of (p, q) -simplices is the set in the p -th row and q -th column. Thus, the p -th column of NNG , $\text{NNG}_{p,*}$, is the nerve of the “vertical” groupoid whose objects are strings of p composable horizontal morphisms $(a_0 \xleftarrow{f_1} a_1 \xleftarrow{f_2} \dots \xleftarrow{f_p} a_p)$ and whose arrows are depicted as

$$\begin{array}{cccc}
 \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot \\
 \uparrow \sigma_1 & \uparrow \sigma_2 & \uparrow & \uparrow \sigma_p & \uparrow \\
 \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot < \cdot
 \end{array}$$

And, similarly, the q -th row, $\text{NNG}_{*,q}$, is the “horizontal” groupoid whose objects are the length q sequences of composable vertical morphisms in \mathcal{C} and whose arrows are sequences of q vertically composable squares. In particular, $\text{NNG}_{0,*}$ and $\text{NNG}_{*,0}$ are, respectively, the nerves of the groupoids of vertical and horizontal morphisms of \mathbb{G} .

Example. Let A, B be subgroups of a finite group G , such that $AB \neq BA$, where

$$AB = \{ab \mid a \in A, b \in B\} \text{ and } BA = \{ba \mid a \in A, b \in B\}$$

(for instance, we can take $G = S_3$, $A = \{id, (1, 2)\}$ and $B = \{id, (1, 3)\}$).

Then, let $\mathcal{C} = \mathcal{C}(A, B)$ denote the double groupoid with only one object, say “.”, with horizontal morphisms $\cdot \xleftarrow{a} \cdot$ the elements $a \in A$, with vertical morphisms $\cdot \uparrow_b \cdot$ are the elements b of B , both with composition given by multiplication in the group, and whose squares

$$\begin{array}{ccc} \cdot & \xleftarrow{a} & \cdot \\ b' \uparrow & & \uparrow b \\ \cdot & \xleftarrow{a'} & \cdot \end{array}$$

are lists (a, b, a', b') of elements $a, a' \in A$ and $b, b' \in B$ such that $ab = b'a'$. Compositions of squares are defined, in the natural way, by

$$(a, b, a', b') \cdot_h (a_1, b_1, a'_1, b_1) = (aa_1, b_1, a'_1 a', b'),$$

$$(a, b, a', b') \cdot_v (a', b_1, a'_1, b'_1) = (a, bb_1, a'_1, b'b'_1),$$

and identities by

$$id^v(a) = (a, e, a, e), \quad id^h(b) = (e, b, e, b),$$

where e is the neutral element of the group.

So defined, we claim that the associated simplicial set diag NNC is not a Kan complex. To prove that, note that the hypothesis $AB \neq BA$ implies the existence of elements $a \in A$ and $b \in B$ such that ab cannot be expressed in the form $b'a'$, for any $a' \in A$ and $b' \in B$. Now, the identity squares $\iota_b = id^h(b)$ and $\iota_a = id^v(a)$ can be placed jointly as in the picture:

$$\begin{array}{ccc} \cdot & \xleftarrow{a} & \cdot \\ \parallel & \iota_a & \parallel \\ \cdot & \xleftarrow{a} & \cdot \\ & b \uparrow & \iota_b \uparrow b \\ & \cdot & \cdot \end{array}$$

This means that, regarded ι_a and ι_b as 1-simplices of diagonal simplicial set diag NNC , they are compatible, in the sense that

$$d_0^h d_0^v \iota_a = \cdot = d_1^h d_1^v \iota_b.$$

Therefore, if we assume that diag NNC is a Kan complex, we should find a diagonal bisimplex, say $x \in \text{NNC}_{2,2}$, such that $d_0^h d_0^v x = \iota_b$ and $d_2^h d_2^v x = \iota_a$. This $(2, 2)$ -simplex

x of the nerve of the double groupoid, should be therefore of the form

$$\begin{array}{ccc}
 \cdot & \xleftarrow{a} & \cdot \\
 \parallel & \wr & \parallel \\
 \cdot & \xleftarrow{a} & \cdot \\
 a' \uparrow & \wr & \uparrow b \\
 \cdot & \xleftarrow{b'} & \cdot
 \end{array}$$

what is clearly impossible. Hence, the diagonal of the bisimplicial nerve of (A, B) is not a Kan complex, in spite of the fact that this bisimplicial set is a pointwise Kan complex in both vertical and horizontal directions.

For a last comment, we shall point out a fact suggested by a referee: the property proved in Theorem 1 for diagonal fibrations is not true for diagonal trivial fibrations. For instance: Let EG be the universal cover of a non-trivial discrete group G , that is, the simplicial set with $EG_n = G^{n+1}$ and faces given by $d_i(x_0, \dots, x_n) = (x_0, \dots, x_{i-1}, x_{i+1}, \dots, x_n)$, $0 \leq i \leq n$, and let $X = EG \otimes EG$, the bisimplicial set defined by $X_{p,q} = EG_p \times EG_q$. Then, $\text{diag } X$ is contractible to a one point Kan complex and, however, each simplicial set $X_{*,q} = EG \times G^q$ is homotopy equivalent to the constant simplicial set G^q , and therefore it is not contractible. In conclusion, $X \rightarrow pt$ is a diagonal trivial fibration but it is not a pointwise trivial fibration.

References

- [1] A. K. BOUSFIELD AND D. M. KAN, *Homotopy limits, completions and localizations*, Lecture Notes in Math., 304, Springer, 1972.
- [2] R. BROWN AND C. B. SPENCER, *Double groupoids and crossed modules*, Cahiers Topologie Géom. Différentielle, XVII (1976), pp. 343–362.
- [3] P. GOERSS AND J. JARDINE, *Simplicial Homotopy Theory*, Progress in Mathematics 174, Birkhäuser, 1999.
- [4] G. M. KELLY AND R. S. STREET, *Review of the elements of 2-categories*, in Category Seminar (Proc. Sem., Sydney, 1972/1973), Lecture Notes in Math., 420, Springer, 1974, pp. 75–103.
- [5] S. MAC LANE, *Categories for the working mathematician*, Graduate Texts in Mathematics, Vol. 5, Springer-Verlag, 1971.

- [6] J. P. MAY, *Simplicial objects in Algebraic Topology*, Van Nostrand Mathematical Studies, No. 11, D. Van Nostrand Co., 1967.
- [7] I. MOERDIJK, *Bisimplicial sets and the group-completion theorem*, in Algebraic K-Theory: connections with geometry and topology, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 279, Kluwer Academic Publishers, 1989, pp. 225–240.

<http://www.emis.de/ZMATH/>
<http://www.ams.org/mathscinet>

This article may be accessed via WWW at <http://jhrrs.rmi.acnet.ge>

A.M. Cegarra
acegarra@ugr.es

Departamento de Álgebra
Facultad de Ciencias
Universidad de Granada
18071 Granada
Spain

J. Remedios
jremed@ull.es

Departamento de Matemática Fundamental
Universidad de La Laguna
38271 La Laguna
Spain