

Journal of Inequalities in Pure and Applied Mathematics



APPROXIMATION OF ENTIRE FUNCTIONS OF TWO COMPLEX VARIABLES IN BANACH SPACES

RAMESH GANTI AND G.S. SRIVASTAVA

Department of Mathematics
Indian Institute of Technology Roorkee
Roorkee - 247 667, India.

EMail: girssfma@iitr.ernet.in

volume 7, issue 2, article 51,
2006.

*Received 05 January, 2006;
accepted 27 January, 2006.*

Communicated by: S.P. Singh

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)



Abstract

In the present paper, we study the polynomial approximation of entire functions of two complex variables in Banach spaces. The characterizations of order and type of entire functions of two complex variables have been obtained in terms of the approximation errors.

2000 Mathematics Subject Classification: 30B10, 30D15.

Key words: Entire function, Order, type, Approximation, Error.

Contents

1	Introduction	3
2	Main Results	7
References		

Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 2 of 26

1. Introduction

Let $f(z_1, z_2) = \sum a_{m_1 m_2} z_1^{m_1} z_2^{m_2}$ be a function of the complex variables z_1 and z_2 , regular for $|z_t| \leq r_t$, $t = 1, 2$. If r_1 and r_2 can be taken arbitrarily large, then $f(z_1, z_2)$ represents an entire function of the complex variables z_1 and z_2 . Following Bose and Sharma [1], we define the maximum modulus of $f(z_1, z_2)$ as

$$M(r_1, r_2) = \max_{|z_t| \leq r_t} |f(z_1, z_2)|, \quad t = 1, 2.$$

The order ρ of the entire function $f(z_1, z_2)$ is defined as [1, p. 219]:

$$\limsup_{r_1, r_2 \rightarrow \infty} \frac{\log \log M(r_1, r_2)}{\log(r_1 r_2)} = \rho.$$

For $0 < \rho < \infty$, the type τ of an entire function $f(z_1, z_2)$ is defined as [1, p. 223]:


$$\limsup_{r_1, r_2 \rightarrow \infty} \frac{\log M(r_1, r_2)}{r_1^\rho + r_2^\rho} = \tau.$$

Bose and Sharma [1], obtained the following characterizations for order and type of entire functions of two complex variables.

Theorem 1.1. *The entire function $f(z_1, z_2) = \sum_{m_1, m_2=0}^{\infty} a_{m_1 m_2} z_1^{m_1} z_2^{m_2}$ is of finite order if and only if*

$$\mu = \limsup_{m_1, m_2 \rightarrow \infty} \frac{\log(m_1^{m_1} m_2^{m_2})}{\log(|a_{m_1 m_2}|^{-1})}$$

is finite and then the order ρ of $f(z_1, z_2)$ is equal to μ .



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 3 of 26](#)

Define

$$\alpha = \limsup_{m_1, m_2 \rightarrow \infty} \sqrt[m_1 + m_2]{m_1^{m_1} m_2^{m_2} |a_{m_1 m_2}|^\rho}.$$

We have

Theorem 1.2. If $0 < \alpha < \infty$, the function $f(z_1, z_2) = \sum_{m_1, m_2=0}^{\infty} a_{m_1 m_2} z_1^{m_1} z_2^{m_2}$ is an entire function of order ρ and type τ if and only if $\alpha = e\rho\tau$.

Let $H_q, q > 0$ denote the space of functions $f(z_1, z_2)$ analytic in the unit bi-disc $U = \{z_1, z_2 \in C : |z_1| < 1, |z_2| < 1\}$ such that

$$\|f\|_{H_q} = \lim_{r_1, r_2 \rightarrow 1^-} M_q(f; r_1, r_2) < \infty,$$

where

$$M_q(f; r_1, r_2) = \left\{ \frac{1}{4\pi^2} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} |f(r_1 e^{it_1}, r_2 e^{it_2})|^q dt_1 dt_2 \right\}^{\frac{1}{q}},$$

and let $H'_q, q > 0$ denote the space of functions $f(z_1, z_2)$ analytic in U and satisfying the condition

$$\|f\|_{H'_q} = \left\{ \frac{1}{\pi^2} \int_{|z_1|<1} \int_{|z_2|<1} |f(z_1, z_2)|^q dx_1 dy_1 dx_2 dy_2 \right\}^{\frac{1}{q}} < \infty.$$

Set

$$\|f\|_{H'_\infty} = \|f\|_{H_\infty} = \sup \{|f(z_1, z_2)| : z_1, z_2 \in U\}.$$

H_q and H'_q are Banach spaces for $q \geq 1$. In analogy with spaces of functions of one variable, we call H_q and H'_q the Hardy and Bergman spaces respectively.



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 4 of 26](#)

The function $f(z_1, z_2)$ analytic in U belongs to the space $\mathbf{B}(p, q, \kappa)$, where $0 < p < q \leq \infty$, and $0 < \kappa \leq \infty$, if

$$\|f\|_{p,q,\kappa} = \left\{ \int_0^1 \int_0^1 \{(1-r_1)(1-r_2)\}^{\kappa(1/p-1/q)-1} M_q^\kappa(f, r_1, r_2) dr_1 dr_2 \right\}^{\frac{1}{\kappa}} < \infty,$$

$0 < \kappa < \infty,$

$$\|f\|_{p,q,\infty} = \sup \{(1-r_1)(1-r_2)\}^{(1/p-1/q)-1} M_q(f, r_1, r_2) : 0 < r_1, r_2 < 1\} < \infty.$$

The space $\mathbf{B}(p, q, \kappa)$ is a Banach space for $p > 0$ and $q, \kappa \geq 1$, otherwise it is a Fréchet space. Further, we have

$$(1.1) \quad H_q \subset H'_q = \mathbf{B}\left(\frac{q}{2}, q, q\right), \quad 1 \leq q < \infty.$$

Let X be a Banach space and let $E_{m,n}(f, X)$ be the best approximation of a function $f(z_1, z_2) \in X$ by elements of the space P that consists of algebraic polynomials of degree $\leq m + n$ in two complex variables:

$$(1.2) \quad E_{m,n}(f, X) = \inf \{\|f - p\|_x; p \in P\}.$$

To the best of our knowledge, characterizations for the order and type of entire functions of two complex variables in Banach spaces have not been obtained so far. In this paper, we have made an attempt to bridge this gap.



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 5 of 26](#)

Notation: For reducing the length of expressions we use the following notations in the main results.

$$B^{1/\kappa} \left[(n+1)\kappa + 1; \kappa \left(\frac{1}{p} - \frac{1}{2} \right) \right] = B[n, p, 2, \kappa]$$

$$B^{1/\kappa} \left[(m+1)\kappa + 1; \kappa \left(\frac{1}{p} - \frac{1}{2} \right) \right] = B[m, p, 2, \kappa]$$

$$B^{1/\kappa} \left[(n+1)\kappa + 1; \kappa \left(\frac{1}{p} - \frac{1}{q} \right) \right] = B[n, p, q, \kappa]$$

$$B^{1/\kappa} \left[(m+1)\kappa + 1; \kappa \left(\frac{1}{p} - \frac{1}{q} \right) \right] = B[m, p, q, \kappa].$$



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 6 of 26

2. Main Results

Theorem 2.1. Let $f(z_1, z_2) = \sum_{m,n=0}^{\infty} a_{mn} z_1^m z_2^n$, then the entire function $f(z_1, z_2)$ $\in \mathbf{B}(p, q, \kappa)$ is of finite order ρ , if and only if

$$(2.1) \quad \rho = \limsup_{m,n \rightarrow \infty} \frac{\ln(m^m n^n)}{-\ln E_{m,n}(f, \mathbf{B}(p, q, \kappa))}.$$

Proof. We prove the above result in two steps. First we consider the space $\mathbf{B}(p, q, \kappa)$, $q = 2$, $0 < p < 2$ and $\kappa \geq 1$. Let $f(z_1, z_2) \in \mathbf{B}(p, q, \kappa)$ be of order ρ . From Theorem 1.1, for any $\epsilon > 0$, there exists a natural number $n_0 = n_0(\epsilon)$ such that

$$(2.2) \quad |a_{mn}| \leq m^{-m/\rho+\epsilon} n^{-n/\rho+\epsilon} \quad m, n > n_0.$$

We denote the partial sum of the Taylor series of a function $f(z_1, z_2)$ by

$$T_{m,n}(f, z_1, z_2) = \sum_{j_1=0}^m \sum_{j_2=0}^n a_{j_1 j_2} z_1^{j_1} z_2^{j_2}.$$

We write

$$(2.3) \quad \begin{aligned} E_{m,n}(f, \mathbf{B}(p, 2, \kappa)) &= \|f - T_{m,n}(f)\|_{p,2,\kappa} \\ &= \left\{ \int_0^1 \int_0^1 \{(1-r_1)(1-r_2)\}^{\kappa(1/p-1/2)-1} \right. \\ &\quad \times \left. \left(\sum_{j_1} \sum_{j_2} r_1^{2j_1} r_2^{2j_2} |a_{j_1 j_2}|^2 \right)^{\frac{\kappa}{2}} dr_1 dr_2 \right\}^{\frac{1}{\kappa}}, \end{aligned}$$



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 7 of 26](#)

where

$$\sum_{j_1} \sum_{j_2} r_1^{2j_1} r_2^{2j_2} |a_{j_1 j_2}|^2 = S_1 + S_2 + \sum_{j_1=m+1}^{\infty} \sum_{j_2=n+1}^{\infty} r_1^{2j_1} r_2^{2j_2} |a_{j_1 j_2}|^2,$$

$$S_1 = \sum_{j_1=0}^m \sum_{j_2=n+1}^{\infty} r_1^{2j_1} r_2^{2j_2} |a_{j_1 j_2}|^2 \quad \text{and} \quad S_2 = \sum_{j_1=m+1}^{\infty} \sum_{j_2=0}^n r_1^{2j_1} r_2^{2j_2} |a_{j_1 j_2}|^2.$$

Since S_1, S_2 are bounded and $r_1, r_2 < 1$, therefore the above expression (2.3) becomes

$$E_{m,n}(f, \mathbf{B}(p, 2, \kappa)) \leq C \left\{ \int_0^1 \{(1-r)^{\kappa(1/p-1/2)-1}\} r^{(s+1)\kappa} dr \right\} \left\{ \sum_{j_1=m+1}^{\infty} \sum_{j_2=n+1}^{\infty} |a_{j_1 j_2}|^2 \right\}^{\frac{1}{2}},$$

where

$$\begin{aligned} & \left\{ \int_0^1 \{(1-r)^{\kappa(1/p-1/2)-1}\} r^{(s+1)\kappa} dr \right\} \\ &= \left\{ \int_0^1 \{(1-r_1)^{\kappa(1/p-1/2)-1}\} r_1^{(m+1)\kappa} dr_1 \right\} \\ & \quad \times \left\{ \int_0^1 \{(1-r_2)\}^{\kappa(1/p-1/2)-1} r_2^{(n+1)\kappa} dr_2 \right\}. \end{aligned}$$



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 8 of 26](#)

Therefore

$$(2.4) \quad E_{m,n}(f, \mathbf{B}(p, 2, \kappa))$$

$$\leq CB[m, p, 2, \kappa]B[n, p, 2, \kappa] \left\{ \sum_{j_1=m+1}^{\infty} \sum_{j_2=n+1}^{\infty} |a_{j_1 j_2}|^2 \right\}^{\frac{1}{2}},$$

where C is a constant and $B(a, b)$ ($a, b > 0$) denotes the beta function.

By using (2.2), we have

$$\begin{aligned} \sum_{j_1=m+1}^{\infty} \sum_{j_2=n+1}^{\infty} |a_{j_1 j_2}|^2 &\leq \sum_{j_1=m+1}^{\infty} \sum_{j_2=n+1}^{\infty} j_1^{-\frac{2j_1}{\rho+\epsilon}} j_2^{-\frac{2j_2}{\rho+\epsilon}} \\ &\leq \sum_{j_1=m+1}^{\infty} j_1^{-\frac{2j_1}{\rho+\epsilon}} \sum_{j_2=n+1}^{\infty} j_2^{-\frac{2j_2}{\rho+\epsilon}} \\ &\leq O(1)(m+1)^{-2(m+1)/\rho+\epsilon}(n+1)^{-2(n+1)/\rho+\epsilon}. \end{aligned}$$

Using the above inequality in (2.4), we have

$$\begin{aligned} E_{m,n}(f, \mathbf{B}(p, 2, \kappa)) &\leq CB[m, p, 2, \kappa]B[n, p, 2, \kappa](m+1)^{-(m+1)/\rho+\epsilon}(n+1)^{-(n+1)/\rho+\epsilon}. \\ \Rightarrow \rho+\epsilon &\geq \frac{\ln [(m+1)^{(m+1)}(n+1)^{(n+1)}]}{-\ln \{E_{m,n}(f, \mathbf{B}(p, 2, \kappa))\} + \ln \{B[m, p, 2, \kappa]\} + \ln \{B[n, p, 2, \kappa]\}}. \end{aligned}$$

Now

$$B\left[(n+1)\kappa + 1; \kappa \left(\frac{1}{p} - \frac{1}{2}\right)\right] = \frac{\Gamma((n+1)\kappa + 1)\Gamma\left(\kappa \left(\frac{1}{p} - \frac{1}{2}\right)\right)}{\Gamma\left(\left(n + \frac{1}{2} + \frac{1}{p}\right)\kappa + 1\right)}.$$



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 9 of 26](#)

Hence

$$B \left[(n+1)\kappa + 1; \kappa \left(\frac{1}{p} - \frac{1}{2} \right) \right] \\ \simeq \frac{e^{-[(n+1)\kappa+1]} [(n+1)\kappa + 1]^{(n+1)\kappa+3/2} \Gamma \left(\frac{1}{p} - \frac{1}{2} \right)}{e^{[(n+1/2+1/p)\kappa+1]} [(n+\frac{1}{2}+\frac{1}{p})\kappa + 1]^{(n+1/2+1/p)\kappa+3/2}}.$$

Thus

$$(2.5) \quad \left\{ B \left[(n+1)\kappa + 1; \kappa \left(\frac{1}{p} - \frac{1}{2} \right) \right] \right\}^{\frac{1}{(n+1)}} \simeq 1.$$

Now proceeding to limits, we obtain


$$(2.6) \quad \rho \geq \limsup_{m,n \rightarrow \infty} \frac{\ln(m^m n^n)}{-\ln \{E_{m,n}(f, \mathbf{B}(p, 2, \kappa))\}}.$$

For the reverse inequality, since from the right hand side of the inequality (2.4), we have

$$(2.7) \quad |a_{m+1n+1}| B[m, p, 2, \kappa] B[n, p, 2, \kappa] \leq E_{m,n}(f, \mathbf{B}(p, 2, \kappa)),$$

we have

$$\frac{\ln(m^m n^n)}{-\ln E_{m,n}(f, \mathbf{B}(p, 2, \kappa))} \\ \geq \frac{\ln(m^m n^n)}{-\ln |a_{m+1n+1}| + \ln \{B[m, p, 2, \kappa]\} + \ln \{B[n, p, 2, \kappa]\}}.$$



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 10 of 26

Now proceeding to limits, we obtain

$$(2.8) \quad \limsup_{m,n \rightarrow \infty} \frac{\ln(m^m n^n)}{-\ln E_{m,n}(f, \mathbf{B}(p, 2, \kappa))} \geq \rho.$$

From (2.6) and (2.8), we get the required result.

In the second step, for the general case $\mathbf{B}(p, q, \kappa)$, $q \neq 2$, we have

$$(2.9) \quad \begin{aligned} E_{m,n}(f, \mathbf{B}(p, q, \kappa)) &\leq \|f - T_{m,n}(f)\|_{p,q,\kappa} \\ &= \left\{ \int_0^1 \int_0^1 \{(1-r_1)(1-r_2)\}^{\kappa(1/p-1/q)-1} \right. \\ &\quad \times \left(\sum_{j_1} \sum_{j_2} r_1^{qj_1} r_2^{qj_2} |a_{j_1 j_2}|^q \right)^{\frac{1}{q}} dr_1 dr_2 \right\}^{\frac{1}{\kappa}}, \end{aligned}$$

where

$$\sum_{j_1} \sum_{j_2} r_1^{2j_1} r_2^{2j_2} |a_{j_1 j_2}|^2 = S_1 + S_2 + \sum_{j_1=m+1}^{\infty} \sum_{j_2=n+1}^{\infty} r_1^{2j_1} r_2^{2j_2} |a_{j_1 j_2}|^2,$$

$$S_1 = \sum_{j_1=0}^m \sum_{j_2=n+1}^{\infty} r_1^{2j_1} r_2^{2j_2} |a_{j_1 j_2}|^2 \quad \text{and} \quad S_2 = \sum_{j_1=m+1}^{\infty} \sum_{j_2=0}^n r_1^{2j_1} r_2^{2j_2} |a_{j_1 j_2}|^2.$$

Since S_1, S_2 are bounded and $r_1, r_2 < 1$, therefore the above expression (2.9)



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 1 of 26](#)

becomes

$$E_{m,n}(f, \mathbf{B}(p, q, \kappa)) \leq C' \left\{ \int_0^1 \{(1-r)^{\kappa(1/p-1/q)-1}\} r^{(s+1)\kappa} dr \right\} \left\{ \sum_{j_1=m+1}^{\infty} \sum_{j_2=n+1}^{\infty} |a_{j_1 j_2}|^q \right\}^{\frac{1}{q}},$$


where

$$\begin{aligned} & \left\{ \int_0^1 \{(1-r)^{\kappa(1/p-1/q)-1}\} r^{(s+1)\kappa} dr \right\} \\ &= \left\{ \int_0^1 \{(1-r_1)^{\kappa(1/p-1/q)-1}\} r_1^{(m+1)\kappa} dr_1 \right\} \\ &\quad \times \left\{ \int_0^1 \{(1-r_2)\}^{\kappa(1/p-1/q)-1} r_2^{(n+1)\kappa} dr_2 \right\}. \end{aligned}$$

Therefore

$$(2.10) \quad E_{m,n}(f, \mathbf{B}(p, q, \kappa)) \leq C' B[m, p, q, \kappa] B[n, p, q, \kappa] \left\{ \sum_{j_1=m+1}^{\infty} \sum_{j_2=n+1}^{\infty} |a_{j_1 j_2}|^q \right\}^{\frac{1}{q}},$$

where C' is constant and $B[m, p, q, \kappa]$ is Euler's integral of the first kind. By



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 12 of 26](#)

using (2.2), we get

$$\begin{aligned} \sum_{j_1=m+1}^{\infty} \sum_{j_2=n+1}^{\infty} |a_{j_1 j_2}|^q &\leq \sum_{j_1=m+1}^{\infty} j_1^{\frac{-q j_1}{(\rho+\epsilon)}} \sum_{j_2=n+1}^{\infty} j_2^{\frac{-q j_2}{(\rho+\epsilon)}} \\ &\leq O(1)(m+1)^{\frac{-q(m+1)}{(\rho+\epsilon)}}(n+1)^{\frac{-q(n+1)}{(\rho+\epsilon)}}. \end{aligned}$$

Using above inequality in (2.10), we get

$$\begin{aligned} E_{m,n}(f, \mathbf{B}(p, q, \kappa)) \\ \leq C' B[m, p, q, \kappa] B[n, p, q, \kappa] (m+1)^{-(m+1)/\rho+\epsilon} (n+1)^{-(n+1)/\rho+\epsilon}. \\ \Rightarrow \rho+\epsilon \geq \frac{\ln [(m+1)^{m+1} (n+1)^{n+1}]}{-\ln E_{m,n}(f, \mathbf{B}(p, q, \kappa)) + \ln \{B[m, p, q, \kappa]\} + \ln \{B[n, p, q, \kappa]\}}. \end{aligned}$$


Now proceeding to limits, we obtain

$$(2.11) \quad \rho \geq \limsup_{m,n \rightarrow \infty} \frac{\ln (m^m n^n)}{-\ln E_{m,n}(f, \mathbf{B}(p, q, \kappa))}.$$

Let $0 < p < q < 2$, and $\kappa, q \geq 1$. Since

$$E_{m,n}(f, \mathbf{B}(p_1, q_1, \kappa_1)) \leq 2^{1/q-1/q_1} \left[\kappa \left(\frac{1}{p} - \frac{1}{q} \right) \right]^{\frac{1}{\kappa} - \frac{1}{\kappa_1}} E_{m,n}(f, \mathbf{B}(p, q, \kappa)),$$

where $p_1 = p, q_1 = 2$ and $\kappa_1 = \kappa$, and the condition (2.1) is already proved for



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 13 of 26](#)

the space $\mathbf{B}(p, 2, \kappa)$, we get

$$(2.12) \quad \limsup_{m,n \rightarrow \infty} \frac{\ln(m^m n^n)}{-\ln E_{m,n}(f, \mathbf{B}(p, q, \kappa))} \\ \geq \limsup_{m,n \rightarrow \infty} \frac{\ln(m^m n^n)}{-\ln E_{m,n}(f, \mathbf{B}(p, 2, \kappa))} = \rho.$$

Now let $0 < p \leq 2 < q$. Since

$$M_2(f, r_1, r_2) \leq M_q(f, r_1, r_2), \quad 0 < r_1, r_2 < 1,$$

therefore

$$(2.13) \quad E_{m,n}(f, \mathbf{B}(p, q, \kappa)) \\ \geq \left\{ \int_0^1 \int_0^1 \{(1-r_1)(1-r_2)\}^{\kappa(1/p-1/q)-1} Q dr_1 dr_2 \right\}^{\frac{1}{\kappa}} \\ \geq |a_{m+1n+1}| B[m, p, q, \kappa] B[n, p, q, \kappa],$$

where $Q = \inf [M_2^\kappa(f - p; r_1, r_2) : p \in P]$. Hence we have

$$\frac{\ln(m^m n^n)}{-\ln E_{m,n}(f, \mathbf{B}(p, q, \kappa))} \\ \geq \frac{\ln(m^m n^n)}{-\ln |a_{m+1n+1}| + \ln \{B[m, p, q, \kappa]\} + \ln \{B[n, p, q, \kappa]\}}.$$

Now proceeding to limits, we obtain

$$(2.14) \quad \limsup_{m,n \rightarrow \infty} \frac{\ln(m^m n^n)}{-\ln E_{m,n}(f, \mathbf{B}(p, q, \kappa))} \geq \rho.$$



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 14 of 26

From (2.11) and (2.14), we get the required result. \square

Now we prove

Theorem 2.2. *Let*

$$f(z_1, z_2) = \sum_{m,n=0}^{\infty} a_{mn} z_1^m z_2^n,$$

then the entire function $f(z_1, z_2) \in H_q$ is of finite order ρ , if and only if

$$(2.15) \quad \rho = \limsup_{m,n \rightarrow \infty} \frac{\ln(m^m n^n)}{-\ln E_{m,n}(f, H_q)}.$$

Proof. Let $f(z_1, z_2) = \sum_{m,n=0}^{\infty} a_{mn} z_1^m z_2^n \in H_q$ be an entire transcendental function. Since f is entire, we have

$$(2.16) \quad \lim_{m,n \rightarrow \infty} \sqrt[m+n]{|a_{mn}|} = 0,$$

and $f \in H_q$, therefore


$$M_q(f; r_1, r_2) < \infty,$$

and $f(z_1, z_2) \in \mathbf{B}(p, q, \kappa)$, $0 < p < q \leq \infty$; $q, \kappa \geq 1$. By (1.1) we obtain

$$(2.17) \quad E_{m,n}(f, \mathbf{B}(q/2, q, q)) \leq \varsigma_q E_{m,n}(f, H_q), \quad 1 \leq q < \infty,$$

where ς_q is a constant independent of m, n and f . In the case of space H_∞ ,

$$(2.18) \quad E_{m,n}(f, \mathbf{B}(p, \infty, \infty)) \leq E_{m,n}(f, H_\infty), \quad 0 < p < \infty.$$



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 15 of 26](#)

From (2.17), we have

$$\begin{aligned}
 (2.19) \quad \xi(f) &= \limsup_{m,n \rightarrow \infty} \frac{\ln(m^m n^n)}{-\ln E_{m,n}(f, H_q)} \\
 &\geq \limsup_{m,n \rightarrow \infty} \frac{\ln(m^m n^n)}{-\ln E_{m,n}(f, \mathbf{B}(q/2, q, q))} \\
 &\geq \rho, \quad 1 \leq q < \infty,
 \end{aligned}$$

and using estimate (2.18) we prove inequality (2.19) for the case $q = \infty$.

For the reverse inequality


$$(2.20) \quad \xi(f) \leq \rho,$$

since

$$E_{m,n}(f, H_q) \leq O(1) \sum_{j_1=m+1}^{\infty} \sum_{j_2=n+1}^{\infty} |a_{j_1 j_2}(f)|,$$

using (2.2), we have

$$\begin{aligned}
 E_{m,n}(f, H_q) &\leq O(1) \sum_{j_1=m+1}^{\infty} \sum_{j_2=n+1}^{\infty} j_1^{-\frac{j_1}{\rho+\epsilon}} j_2^{-\frac{j_2}{\rho+\epsilon}} \\
 &\leq O(1) \sum_{j_1=m+1}^{\infty} j_1^{-\frac{j_1}{\rho+\epsilon}} \sum_{j_2=n+1}^{\infty} j_2^{-\frac{j_2}{\rho+\epsilon}} \\
 &\leq O(1)(m+1)^{-(m+1)/\rho+\epsilon} (n+1)^{-(n+1)/\rho+\epsilon}.
 \end{aligned}$$



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 16 of 26

$$\Rightarrow \rho + \epsilon \geq \frac{\ln [(m+1)^{(m+1)}(n+1)^{(n+1)}]}{-\ln [E_{m,n}(f, H_q)]}.$$

Now proceeding to limits and since ϵ is arbitrary, then we will get (2.20). From (2.19) and (2.20) we will obtain the required result.

Now we prove sufficiency. Assume that the condition (2.15) is satisfied. Then it follows that $\ln [1/E_{m,n}(f, H_q)]^{1/(m+n)} \rightarrow \infty$ as $m, n \rightarrow \infty$.

This yields

$$\lim_{m,n \rightarrow \infty} \sqrt[m+n]{E_{m,n}(f, H_q)} = 0.$$

This relation and the estimate $|a_{m+1,n+1}(f)| \leq E_{m,n}(f, H_q)$ yield the relation (2.16). This means that $f(z_1, z_2) \in H_q$ is an entire transcendental function. \square

Now we prove


Theorem 2.3. Let $f(z_1, z_2) = \sum_{m,n=0}^{\infty} a_{mn} z_1^m z_2^n$, then the entire function $f(z_1, z_2) \in \mathbf{B}(p, q, \kappa)$ of finite order ρ , is of type τ if and only if

$$(2.21) \quad \tau = \frac{1}{e\rho} \limsup_{m,n \rightarrow \infty} \{m^m n^n E_{m,n}^{\rho}(f, \mathbf{B}(p, q, \kappa))\}^{\frac{1}{m+n}}.$$

Proof. We prove the above result in two steps.

First we consider the space $\mathbf{B}(p, q, \kappa)$, $q = 2$, $0 < p < 2$ and $\kappa \geq 1$. Let $f(z) \in \mathbf{B}(p, q, \kappa)$ be of order ρ . From Theorem 1.2, for any $\epsilon > 0$, there exists a natural number $n_0 = n_0(\epsilon)$ such that

$$(2.22) \quad |a_{mn}| \leq m^{-m/\rho} n^{-n/\rho} [e\rho(\tau + \epsilon)]^{\frac{m+n}{\rho}}.$$



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 17 of 26](#)

We denote the partial sum of the Taylor series of a function $f(z_1, z_2)$ by

$$T_{m,n}(f, z_1, z_2) = \sum_{j_1=0}^m \sum_{j_2=0}^n a_{j_1 j_2} z_1^{j_1} z_2^{j_2},$$

we write


$$\begin{aligned} (2.23) \quad E_{m,n}(f, \mathbf{B}(p, 2, \kappa)) &= \|f - T_{m,n}(f)\|_{p,2,\kappa} \\ &= \left\{ \int_0^1 \int_0^1 \{(1-r_1)(1-r_2)\}^{\kappa(1/p-1/2)-1} \right. \\ &\quad \times \left. \left(\sum_{j_1} \sum_{j_2} r_1^{2j_1} r_2^{2j_2} |a_{j_1 j_2}|^2 \right)^{\frac{\kappa}{2}} dr_1 dr_2 \right\}^{\frac{1}{\kappa}}, \end{aligned}$$

where

$$\sum_{j_1} \sum_{j_2} r_1^{2j_1} r_2^{2j_2} |a_{j_1 j_2}|^2 = S_1 + S_2 + \sum_{j_1=m+1}^{\infty} \sum_{j_2=n+1}^{\infty} r_1^{2j_1} r_2^{2j_2} |a_{j_1 j_2}|^2,$$

$$S_1 = \sum_{j_1=0}^m \sum_{j_2=n+1}^{\infty} r_1^{2j_1} r_2^{2j_2} |a_{j_1 j_2}|^2 \quad \text{and} \quad S_2 = \sum_{j_1=m+1}^{\infty} \sum_{j_2=0}^n r_1^{2j_1} r_2^{2j_2} |a_{j_1 j_2}|^2.$$

Since S_1, S_2 are bounded, and $r_1, r_2 < 1$ therefore the above expression (2.23)



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 18 of 26](#)

becomes

$$(2.24) \quad E_{m,n}(f, \mathbf{B}(p, 2, \kappa))$$

$$\leq DB[m, p, 2, \kappa]B[n, p, 2, \kappa] \left\{ \sum_{j_1=m+1}^{\infty} \sum_{j_2=n+1}^{\infty} |a_{j_1 j_2}|^2 \right\}^{\frac{1}{2}},$$

where D is a constant and $B(a, b)$ ($a, b > 0$) denotes the beta function. By using (2.22), we have

$$\begin{aligned} & \sum_{j_1=m+1}^{\infty} \sum_{j_2=n+1}^{\infty} |a_{j_1 j_2}|^2 \\ & \leq \sum_{j_1=m+1}^{\infty} \sum_{j_2=n+1}^{\infty} j_1^{-\frac{2j_1}{\rho}} j_2^{-\frac{2j_2}{\rho}} [e\rho(\tau + \epsilon)]^{\frac{2(j_1+j_2)}{\rho}} \\ & \leq \sum_{j_1=m+1}^{\infty} j_1^{-\frac{2j_1}{\rho}} [e\rho(\tau + \epsilon)]^{\frac{2j_1}{\rho}} \sum_{j_2=n+1}^{\infty} j_2^{-\frac{2j_2}{\rho+\epsilon}} [e\rho(\tau + \epsilon)]^{\frac{2j_2}{\rho}} \\ & \leq O(1)(m+1)^{-2(m+1)/\rho}(n+1)^{-2(n+1)/\rho}[e\rho(\tau + \epsilon)]^{\frac{2(m+n+2)}{\rho}}. \end{aligned}$$


Using the above inequality in (2.24), we get

$$E_{m,n}^{\rho}(f, \mathbf{B}(p, 2, \kappa)) \leq D^{\rho} B^{\rho}[m, p, 2, \kappa] B^{\rho}[n, p, 2, \kappa] Y [e\rho(\tau + \epsilon)]^{(m+n+2)},$$

where $Y = (m+1)^{-(m+1)}(n+1)^{-(n+1)}$.

Now proceeding to limits and since ϵ is arbitrary, we have

$$(2.25) \quad \frac{1}{e\rho} \limsup_{m,n \rightarrow \infty} \{m^m n^n E_{m,n}^{\rho}(f, \mathbf{B}(p, 2, \kappa))\}^{\frac{1}{m+n}} \leq \tau.$$



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 19 of 26

For the reverse inequality, since from the right hand side of (2.24),

$$|a_{m+1n+1}|B[m, p, 2, \kappa]B[n, p, 2, \kappa] \leq E_{m,n}(f, \mathbf{B}(p, 2, \kappa))$$

we have

$$\begin{aligned} m^{m/(m+n)} n^{n/(m+n)} |a_{m+1n+1}|^{\rho/(m+n)} B^{\frac{\rho}{(m+n)}} [m, p, 2, \kappa] B^{\frac{\rho}{(m+n)}} [n, p, 2, \kappa] \\ \leq \{E_{m,n}^\rho m^m n^n\}^{1/(m+n)}. \end{aligned}$$

Now proceeding to limits, we obtain

$$(2.26) \quad \tau \leq \frac{1}{e\rho} \limsup_{m,n \rightarrow \infty} \{m^m n^n E_{m,n}^\rho(f, \mathbf{B}(p, 2, \kappa))\}^{\frac{1}{m+n}}.$$

From (2.25) and (2.26), we get the required result.

In the second step, for the general case $\mathbf{B}(p, q, \kappa)$, $q \neq 2$, we have

$$\begin{aligned} (2.27) \quad E_{m,n}(f, \mathbf{B}(p, q, \kappa)) &\leq \|f - T_{m,n}(f)\|_{p,q,\kappa} \\ &= \left\{ \int_0^1 \int_0^1 \{(1-r_1)(1-r_2)\}^{\kappa(1/p-1/q)-1} \right. \\ &\quad \times \left. \left(\sum_{j_1} \sum_{j_2} r_1^{qj_1} r_2^{qj_2} |a_{j_1 j_2}|^q \right)^{\frac{\kappa}{q}} dr_1 dr_2 \right\}^{\frac{1}{\kappa}}, \end{aligned}$$

where

$$\sum_{j_1} \sum_{j_2} r_1^{2j_1} r_2^{2j_2} |a_{j_1 j_2}|^2 = S_1 + S_2 + \sum_{j_1=m+1}^{\infty} \sum_{j_2=n+1}^{\infty} r_1^{2j_1} r_2^{2j_2} |a_{j_1 j_2}|^2,$$



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 20 of 26

$$S_1 = \sum_{j_1=0}^m \sum_{j_2=n+1}^{\infty} r_1^{2j_1} r_2^{2j_2} |a_{j_1 j_2}|^2 \quad \text{and} \quad S_2 = \sum_{j_1=m+1}^{\infty} \sum_{j_2=0}^n r_1^{2j_1} r_2^{2j_2} |a_{j_1 j_2}|^2.$$

Since S_1, S_2 are bounded, therefore the above expression (2.27) becomes


$$\begin{aligned} E_{m,n}(f, \mathbf{B}(p, q, \kappa)) \\ \leq G \left\{ \int_0^1 \{(1-r)^{\kappa(1/p-1/q)-1}\} r^{(s+1)\kappa} dr \right\} \left\{ \sum_{j_1=m+1}^{\infty} \sum_{j_2=n+1}^{\infty} |a_{j_1 j_2}|^q \right\}^{\frac{1}{q}}, \end{aligned}$$

where

$$\begin{aligned} & \left\{ \int_0^1 \{(1-r)^{\kappa(1/p-1/q)-1}\} r^{(s+1)\kappa} dr \right\} \\ &= \left\{ \int_0^1 \{(1-r_1)^{\kappa(1/p-1/q)-1}\} r_1^{(m+1)\kappa} dr_1 \right\} \\ & \quad \times \left\{ \int_0^1 \{(1-r_2)\}^{\kappa(1/p-1/q)-1} r_2^{(n+1)\kappa} dr_2 \right\}. \end{aligned}$$

Since $r_1, r_2 < 1$, therefore we have

$$(2.28) \quad \begin{aligned} E_{m,n}(f, \mathbf{B}(p, q, \kappa)) \\ \leq GB[m, p, q, \kappa] B[n, p, q, \kappa] \left\{ \sum_{j_1=m+1}^{\infty} \sum_{j_2=n+1}^{\infty} |a_{j_1 j_2}|^q \right\}^{\frac{1}{q}}, \end{aligned}$$



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 21 of 26](#)

where G is constant and $B[m, p, q, \kappa]$ is Euler's integral of the first kind. Using (2.22), we get

$$\begin{aligned} & \sum_{j_1=m+1}^{\infty} \sum_{j_2=n+1}^{\infty} |a_{j_1 j_2}|^q \\ & \leq \sum_{j_1=m+1}^{\infty} \sum_{j_2=n+1}^{\infty} j_1^{-\frac{q j_1}{\rho}} j_2^{-\frac{q j_2}{\rho}} [e\rho(\tau + \epsilon)]^{\frac{q(j_1+j_2)}{\rho}} \\ & \leq \sum_{j_1=m+1}^{\infty} j_1^{-\frac{q j_1}{\rho}} [e\rho(\tau + \epsilon)]^{\frac{q j_1}{\rho}} \sum_{j_2=n+1}^{\infty} j_2^{-\frac{q j_2}{\rho+\epsilon}} [e\rho(\tau + \epsilon)]^{\frac{q j_2}{\rho}} \\ & \leq O(1)(m+1)^{-q(m+1)/\rho}(n+1)^{-q(n+1)/\rho}[e\rho(\tau + \epsilon)]^{\frac{q(m+n+2)}{\rho}}. \end{aligned}$$

Using the above inequality in (2.28), we get


$$E_{m,n}^{\rho}(f, \mathbf{B}(p, q, \kappa)) \leq G^{\rho} B^{\rho}[m, p, q, \kappa] B^{\rho}[n, p, q, \kappa] Y [e\rho(\tau + \epsilon)]^{(m+n+2)},$$

where $Y = (m+1)^{-(m+1)}(n+1)^{-(n+1)}$. Now proceeding to limits, since ϵ is arbitrary, we have

$$(2.29) \quad \frac{1}{e\rho} \limsup_{m,n \rightarrow \infty} \{m^m n^n E_{mn}^{\rho}(f, \mathbf{B}(p, q, \kappa))\}^{\frac{1}{m+n}} \leq \tau.$$

Let $0 < p < q < 2$, and $\kappa, q \geq 1$. Since

$$E_{m,n}(f, \mathbf{B}(p_1, q_1, \kappa_1)) \leq 2^{1/q-1/q_1} [\kappa(1/p - 1/q)]^{1/\kappa-1/\kappa_1} E_{m,n}(f, \mathbf{B}(p, q, \kappa)),$$



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 22 of 26](#)

where $p_1 = p$, $q_1 = 2$ and $\kappa_1 = \kappa$, and the condition (2.21) has already been proved for the space $\mathbf{B}(p, 2, \kappa)$, we get

$$\begin{aligned} & \limsup_{m,n \rightarrow \infty} \{m^m n^n E_{m,n}^\rho(f, \mathbf{B}(p, q, \kappa))\}^{\frac{1}{m+n}} \\ & \geq \limsup_{m,n \rightarrow \infty} \{m^m n^n E_{m,n}^\rho(f, \mathbf{B}(p, 2, \kappa))\}^{\frac{1}{m+n}} = \tau. \end{aligned}$$

Now let $0 < p \leq 2 < q$. Since, in this case we have

$$M_2(f, r_1, r_2) \leq M_q(f, r_1, r_2), \quad 0 < r_1, r_2 < 1,$$

therefore


$$\begin{aligned} (2.30) \quad & \limsup_{m,n \rightarrow \infty} \{m^m n^n E_{m,n}^\rho(f, \mathbf{B}(p, q, \kappa))\}^{\frac{1}{m+n}} \\ & \geq \limsup_{m,n \rightarrow \infty} \{m^m n^n |a_{mn}|^\rho\}^{\frac{1}{m+n}} \\ & = e\rho\tau. \end{aligned}$$

From (2.29) and (2.30), we get the required result. \square

Lastly we prove

Theorem 2.4. Let $f(z_1, z_2) = \sum_{m,n=0}^{\infty} a_{mn} z_1^m z_2^n$, then the entire function $f(z_1, z_2) \in H_q$ having finite order ρ is of type τ if and only if

$$(2.31) \quad \tau = \frac{1}{e\rho} \limsup_{m,n \rightarrow \infty} \{m^m n^n E_{m,n}^\rho(f, H_q)\}^{\frac{1}{m+n}}.$$



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 23 of 26](#)

Proof. Since $f(z_1, z_2) = \sum_{m,n=0}^{\infty} a_{mn} z_1^m z_2^n$ is an entire transcendental function, we have

$$(2.32) \quad \lim_{m,n \rightarrow \infty} \sqrt[m+n]{|a_{mn}|} = 0.$$

Therefore $f(z_1, z_2) \in \mathbf{B}(p, q, \kappa)$, $0 < p < q \leq \infty$; $q, \kappa \geq 1$. We have

$$(2.33) \quad \begin{aligned} \xi(f) &= \frac{1}{e\rho} \limsup_{m,n \rightarrow \infty} \{m^m n^n E_{m,n}^\rho(f, H_q)\}^{\frac{1}{m+n}} \\ &\geq \frac{1}{e\rho} \limsup_{m,n \rightarrow \infty} \left\{ m^m n^n E_{m,n}^\rho \left(f, \mathbf{B} \left(\frac{q}{2}, q, q \right) \right) \right\}^{\frac{1}{m+n}} = \tau \end{aligned}$$

for $1 \leq q < \infty$. Using the estimate (2.18) we prove inequality (2.33) in the case $q = \infty$. For the reverse inequality

$$(2.34) \quad \xi(f) \leq \tau,$$

we have

$$E_{m,n}(f, H_q) \leq \sum_{j_1=m+1}^{\infty} \sum_{j_2=n+1}^{\infty} |a_{j_1 j_2}(f)|.$$

Using (2.22), we get

$$\begin{aligned} E_{m,n}^\rho(f, H_q) &\leq O(1)(m+1)^{-(m+1)}(n+1)^{-(n+1)}[e\rho(\tau + \epsilon)]^{(m+n+2)} \\ &\Rightarrow \tau + \epsilon \geq \frac{1}{e\rho} \{(m+1)^{(m+1)}(n+1)^{(n+1)} E_{m,n}^\rho(f, H_q)\}^{\frac{1}{(m+n+2)}}. \end{aligned}$$



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 24 of 26

Now proceeding to limits, since ϵ is arbitrary, we get

$$(2.35) \quad \tau \geq \frac{1}{e\rho} \limsup_{m,n \rightarrow \infty} \{m^m n^n E_{m,n}^\rho(f, H_q)\}^{\frac{1}{m+n}}.$$


From (2.33) and (2.35), we obtain the required result.

Now we prove sufficiency. Assume that the condition (2.31) is satisfied. Then it follows that $\{E_{m,n}^\rho(f, H_q)\}^{1/(m+n)} \rightarrow 0$ as $m, n \rightarrow \infty$. This yields

$$\lim_{m,n \rightarrow \infty} \sqrt[m+n]{E_{m,n}(f, H_q)} = 0.$$

This relation and the estimate $|a_{m+1n+1}(f)| \leq E_{m,n}(f, H_q)$ yield the inequality (2.32). This implies that $f(z_1, z_2) \in H_q$ is an entire transcendental function.

□



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

[Page 25 of 26](#)

References

- [1] S.K. BOSE, AND D. SHARMA, Integral functions of two complex variables, *Compositio Math.*, **15** (1963), 210–226.
- [2] S.B. VAKARCHUK AND S.I. ZHIR, On some problems of polynomial approximation of entire transcendental functions, *Ukrainian Mathem. J.*, **54**(9) (2002), 1393–1401.



Approximation of Entire Functions of Two Complex Variables in Banach Spaces

Ramesh Ganti and G.S. Srivastava

[Title Page](#)

[Contents](#)



[Go Back](#)

[Close](#)

[Quit](#)

Page 26 of 26