

APPROXIMATING THE FINITE HILBERT TRANSFORM VIA AN OSTROWSKI TYPE INEQUALITY FOR FUNCTIONS OF BOUNDED VARIATION

S.S. DRAGOMIR

School of Communications and Informatics
Victoria University of Technology
PO Box 14428
Melbourne City MC
8001 Victoria, Australia

EMail: sever@matilda.vu.edu.au

URL: <http://rgmia.vu.edu.au/SSDragomirWeb.html>



volume 3, issue 4, article 51,
2002.

*Received 5 April, 2002;
accepted 20 May, 2002.*

Communicated by: B. Mond

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract


Using the Ostrowski type inequality for functions of bounded variation, an approximation of the finite Hilbert Transform is given. Some numerical experiments are also provided.

2000 Mathematics Subject Classification: Primary 26D10, 26D15; Secondary 41A55, 47A99.

Key words: Finite Hilbert Transform, Ostrowski's Inequality.

Contents

1	Introduction	3
2	Some Inequalities on the Interval $[a, b]$	4
3	A Quadrature Formula for Equidistant Divisions	15
4	A More General Quadrature Formula	21
5	Numerical Experiments	28
	References	



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 2 of 37

1. Introduction

Cauchy principal value integrals of the form

$$(1.1) \quad (Tf)(a, b; t) = PV \int_a^b \frac{f(\tau)}{\tau - t} d\tau \\ := \lim_{\varepsilon \rightarrow 0^+} \left[\int_a^{t-\varepsilon} \frac{f(\tau)}{\tau - t} d\tau + \int_{t+\varepsilon}^b \frac{f(\tau)}{\tau - t} d\tau \right]$$


play an important role in fields like aerodynamics, the theory of elasticity and other areas of the engineering sciences. They are also helpful tools in some methods for the solution of differential equations (cf., e.g. [23]).

For different approaches in approximating the finite Hilbert transform (1.1) including: interpolatory, noninterpolatory, Gaussian, Chebychevian and spline methods, see for example the papers [1] – [12], [14] – [22], [24] – [33] and the references therein.

In contrast with all these methods, we point out here a new method in approximating the finite Hilbert transform by the use of the Ostrowski inequality for functions of bounded variation established in [13].

For a comprehensive list of papers on Ostrowski's inequality, visit the site <http://rgmia.vu.edu.au>.

Estimates for the error bounds and some numerical examples for the obtained approximation are also presented.



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 3 of 37

2. Some Inequalities on the Interval $[a, b]$

We start with the following lemma proved in [13] dealing with an Ostrowski type inequality for functions of bounded variation.

Lemma 2.1. *Let $u : [a, b] \rightarrow \mathbb{R}$ be a function of bounded variation on $[a, b]$. Then, for all $x \in [a, b]$, we have the inequality:*

$$(2.1) \quad \left| u(x)(b-a) - \int_a^b u(t) dt \right| \leq \left[\frac{1}{2}(b-a) + \left| x - \frac{a+b}{2} \right| \right] \bigvee_a^b(u),$$

where $\bigvee_a^b(u)$ denotes the total variation of u on $[a, b]$.
The constant $\frac{1}{2}$ is the best possible one.

Proof. For the sake of completeness and since this result will be essentially used in what follows, we give here a short proof.

Using the integration by parts formula for the Riemann-Stieltjes integral we have


$$\int_a^x (t-a) du(t) = u(x)(x-a) - \int_a^x u(t) dt$$

and

$$\int_x^b (t-b) du(t) = u(x)(b-x) - \int_x^b u(t) dt.$$

If we add the above two equalities, we get

$$(2.2) \quad u(x)(b-a) - \int_a^b u(t) dt = \int_a^x (t-a) du(t) + \int_x^b (t-b) du(t)$$



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 4 of 37

for any $x \in [a, b]$.

If $p : [c, d] \rightarrow \mathbb{R}$ is continuous on $[c, d]$ and $v : [c, d] \rightarrow \mathbb{R}$ is of bounded variation on $[c, d]$, then:

$$(2.3) \quad \left| \int_c^d p(x) dv(x) \right| \leq \sup_{x \in [c, d]} |p(x)| \bigvee_c^d(u).$$

Using (2.2) and (2.3), we deduce


$$\begin{aligned} \left| u(x)(b-a) - \int_a^b u(t) dt \right| &\leq \left| \int_a^x (t-a) du(t) \right| + \left| \int_x^b (t-b) du(t) \right| \\ &\leq (x-a) \bigvee_a^x(u) + (b-x) \bigvee_x^b(u) \\ &\leq \max\{x-a, b-x\} \left[\bigvee_a^x(u) + \bigvee_x^b(u) \right] \\ &= \left[\frac{1}{2}(b-a) + \left| x - \frac{a+b}{2} \right| \right] \bigvee_a^b(u) \end{aligned}$$

and the inequality (2.1) is proved.

Now, assume that the inequality (2.2) holds with a constant $c > 0$, i.e.,

$$(2.4) \quad \left| u(x)(b-a) - \int_a^b u(t) dt \right| \leq \left[c(b-a) + \left| x - \frac{a+b}{2} \right| \right] \bigvee_a^b(u)$$

for all $x \in [a, b]$.



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 5 of 37

Consider the function $u_0 : [a, b] \rightarrow \mathbb{R}$ given by

$$u_0(x) = \begin{cases} 0 & \text{if } x \in [a, b] \setminus \left\{ \frac{a+b}{2} \right\} \\ 1 & \text{if } x = \frac{a+b}{2}. \end{cases}$$

Then u_0 is of bounded variation on $[a, b]$ and

$$\bigvee_a^b(u_0) = 2, \quad \int_a^b u_0(t) dt = 0.$$

If we apply (2.4) for u_0 and choose $x = \frac{a+b}{2}$, then we get $2c \geq 1$ which implies that $c \geq \frac{1}{2}$ showing that $\frac{1}{2}$ is the best possible constant in (2.1). \square


The best inequality we can get from (2.1) is the following midpoint inequality.

Corollary 2.2. *With the assumptions in Lemma 2.1, we have*

$$(2.5) \quad \left| u\left(\frac{a+b}{2}\right)(b-a) - \int_a^b u(t) dt \right| \leq \frac{1}{2}(b-a) \bigvee_a^b(u).$$

The constant $\frac{1}{2}$ is best possible.

Using the above Ostrowski type inequality we may point out the following result in estimating the finite Hilbert transform.



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 6 of 37

Theorem 2.3. Let $f : [a, b] \rightarrow \mathbb{R}$ be a function such that its derivative $f' : [a, b] \rightarrow \mathbb{R}$ is of bounded variation on $[a, b]$. Then we have the inequality:

$$(2.6) \quad \left| (Tf)(a, b; t) - \frac{f(t)}{\pi} \ln \left(\frac{b-t}{t-a} \right) - \frac{b-a}{\pi} [f; \lambda t + (1-\lambda)b, \lambda t + (1-\lambda)a] \right| \leq \frac{1}{\pi} \left[\frac{1}{2} + \left| \lambda - \frac{1}{2} \right| \right] \left[\frac{1}{2}(b-a) + \left| t - \frac{a+b}{2} \right| \right] \bigvee_a^b(f'),$$

for any $t \in (a, b)$ and $\lambda \in [0, 1)$, where $[f; \alpha, \beta]$ is the divided difference, i.e.,

$$[f; \alpha, \beta] := \frac{f(\alpha) - f(\beta)}{\alpha - \beta}.$$


Proof. Since f' is bounded on $[a, b]$, it follows that f is Lipschitzian on $[a, b]$ and thus the finite Hilbert transform exists everywhere in (a, b) .

As for the function $f_0 : (a, b) \rightarrow \mathbb{R}$, $f_0(t) = 1$, $t \in (a, b)$, we have

$$(Tf_0)(a, b; t) = \frac{1}{\pi} \ln \left(\frac{b-t}{t-a} \right), \quad t \in (a, b),$$

then obviously

$$(2.7) \quad (Tf)(a, b; t) - \frac{f(t)}{\pi} \ln \left(\frac{b-t}{t-a} \right) = \frac{1}{\pi} PV \int_a^b \frac{f(\tau) - f(t)}{\tau - t} d\tau.$$



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 7 of 37

Now, if we choose in (2.1), $u = f'$, $x = \lambda c + (1 - \lambda) d$, $\lambda \in [0, 1]$, then we get

$$|f(d) - f(c) - (d - c)f'(\lambda c + (1 - \lambda)d)| \leq \left[\frac{1}{2}|d - c| + \left| \lambda c + (1 - \lambda)d - \frac{c + d}{2} \right| \right] \left| \bigvee_c^d(f') \right|$$


where $c, d \in (a, b)$, which is equivalent to

$$(2.8) \quad \left| \frac{f(d) - f(c)}{d - c} - f'(\lambda c + (1 - \lambda)d) \right| \leq \left[\frac{1}{2} + \left| \lambda - \frac{1}{2} \right| \right] \left| \bigvee_c^d(f') \right|$$

for any $c, d \in (a, b)$, $c \neq d$.

Using (2.8), we may write

$$(2.9) \quad \left| \frac{1}{\pi} PV \int_a^b \frac{f(\tau) - f(t)}{\tau - t} d\tau - \frac{1}{\pi} PV \int_a^b f'(\lambda t + (1 - \lambda)\tau) d\tau \right| \leq \frac{1}{\pi} \left[\frac{1}{2} + \left| \lambda - \frac{1}{2} \right| \right] PV \int_a^b \left| \bigvee_{\tau}^t(f') \right| dt = \frac{1}{\pi} \left[\frac{1}{2} + \left| \lambda - \frac{1}{2} \right| \right] \left[\int_a^t \left(\bigvee_{\tau}^t(f') \right) dt + \int_t^b \left(\bigvee_t^{\tau}(f') \right) dt \right] \leq \frac{1}{\pi} \left[\frac{1}{2} + \left| \lambda - \frac{1}{2} \right| \right] \left[(t - a) \bigvee_a^t(f') + (b - t) \bigvee_t^b(f') \right] \leq \frac{1}{\pi} \left[\frac{1}{2} + \left| \lambda - \frac{1}{2} \right| \right] \left[\frac{1}{2}(b - a) + \left| t - \frac{a + b}{2} \right| \right] \bigvee_a^b(f').$$



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 8 of 37

Since (for $\lambda \neq 1$)


$$\begin{aligned}
 & \frac{1}{\pi} PV \int_a^b f'(\lambda t + (1 - \lambda)\tau) d\tau \\
 &= \frac{1}{\pi} \lim_{\varepsilon \rightarrow 0^+} \left[\int_a^{t-\varepsilon} + \int_{t+\varepsilon}^b \right] (f'(\lambda t + (1 - \lambda)\tau) d\tau) \\
 &= \frac{1}{\pi} \lim_{\varepsilon \rightarrow 0^+} \left[\frac{1}{1 - \lambda} f(\lambda t + (1 - \lambda)\tau) \Big|_a^{t-\varepsilon} + \frac{1}{1 - \lambda} f(\lambda t + (1 - \lambda)\tau) \Big|_{t+\varepsilon}^b \right] \\
 &= \frac{1}{\pi} \cdot \frac{f(t) - f(\lambda t + (1 - \lambda)a) + f(\lambda t + (1 - \lambda)b) - f(t)}{1 - \lambda} \\
 &= \frac{b - a}{\pi} [f; \lambda t + (1 - \lambda)b, \lambda t + (1 - \lambda)a].
 \end{aligned}$$

Using (2.9) and (2.7), we deduce the desired result (2.6). □

It is obvious that the best inequality we can get from (2.6) is the one for $\lambda = \frac{1}{2}$. Thus, we may state the following corollary.

Corollary 2.4. *With the assumptions of Theorem 2.3, we have*

$$\begin{aligned}
 (2.10) \quad & \left| (Tf)(a, b; t) - \frac{f(t)}{\pi} \ln \left(\frac{b-t}{t-a} \right) - \frac{b-a}{\pi} \left[f; \frac{t+b}{2}, \frac{a+t}{2} \right] \right| \\
 & \leq \frac{1}{2\pi} \left[\frac{1}{2}(b-a) + \left| t - \frac{a+b}{2} \right| \right] \bigvee_a^b(f').
 \end{aligned}$$



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 9 of 37

The above Theorem 2.3 may be used to point out some interesting inequalities for the functions for which the finite Hilbert transforms $(Tf)(a, b; t)$ can be expressed in terms of special functions.

For instance, we have:

1) Assume that $f : [a, b] \subset (0, \infty) \rightarrow \mathbb{R}$, $f(x) = \frac{1}{x}$. Then


$$(Tf)(a, b; t) = \frac{1}{\pi t} \ln \left[\frac{(b-t)a}{(t-a)b} \right], \quad t \in (a, b),$$

$$\begin{aligned} \frac{b-a}{\pi} \cdot [f; \lambda t + (1-\lambda)b, \lambda t + (1-\lambda)a] \\ = -\frac{1}{\pi} \cdot \frac{b-a}{[\lambda t + (1-\lambda)b][\lambda t + (1-\lambda)a]}, \end{aligned}$$

$$\bigvee_a^b(f') = \int_a^b |f''(t)| dt = \frac{b^2 - a^2}{a^2 b^2}.$$

Using the inequality (2.6) we may write that

$$\begin{aligned} \left| \frac{1}{\pi t} \ln \left[\frac{(b-t)a}{(t-a)b} \right] - \frac{1}{\pi t} \ln \left(\frac{b-t}{t-a} \right) + \frac{b-a}{\pi [\lambda t + (1-\lambda)b][\lambda t + (1-\lambda)a]} \right| \\ \leq \frac{1}{\pi} \left[\frac{1}{2} + \left| \lambda - \frac{1}{2} \right| \right] \left[\frac{1}{2} (b-a) + \left| t - \frac{a+b}{2} \right| \right] \cdot \frac{b^2 - a^2}{a^2 b^2} \end{aligned}$$



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents




Go Back

Close

Quit

Page 10 of 37



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 11 of 37

which is equivalent to:

$$(2.11) \quad \left| \frac{b-a}{[\lambda t + (1-\lambda)b][\lambda t + (1-\lambda)a]} - \frac{1}{t} \ln \left(\frac{b}{a} \right) \right| \leq \left[\frac{1}{2} + \left| \lambda - \frac{1}{2} \right| \right] \left[\frac{1}{2}(b-a) + \left| t - \frac{a+b}{2} \right| \right] \cdot \frac{b^2 - a^2}{a^2 b^2}.$$

If we use the notations

$$L(a, b) := \frac{b-a}{\ln b - \ln a} \quad (\text{the logarithmic mean})$$

$$A_\lambda(x, y) := \lambda x + (1-\lambda)y \quad (\text{the weighted arithmetic mean})$$


$$G(a, b) := \sqrt{ab} \quad (\text{the geometric mean})$$

$$A(a, b) := \frac{a+b}{2} \quad (\text{the arithmetic mean})$$

then by (2.11) we deduce

$$\left| \frac{1}{A_\lambda(t, b) A_\lambda(t, a)} - \frac{1}{tL(a, b)} \right| \leq \left[\frac{1}{2} + \left| \lambda - \frac{1}{2} \right| \right] \left[\frac{1}{2}(b-a) + |t - A(a, b)| \right] \frac{2A(a, b)}{G^4(a, b)},$$

giving the following proposition:



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 12 of 37

Proposition 2.5. *With the above assumption, we have*

$$(2.12) \quad |tL(a, b) - A_\lambda(t, b)A_\lambda(t, a)| \\ \leq \frac{2A(a, b)}{G^4(a, b)} \left[\frac{1}{2} + \left| \lambda - \frac{1}{2} \right| \right] \left[\frac{1}{2} (b - a) + \left| t - \frac{a + b}{2} \right| \right] \\ \times tA_\lambda(t, b)A_\lambda(t, a)L(a, b)$$

for any $t \in (a, b)$, $\lambda \in [0, 1)$.

In particular, for $t = A(a, b)$ and $\lambda = \frac{1}{2}$, we get

$$(2.13) \quad \left| A(a, b)L(a, b) - \frac{(A(a, b) + a)(A(a, b) + b)}{4} \right| \\ \leq \frac{1}{2} \cdot \frac{A^2(a, b)}{G^4(a, b)} \cdot \frac{(A(a, b) + a)(A(a, b) + b)}{4} L(a, b).$$

2) Assume that $f : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = \exp(x)$. Then

$$(Tf)(a, b; t) = \frac{\exp(t)}{\pi} [Ei(b - t) - Ei(a - t)],$$

where

$$Ei(z) := PV \int_{-\infty}^z \frac{\exp(t)}{t} dt, \quad z \in \mathbb{R}.$$

Also, we have:

$$\frac{b - a}{\pi} [\exp; \lambda t + (1 - \lambda)b, \lambda t + (1 - \lambda)a] \\ = \frac{1}{\pi} \cdot \frac{\exp(\lambda t + (1 - \lambda)b) - \exp(\lambda t + (1 - \lambda)a)}{1 - \lambda},$$

$$\bigvee_a^b (f') = \int_a^b |f''(t)| dt = \exp(b) - \exp(a).$$

Using the inequality (2.6) we may write:

$$(2.14) \quad \left| \exp(t) \left[Ei(b-t) - Ei(a-t) - \ln\left(\frac{b-t}{t-a}\right) \right] - \frac{\exp(\lambda t + (1-\lambda)b) - \exp(\lambda t + (1-\lambda)a)}{1-\lambda} \right| \leq \left[\frac{1}{2} + \left| \lambda - \frac{1}{2} \right| \right] \left[\frac{1}{2}(b-a) + \left| t - \frac{a+b}{2} \right| \right] [\exp(b) - \exp(a)]$$


for any $t \in (a, b)$.

If in (2.14) we make $\lambda = \frac{1}{2}$ and $t = \frac{a+b}{2}$, we get

$$\left| \exp\left(\frac{a+b}{2}\right) Ei\left(\frac{b-a}{2}\right) - 2 \left[\exp\left(\frac{a+3b}{4}\right) - \exp\left(\frac{3a+b}{4}\right) \right] \right| \leq \frac{1}{4}(b-a) [\exp(b) - \exp(a)],$$

which is equivalent to:

$$\left| Ei\left(\frac{b-a}{2}\right) - 2 \left[\exp\left(\frac{b-a}{4}\right) - \exp\left(-\frac{b-a}{4}\right) \right] \right| \leq \frac{1}{4}(b-a) \left[\exp\left(\frac{b-a}{2}\right) - \exp\left(-\frac{b-a}{2}\right) \right].$$



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents




Go Back

Close

Quit

Page 13 of 37



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

If in this inequality we make $\frac{b-a}{2} = z > 0$, then we get

$$(2.15) \quad \left| Ei(z) - 2 \left[\exp\left(\frac{z}{2}\right) - \exp\left(-\frac{z}{2}\right) \right] \right| \leq \frac{1}{2} z [\exp(z) - \exp(-z)]$$

for any $z > 0$.

Consequently, we may state the following proposition.

Proposition 2.6. *With the above assumptions, we have*

$$(2.16) \quad \left| Ei(z) - 4 \sinh\left(\frac{1}{2}z\right) \right| \leq z \sinh(z)$$

for any $z > 0$.

The reader may get other similar inequalities for special functions if appropriate examples of functions f are chosen.

Title Page

Contents



Go Back

Close

Quit

Page 14 of 37

3. A Quadrature Formula for Equidistant Divisions

The following lemma is of interest in itself.

Lemma 3.1. *Let $u : [a, b] \rightarrow \mathbb{R}$ be a function of bounded variation on $[a, b]$. Then for all $n \geq 1$, $\lambda_i \in [0, 1)$ ($i = 0, \dots, n-1$) and $t, \tau \in [a, b]$ with $t \neq \tau$, we have the inequality:*


$$(3.1) \quad \left| \frac{1}{\tau - t} \int_t^\tau u(s) ds - \frac{1}{n} \sum_{i=0}^{n-1} u \left[t + (i+1 - \lambda_i) \frac{\tau - t}{n} \right] \right| \leq \frac{1}{n} \left[\frac{1}{2} + \max_{i=0, n-1} \left| \lambda_i - \frac{1}{2} \right| \right] \left| \bigvee_t^\tau(u) \right|.$$

Proof. Consider the equidistant division of $[t, \tau]$ (if $t < \tau$) or $[\tau, t]$ (if $\tau < t$) given by

$$(3.2) \quad E_n : x_i = t + i \cdot \frac{\tau - t}{n}, \quad i = \overline{0, n}.$$

Then the points $\xi_i = \lambda_i \left[t + i \cdot \frac{\tau - t}{n} \right] + (1 - \lambda_i) \left[t + (i+1) \cdot \frac{\tau - t}{n} \right]$ ($\lambda_i \in [0, 1]$, $i = \overline{0, n-1}$) are between x_i and x_{i+1} . We observe that we may write for simplicity $\xi_i = t + (i+1 - \lambda_i) \frac{\tau - t}{n}$ ($i = \overline{0, n-1}$). We also have

$$\begin{aligned} \xi_i - \frac{x_i + x_{i+1}}{2} &= \frac{\tau - t}{2n} (1 - 2\lambda_i), \\ \xi_i - x_i &= (1 - \lambda_i) \frac{\tau - t}{n} \end{aligned}$$



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 15 of 37

and

$$x_{i+1} - \xi_i = \lambda_i \cdot \frac{\tau - t}{n}$$

for any $i = \overline{0, n-1}$.

If we apply the inequality (2.1) on the interval $[x_i, x_{i+1}]$ and the intermediate point ξ_i ($i = \overline{0, n-1}$), then we may write that


$$(3.3) \quad \left| \frac{\tau - t}{n} u \left(t + (i + 1 - \lambda_i) \frac{\tau - t}{n} \right) - \int_{x_i}^{x_{i+1}} u(s) ds \right| \\ \leq \left[\frac{1}{2} \cdot \frac{|\tau - t|}{n} + \left| \frac{\tau - t}{2n} (1 - 2\lambda_i) \right| \right] \left| \bigvee_{x_i}^{x_{i+1}}(u) \right|.$$

Summing, we get

$$\left| \int_t^\tau u(s) ds - \frac{\tau - t}{n} \sum_{i=0}^{n-1} u \left[t + (i + 1 - \lambda_i) \frac{\tau - t}{n} \right] \right| \\ \leq \frac{|\tau - t|}{2n} \sum_{i=0}^{n-1} [1 + |1 - 2\lambda_i|] \left| \bigvee_{x_i}^{x_{i+1}}(u) \right| \\ = \frac{|\tau - t|}{n} \left[\frac{1}{2} + \max_{i=0, n-1} \left| \lambda_i - \frac{1}{2} \right| \right] \left| \bigvee_t^\tau(u) \right|,$$

which is equivalent to (3.1). □

We may now state the following theorem in approximating the finite Hilbert transform of a differentiable function with the derivative of bounded variation on $[a, b]$.



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 16 of 37

Theorem 3.2. Let $f : [a, b] \rightarrow \mathbb{R}$ be a differentiable function such that its derivative f' is of bounded variation on $[a, b]$. If $\lambda = (\lambda_i)_{i=0, n-1}$, $\lambda_i \in [0, 1]$ ($i = \overline{0, n-1}$) and

$$(3.4) \quad S_n(f; \lambda, t)$$

$$:= \frac{b-a}{\pi n} \sum_{i=0}^{n-1} \left[f; (i+1-\lambda_i) \frac{b-t}{n} + t, (i+1-\lambda_i) \frac{a-t}{n} + t \right],$$

then we have the estimate:

$$(3.5) \quad \left| (Tf)(a, b; t) - \frac{f(t)}{\pi} \ln \left(\frac{b-t}{t-a} \right) - S_n(f; \lambda, t) \right|$$


$$\leq \frac{b-a}{n\pi} \left[\frac{1}{2} + \max_{i=0, n-1} \left| \lambda_i - \frac{1}{2} \right| \right] \left[\frac{1}{2}(b-a) + \left| t - \frac{a+b}{2} \right| \right] \bigvee_a^b(f')$$

$$\leq \frac{b-a}{n\pi} \bigvee_a^b(f').$$

Proof. Applying Lemma 3.1 for the function f' , we may write that

$$(3.6) \quad \left| \frac{f(\tau) - f(t)}{\tau - t} - \frac{1}{n} \sum_{i=0}^{n-1} f' \left[t + (i+1-\lambda_i) \frac{\tau-t}{n} \right] \right|$$

$$\leq \frac{1}{n} \left[\frac{1}{2} + \max_{i=0, n-1} \left| \lambda_i - \frac{1}{2} \right| \right] \left| \bigvee_t^\tau(f') \right|$$



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 17 of 37


for any $t, \tau \in [a, b], t \neq \tau$.

Consequently, we have

$$\begin{aligned}
 (3.7) \quad & \left| \frac{1}{\pi} PV \int_a^b \frac{f(\tau) - f(t)}{\tau - t} d\tau \right. \\
 & \quad \left. - \frac{1}{\pi n} \sum_{i=0}^{n-1} PV \int_a^b f' \left[t + (i+1 - \lambda_i) \frac{\tau - t}{n} \right] d\tau \right| \\
 & \leq \frac{1}{n\pi} \left[\frac{1}{2} + \max_{i=0, n-1} \left| \lambda_i - \frac{1}{2} \right| \right] PV \int_a^b \left| \bigvee_t^\tau (f') \right| d\tau \\
 & \leq \frac{1}{n\pi} \left[\frac{1}{2} + \max_{i=0, n-1} \left| \lambda_i - \frac{1}{2} \right| \right] \left[\frac{1}{2} (b-a) + \left| t - \frac{a+b}{2} \right| \right] \bigvee_a^b (f').
 \end{aligned}$$

On the other hand

$$\begin{aligned}
 (3.8) \quad & PV \int_a^b f' \left[t + (i+1 - \lambda_i) \frac{\tau - t}{n} \right] d\tau \\
 & = \lim_{\varepsilon \rightarrow 0^+} \left[\int_a^{t-\varepsilon} + \int_{t+\varepsilon}^b \right] \left(f' \left[t + (i+1 - \lambda_i) \frac{\tau - t}{n} \right] d\tau \right) \\
 & = \lim_{\varepsilon \rightarrow 0^+} \left[\frac{n}{i+1 - \lambda_i} f \left(t + (i+1 - \lambda_i) \frac{\tau - t}{n} \right) \Big|_a^{t-\varepsilon} \right. \\
 & \quad \left. + \frac{n}{i+1 - \lambda_i} f \left(t + (i+1 - \lambda_i) \frac{\tau - t}{n} \right) \Big|_{t+\varepsilon}^b \right]
 \end{aligned}$$



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 18 of 37

$$\begin{aligned}
&= \frac{n}{i+1-\lambda_i} \left[f \left(t + (i+1-\lambda_i) \frac{b-t}{n} \right) \right. \\
&\quad \left. - f \left(t + (i+1-\lambda_i) \frac{a-t}{n} \right) \right] \\
&= (b-a) \left[f; t + (i+1-\lambda_i) \frac{b-t}{n}, (i+1-\lambda_i) \frac{a-t}{n} + t \right].
\end{aligned}$$

Since (see for example (2.7)),

$$(Tf)(a, b; t) = \frac{1}{\pi} PV \int_a^b \frac{f(\tau) - f(t)}{\tau - t} d\tau + \frac{f(t)}{\pi} \ln \left(\frac{b-t}{t-a} \right)$$

for $t \in (a, b)$, then by (3.7) and (3.8) we deduce the desired estimate (3.5). \square

Remark 3.1. For $n = 1$, we recapture the inequality (2.6).


Corollary 3.3. With the assumptions of Theorem 3.2, we have

$$(3.9) \quad (Tf)(a, b; t) = \frac{f(t)}{\pi} \ln \left(\frac{b-t}{t-a} \right) + \lim_{n \rightarrow \infty} S_n(f; \lambda, t)$$

uniformly by rapport of $t \in (a, b)$ and λ with $\lambda_i \in [0, 1)$ ($i \in \mathbb{N}$).

Remark 3.2. If one needs to approximate the finite Hilbert Transform $(Tf)(a, b; t)$ in terms of

$$\frac{f(t)}{\pi} \ln \left(\frac{b-t}{t-a} \right) + S_n(f; \lambda, t)$$



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 19 of 37

with the accuracy $\varepsilon > 0$ (ε small), then the theoretical minimal number n_ε to be chosen is:

$$(3.10) \quad n_\varepsilon := \left[\frac{b-a}{\varepsilon\pi} \bigvee_a^b (f') \right] + 1$$

where $[\alpha]$ is the integer part of α .

It is obvious that the best inequality we can get in (3.5) is for $\lambda_i = \frac{1}{2}$ ($i = \overline{0, n-1}$) obtaining the following corollary.

Corollary 3.4. Let f be as in Theorem 3.2. Define


$$(3.11) \quad M_n(f; t) := \frac{b-a}{\pi n} \sum_{i=0}^{n-1} \left[f; \left(i + \frac{1}{2} \right) \frac{b-t}{n} + t, \left(i + \frac{1}{2} \right) \frac{a-t}{n} + t \right].$$

Then we have the estimate

$$(3.12) \quad \left| (Tf)(a, b; t) - \frac{f(t)}{\pi} \ln \left(\frac{b-t}{t-a} \right) - M_n(f; t) \right| \leq \frac{b-a}{2n\pi} \left[\frac{1}{2} (b-a) + \left| t - \frac{a+b}{2} \right| \right] \bigvee_a^b (f')$$

for any $t \in (a, b)$.

This rule will be numerically implemented in Section 5 for different choices of f and n .



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 20 of 37

4. A More General Quadrature Formula

We may state the following lemma.

Lemma 4.1. *Let $u : [a, b] \rightarrow \mathbb{R}$ be a function of bounded variation on $[a, b]$, $0 = \mu_0 < \mu_1 < \dots < \mu_{n-1} < \mu_n = 1$ and $\nu_i \in [\mu_i, \mu_{i+1}]$, $i = \overline{0, n-1}$. Then for any $t, \tau \in [a, b]$ with $t \neq \tau$, we have the inequality:*

$$(4.1) \quad \left| \frac{1}{\tau - t} \int_t^\tau u(s) ds - \sum_{i=0}^{n-1} (\mu_{i+1} - \mu_i) u[(1 - \nu_i)t + \nu_i\tau] \right| \\ \leq \left[\frac{1}{2} \Delta_n(\mu) + \max_{i=0, n-1} \left| \nu_i - \frac{\mu_i + \mu_{i+1}}{2} \right| \right] \left| \bigvee_t^\tau(u) \right|,$$

where $\Delta_n(\mu) := \max_{i=0, n-1} (\mu_{i+1} - \mu_i)$.

Proof. Consider the division of $[t, \tau]$ (if $t < \tau$) or $[\tau, t]$ (if $\tau < t$) given by


$$(4.2) \quad I_n : x_i := (1 - \mu_i)t + \mu_i\tau \quad (i = \overline{0, n}).$$

Then the points $\xi_i := (1 - \nu_i)t + \nu_i\tau$ ($i = \overline{0, n-1}$) are between x_i and x_{i+1} . We have

$$x_{i+1} - x_i = (\mu_{i+1} - \mu_i)(\tau - t) \quad (i = \overline{0, n-1})$$

and

$$\xi_i - \frac{x_i + x_{i+1}}{2} = \left(\nu_i - \frac{\mu_i + \mu_{i+1}}{2} \right) (\tau - t) \quad (i = \overline{0, n-1}).$$



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 21 of 37

Applying the inequality (2.1) on $[x_i, x_{i+1}]$ with the intermediate points ξ_i ($i = \overline{0, n-1}$), we get

$$\left| \int_{x_i}^{x_{i+1}} u(s) ds - (\mu_{i+1} - \mu_i) (\tau - t) u [(1 - \nu_i) t + \nu_i \tau] \right| \\ \leq \left[\frac{1}{2} (\mu_{i+1} - \mu_i) |\tau - t| + |\tau - t| \left| \nu_i - \frac{\mu_i + \mu_{i+1}}{2} \right| \right] \left| \bigvee_{x_i}^{x_{i+1}} (u) \right|$$


for any $i = \overline{0, n-1}$. Summing over i , using the generalised triangle inequality and dividing by $|t - \tau| > 0$, we obtain

$$\left| \frac{1}{\tau - t} \int_a^b u(s) ds - \sum_{i=0}^{n-1} (\mu_{i+1} - \mu_i) u [(1 - \nu_i) t + \nu_i \tau] \right| \\ \leq \sum_{i=0}^{n-1} \left[\frac{1}{2} (\mu_{i+1} - \mu_i) + \left| \nu_i - \frac{\mu_i + \mu_{i+1}}{2} \right| \right] \left| \bigvee_{x_i}^{x_{i+1}} (u) \right| \\ \leq \left[\frac{1}{2} \Delta_n(\mu) + \max_{i=0, n-1} \left| \nu_i - \frac{\mu_i + \mu_{i+1}}{2} \right| \right] \left| \bigvee_t^\tau (u) \right|$$

and the inequality (4.1) is proved. \square

The following theorem holds.

Theorem 4.2. Let $f : [a, b] \rightarrow \mathbb{R}$ be a differentiable function such that its derivative f' is of bounded variation on $[a, b]$. If $0 = \mu_0 < \mu_1 < \dots < \mu_{n-1} <$



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 22 of 37

$\mu_n = 1$ and $\nu_i \in [\mu_i, \mu_{i+1}]$, ($i = \overline{0, n-1}$), then

$$(4.3) \quad (Tf)(a, b; t) = \frac{f(t)}{\pi} \ln \left(\frac{b-t}{t-a} \right) + \frac{1}{\pi} Q_n(\mu, \nu, t) + W_n(\mu, \nu, t)$$

for any $t \in (a, b)$, where


$$(4.4) \quad Q_n(\mu, \nu, t) := \mu_1 f'(t)(b-a) + (b-a) \sum_{i=1}^{n-2} \left\{ (\mu_{i+1} - \mu_i) \times [f; (1-\nu_i)t + \nu_i b, (1-\nu_i)t + \nu_i a] \right\} + (1 - \mu_{n-1}) [f(b) - f(a)]$$

if $\nu_0 = 0, \nu_{n-1} = 1$,

$$(4.5) \quad Q_n(\mu, \nu, t) := \mu_1 f'(t)(b-a) + (b-a) \sum_{i=1}^{n-1} (\mu_{i+1} - \mu_i) \times [f; (1-\nu_i)t + \nu_i b, (1-\nu_i)t + \nu_i a]$$

if $\nu_0 = 0, \nu_{n-1} < 1$,

$$(4.6) \quad Q_n(\mu, \nu, t) := (b-a) \sum_{i=1}^{n-2} (\mu_{i+1} - \mu_i) \times [f; (1-\nu_i)t + \nu_i b, (1-\nu_i)t + \nu_i a] + (1 - \mu_{n-1}) [f(b) - f(a)]$$



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 23 of 37

if $\nu_0 > 0$, $\nu_{n-1} = 1$ and

$$(4.7) \quad Q_n(\mu, \nu, t)$$

$$:= (b-a) \sum_{i=1}^{n-1} (\mu_{i+1} - \mu_i) [f; (1-\nu_i)t + \nu_i b, (1-\nu_i)t + \nu_i a]$$


if $\nu_0 > 0$, $\nu_{n-1} < 1$.

In all cases, the remainder satisfies the estimate:

$$(4.8) \quad |W_n(\mu, \nu, t)| \leq \frac{1}{\pi} \left[\frac{1}{2} \Delta_n(\mu) + \max_{i=0, n-1} \left| \nu_i - \frac{\mu_i + \mu_{i+1}}{2} \right| \right] \\ \times \left[\frac{1}{2} (b-a) + \left| t - \frac{a+b}{2} \right| \right] \bigvee_a^b(f') \\ \leq \frac{1}{\pi} \Delta_n(\mu) \left[\frac{1}{2} (b-a) + \left| t - \frac{a+b}{2} \right| \right] \left| \bigvee_a^b(f') \right| \\ \leq \frac{1}{\pi} \Delta_n(\mu) (b-a) \bigvee_a^b(f').$$

Proof. If we apply Lemma 4.1 for the function f' , we may write that

$$\left| \frac{f(\tau) - f(t)}{\tau - t} - \sum_{i=0}^{n-1} (\mu_{i+1} - \mu_i) f'[(1-\nu_i)t + \nu_i \tau] \right| \\ \leq \left[\frac{1}{2} \Delta_n(\mu) + \max_{i=0, n-1} \left| \nu_i - \frac{\mu_i + \mu_{i+1}}{2} \right| \right] \left| \bigvee_t^\tau(f') \right|$$



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 24 of 37


for any $t, \tau \in [a, b]$, $t \neq \tau$.

Taking the PV in both sides, we may write that

$$(4.9) \quad \left| \frac{1}{\pi} PV \int_a^b \frac{f(\tau) - f(t)}{\tau - t} d\tau - \frac{1}{\pi} PV \int_a^b \left(\sum_{i=0}^{n-1} (\mu_{i+1} - \mu_i) f' [(1 - \nu_i)t + \nu_i\tau] \right) d\tau \right| \leq \frac{1}{\pi} \left[\frac{1}{2} \Delta_n(\mu) + \max_{i=0, n-1} \left| \nu_i - \frac{\mu_i + \mu_{i+1}}{2} \right| \right] PV \int_a^b \left| \bigvee_t^{\tau} (f') \right| d\tau.$$

If $\nu_0 = 0$, $\nu_{n-1} = 1$, then

$$\begin{aligned} & PV \int_a^b \left(\sum_{i=0}^{n-1} (\mu_{i+1} - \mu_i) f' [(1 - \nu_i)t + \nu_i\tau] \right) d\tau \\ &= PV \int_a^b \mu_1 f'(t) d\tau + \sum_{i=1}^{n-2} (\mu_{i+1} - \mu_i) PV \int_a^b f' [(1 - \nu_i)t + \nu_i\tau] d\tau \\ &\quad + (1 - \mu_{n-1}) PV \int_a^b f'(\tau) d\tau \\ &= \mu_1 f'(t) (b - a) + (b - a) \sum_{i=1}^{n-2} (\mu_{i+1} - \mu_i) \\ &\quad \times [f; (1 - \nu_i)t + \nu_i b, (1 - \nu_i)t + \nu_i a] + (1 - \mu_{n-1}) [f(b) - f(a)]. \end{aligned}$$



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 25 of 37

If $\nu_0 = 0, \nu_{n-1} < 1$, then


$$\begin{aligned}
 PV \int_a^b \left(\sum_{i=0}^{n-1} (\mu_{i+1} - \mu_i) f' [(1 - \nu_i) t + \nu_i \tau] \right) d\tau \\
 = \mu_1 f'(t) (b - a) + (b - a) \sum_{i=1}^{n-1} (\mu_{i+1} - \mu_i) \\
 \times [f; (1 - \nu_i) t + \nu_i b, (1 - \nu_i) t + \nu_i a].
 \end{aligned}$$

If $\nu_0 > 0, \nu_{n-1} = 1$, then

$$\begin{aligned}
 PV \int_a^b \left(\sum_{i=0}^{n-1} (\mu_{i+1} - \mu_i) f' [(1 - \nu_i) t + \nu_i \tau] \right) d\tau \\
 = (b - a) \sum_{i=1}^{n-2} (\mu_{i+1} - \mu_i) [f; (1 - \nu_i) t + \nu_i b, (1 - \nu_i) t + \nu_i a] \\
 + (1 - \mu_{n-1}) [f(b) - f(a)].
 \end{aligned}$$

and, finally, if $\nu_0 > 0, \nu_{n-1} < 1$, then

$$\begin{aligned}
 PV \int_a^b \left(\sum_{i=0}^{n-1} (\mu_{i+1} - \mu_i) f' [(1 - \nu_i) t + \nu_i \tau] \right) d\tau \\
 = (b - a) \sum_{i=1}^{n-1} (\mu_{i+1} - \mu_i) [f; (1 - \nu_i) t + \nu_i b, (1 - \nu_i) t + \nu_i a].
 \end{aligned}$$



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 26 of 37

Since


$$PV \int_a^b \left| \bigvee_t^\tau (f') \right| d\tau \leq \left[\frac{1}{2} (b-a) + \left| t - \frac{a+b}{2} \right| \right] \bigvee_a^b (f')$$

and

$$(Tf)(a, b; t) = \frac{1}{\pi} PV \int_a^b \frac{f(\tau) - f(t)}{\tau - t} d\tau + \frac{f(t)}{\pi} \ln \left(\frac{b-t}{t-a} \right),$$

then by (4.9) we deduce (4.3).

□



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 27 of 37

5. Numerical Experiments

For a function $f : [a, b] \rightarrow \mathbb{R}$, we may consider the quadrature formula

$$E_n(f; a, b, t) := \frac{f(t)}{\pi} \ln \left(\frac{b-t}{t-a} \right) + M_n(f; t), t \in [a, b].$$

As shown above in Section 4, $E_n(f; a, b, t)$ provides an approximation for the Finite Hilbert Transform $(Tf)(a, b; t)$ and the error estimate fulfils the bound described in (2.3).

If we consider the function $f : [-1, 1] \rightarrow \mathbb{R}$, $f(x) = \exp(x)$, then the exact value of the Hilbert transform is


$$(Tf)(a, b; t) = \frac{\exp(t)Ei(1-t) - \exp(t)Ei(-1-t)}{\pi}, t \in [-1, 1]$$

and the plot of this function is embodied in Figure 1.

If we implement the quadrature formula provided by $E_n(f; a, b, t)$ using Maple 6 and chose the value of $n = 100$, then the error $Er(f; a, b, t) := (Tf)(a, b; t) - E_n(f; a, b, t)$ has the variation described in the Figure 2.

For $n = 1,000$, the plot of $Er(f; a, b, t)$ is embodied in the following Figure 3.

Now, if we consider another function, $f : [-1, 1] \rightarrow \mathbb{R}$, $f(x) = \sin x$, then



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 28 of 37




Figure 1:

the exact value of the Hilbert transform is


$$(Tf)(a, b; t) = \frac{-Si(-1+t)\cos(t) + Ci(1-t)\sin(t)}{\pi} + \frac{Si(t+1)\cos(t) - \sin(t)Ci(t+1)}{\pi}, \quad t \in [-1, 1];$$

where

$$Si(x) = \int_0^x \frac{\sin(t)}{t} dt, \quad Ci(x) = \gamma + \ln x + \int_0^x \frac{\cos(t) - 1}{t} dt;$$

having the plot embodied in the following Figure 4.

If we choose the value of $n = 100$, then the error $Er(f; a, b, t)$ for the function $f(x) = \sin x, x \in [-1, 1]$ has the variation described in the Figure 5 below.



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 29 of 37




Figure 2:


Moreover, for $n = 100,000$, the behaviour of $Er(f; a, b, t)$ is plotted in Figure 6.

Finally, if we choose the function $f : [-1, 1] \rightarrow \mathbb{R}$, $f(x) = \sin(x^2)$, the Maple 6 is unable to produce an exact value of the finite Hilbert transform. If we use our formula

$$E_n(f; a, b, t) := \frac{f(t)}{\pi} \ln \left(\frac{b-t}{t-a} \right) + M_n(f; t), t \in [a, b]$$

for $n = 1,000$, then we can produce the plot in Figure 7.

Taking into account the bound (3.12) we know that the accuracy of the plot in Figure 7 is at least of order 10^{-5} .



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 30 of 37




Figure 3:





Figure 4:



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 31 of 37




Figure 5:





Figure 6:



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents




Go Back

Close

Quit

Page 32 of 37



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 33 of 37





Figure 7:

References

- [1] G. CRISCUOLO AND G. MASTROIANNI, Formule gaussiane per il calcolo di integrali a valor principale secondo e Cauchy loro convergenza, *Calcolo*, **22** (1985), 391–411.
- [2] G. CRISCUOLO AND G. MASTROIANNI, Sulla convergenza di alcune formule di quadratura per il calcolo di integrali a valor principale secondo Cauchy, *Anal. Numér. Théor. Approx.*, **14** (1985), 109–116.
- [3] G. CRISCUOLO AND G. MASTROIANNI, On the convergence of an interpolatory product rule for evaluating Cauchy principal value integrals, *Math. Comp.*, **48** (1987), 725–735.
- [4] G. CRISCUOLO AND G. MASTROIANNI, Mean and uniform convergence of quadrature rules for evaluating the finite Hilbert transform, in: A. Pinkus and P. Nevai, Eds., *Progress in Approximation Theory* (Academic Press, Orlando, 1991), 141–175.
- [5] C. DAGNINO, V. DEMICHELIS AND E. SANTI, Numerical integration based on quasi-interpolating splines, *Computing*, **50** (1993), 149–163.
- [6] C. DAGNINO, V. DEMICHELIS AND E. SANTI, An algorithm for numerical integration based on quasi-interpolating splines, *Numer. Algorithms*, **5** (1993), 443–452.
- [7] C. DAGNINO, V. DEMICHELIS AND E. SANTI, Local spline approximation methods for singular product integration, *Approx. Theory Appl.*, **12** (1996), 1–14.



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents




Go Back

Close

Quit

Page 34 of 37

- [8] C. DAGNINO AND P. RABINOWITZ, Product integration of singular integrands based on quasi-interpolating splines, *Comput. Math.*, to appear, special issue dedicated to 100th birthday of Cornelius Lanczos.
- [9] P.J. DAVIS AND P. RABINOWITZ, *Methods of Numerical Integration*, (Academic Press, Orlando, 2nd Ed., 1984).
- [10] K. DIETHELM, Non-optimality of certain quadrature rules for Cauchy principal value integrals, *Z. Angew. Math. Mech.*, **74**(6) (1994), T689–T690.
- [11] K. DIETHELM, Peano kernels and bounds for the error constants of Gaussian and related quadrature rules for Cauchy principal value integrals, *Numer. Math.*, to appear.
- [12] K. DIETHELM, Uniform convergence to optimal order quadrature rules for Cauchy principal value integrals, *J. Comput. Appl. Math.*, **56**(3) (1994), 321–329.
- [13] S.S. DRAGOMIR, On the Ostrowski integral inequality for mappings of bounded variation and applications, *Math. Ineq. & Appl.*, **3**(1) (2001), 59–66.
- [14] D. ELLIOTT, On the convergence of Hunter’s quadrature rule for Cauchy principal value integrals, *BIT*, **19** (1979), 457–462.
- [15] D. ELLIOTT AND D.F. PAGET, On the convergence of a quadrature rule for evaluating certain Cauchy principal value integrals, *Numer. Math.*, **23** (1975), 311–319.



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents




Go Back

Close

Quit

Page 35 of 37

- [16] D. ELLIOTT AND D.F. PAGET, On the convergence of a quadrature rule for evaluating certain Cauchy principal value integrals: An addendum, *Numer. Math.*, **25** (1976), 287–289.
- [17] D. ELLIOTT AND D.F. PAGET, Gauss type quadrature rules for Cauchy principal value integrals, *Math. Comp.*, **33** (1979), 301–309.
- [18] T. HASEGAWA AND T. TORII, An automatic quadrature for Cauchy principal value integrals, *Math. Comp.*, **56** (1991), 741–754.
- [19] T. HASEGAWA AND T. TORII, Hilbert and Hadamard transforms by generalised Chebychev expansion, *J. Comput. Appl. Math.*, **51** (1994), 71–83.
- [20] D.B. HUNTER, Some Gauss type formulae for the evaluation of Cauchy principal value integrals, *Numer. Math.*, **19** (1972), 419–424.
- [21] N.I. IOAKIMIDIS, Further convergence results for two quadrature rules for Cauchy type principal value integrals, *Appl. Mat.*, **27** (1982), 457–466.
- [22] N.I. IOAKIMIDIS, On the uniform convergence of Gaussian quadrature rules for Cauchy principal value integrals and their derivatives, *Math. Comp.*, **44** (1985), 191–198.
- [23] A.I. KALANDIYA, *Mathematical Methods of Two-Dimensional Elasticity*, (Mir, Moscow, 1st English ed., 1978).
- [24] G. MONEGATO, The numerical evaluation of one-dimensional Cauchy principal value integrals, *Computing*, **29** (1982), 337–354.



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents




Go Back

Close

Quit

Page 36 of 37

- [25] G. MONEGATO, On the weights of certain quadratures for the numerical evaluation of Cauchy principal value integrals and their derivatives, *Numer. Math.*, **50** (1987), 273–281.
- [26] D.F. PAGET AND D. ELLIOTT, An algorithm for the numerical evaluation of certain Cauchy principal value integrals, *Numer. Math.*, **19** (1972), 373–385.
- [27] P. RABINOWITZ, On an interpolatory product rule for evaluating Cauchy principal value integrals, *BIT*, **29** (1989), 347–355.
- [28] P. RABINOWITZ, Numerical integration based on approximating splines, *J. Comput. Appl. Math.*, **33** (1990), 73–83.
- [29] P. RABINOWITZ, Application of approximating splines for the solution of Cauchy singular integral equations, *Appl. Numer. Math.*, **15** (1994), 285–297.
- [30] P. RABINOWITZ, Numerical evaluation of Cauchy principal value integrals with singular integrands, *Math. Comput.*, **55** (1990), 265–276.
- [31] P. RABINOWITZ AND E. SANTI, On the uniform convergence of Cauchy principle value of quasi-interpolating splines, *Bit*, **35** (1995), 277–290.
- [32] M.S. SHESHKO, On the convergence of quadrature processes for a singular integral, *Izv. Vyssh. Uohebn. Zaved Mat.*, **20** (1976), 108–118.
- [33] G. TSAMASPHYROS AND P.S. THEOCARIS, On the convergence of some quadrature rules for Cauchy principal-value and finite-part integrals, *Computing*, **31** (1983), 105–114.



Approximating the Finite Hilbert Transform via an Ostrowski Type Inequality for Functions of Bounded Variation

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 37 of 37