

Journal of Inequalities in Pure and Applied Mathematics

ON SOME RETARDED INTEGRAL INEQUALITIES AND APPLICATIONS

B.G. PACHPATTE

57, Shri Niketen Colony
Aurangabad - 431 001,
(Maharashtra) India.

EMail: bgpachpatte@hotmail.com

©2000 Victoria University
ISSN (electronic): 1443-5756
056-01



volume 3, issue 2, article 18,
2002.

Received 11 July, 2001;
accepted 13 November, 2001.
Communicated by: S.S. Dragomir

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

Abstract


The aim of this paper is to establish explicit bounds on certain retarded integral inequalities which can be used as convenient tools in some applications. The two independent variable generalizations of the main results and some applications are also given.

2000 Mathematics Subject Classification: 26D15, 26D20.

Key words: Retarded Integral Inequalities, Explicit Bounds, Two Independent Variable Generalizations, Qualitative Study, Nondecreasing, Change of Variable, Integrodifferential Equation, Uniqueness of Solutions.

Contents

1	Introduction	3
2	Statement of Results	4
3	Proofs of Theorems 2.1 and 2.2	7
4	Some Applications	11
	References	



On Some Retarded Integral Inequalities and Applications

B.G. Pachpatte

Title Page

Contents



Go Back


Close

Quit

Page 2 of 16

1. Introduction

Integral inequalities which provide explicit bounds on unknown functions have played a fundamental role in the development of the theory of differential and integral equations. Over the years, various investigators have discovered many useful integral inequalities in order to achieve a diversity of desired goals, see [1] – [6] and the references given therein. In a recent paper [5] Lipovan has given a useful nonlinear generalisation of the celebrated Gronwall inequality and presented some of its applications. However, the integral inequalities available in the literature do not apply directly in certain general situations and it is desirable to find integral inequalities useful in some new applications. The main purpose of the present paper is to establish explicit bounds on more general retarded integral inequalities which can be used as tools in the qualitative study of certain retarded integrodifferential equations. Some immediate applications of one of the result to convey the importance of our results to the literature are also given.



**On Some Retarded Integral
Inequalities and Applications**

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 3 of 16

2. Statement of Results

In what follows, \mathbb{R} denotes the set of real numbers, $\mathbb{R}_+ = [0, \infty)$, $I = [t_0, T)$, $J_1 = [x_0, X)$, $J_2 = [y_0, Y)$ are the given subsets of \mathbb{R} , $\Delta = J_1 \times J_2$ and $'$ denotes the derivative. The partial derivatives of a function $z(x, y)$, $x, y \in \mathbb{R}$ with respect to x and y are denoted by $D_1 z(x, y)$ and $D_2 z(x, y)$ respectively.

Our main results are given in the following theorems.

Theorem 2.1. *Let $u(t), a(t) \in C(I, \mathbb{R}_+)$, $b(t, s) \in C(I^2, \mathbb{R}_+)$ for $t_0 \leq s \leq t \leq T$ and $\alpha(t) \in C^1(I, I)$ be nondecreasing with $\alpha(t) \leq t$ on I and $k \geq 0$ be a constant.*

(a₁) If

$$(2.1) \quad u(t) \leq k + \int_{\alpha(t_0)}^{\alpha(t)} \left[a(s) u(s) + \int_{\alpha(t_0)}^s b(s, \sigma) u(\sigma) d\sigma \right] ds,$$


for $t \in I$, then

$$(2.2) \quad u(t) \leq k \exp(A(t)),$$

for $t \in I$, where

$$(2.3) \quad A(t) = \int_{\alpha(t_0)}^{\alpha(t)} \left[a(s) + \int_{\alpha(t_0)}^s b(s, \sigma) d\sigma \right] ds,$$

for $t \in I$.



On Some Retarded Integral Inequalities and Applications

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 4 of 16

(a₂) Let $g \in C(\mathbb{R}_+, \mathbb{R}_+)$ be a nondecreasing function with $g(u) > 0$ for $u > 0$.
If

$$(2.4) \quad u(t) \leq k + \int_{\alpha(t_0)}^{\alpha(t)} \left[a(s) g(u(s)) + \int_{\alpha(t_0)}^s b(s, \sigma) g(u(\sigma)) d\sigma \right] ds,$$

for $t \in I$, then for $t_0 \leq t \leq t_1$,

$$(2.5) \quad u(t) \leq G^{-1} [G(k) + A(t)],$$

where $A(t)$ is defined by (2.3), G^{-1} is the inverse function of

$$(2.6) \quad G(r) = \int_{r_0}^r \frac{ds}{g(s)}, \quad r > 0, \quad r_0 > 0,$$

and $t_1 \in I$ is chosen so that


$$G(k) + A(t) \in \text{Dom}(G^{-1}),$$

for all t lying in the interval $[t_0, t_1]$.

Theorem 2.2. Let $u(x, y), a(x, y) \in C(\Delta, \mathbb{R}_+)$, $b(x, y, s, t) \in C(\Delta^2, \mathbb{R}_+)$, for $x_0 \leq s \leq x \leq X$, $y_0 \leq t \leq y \leq Y$, $\alpha(x) \in C^1(J_1, J_1)$, $\beta(y) \in C^1(J_2, J_2)$ be nondecreasing with $\alpha(x) \leq x$ on J_1 , $\beta(y) \leq y$ on J_2 and $k \geq 0$ be a constant.

(b₁) If

$$(2.7) \quad u(x, y) \leq k + \int_{\alpha(x_0)}^{\alpha(x)} \int_{\beta(y_0)}^{\beta(y)} \left[a(s, t) u(s, t) + \int_{\alpha(x_0)}^s \int_{\beta(y_0)}^t b(s, t, \sigma, \eta) u(\sigma, \eta) d\eta d\sigma \right] dt ds,$$



On Some Retarded Integral
Inequalities and Applications

B.G. Pachpatte

Title Page

Contents




Go Back

Close

Quit

Page 5 of 16



Title Page

Contents



Go Back

Close

Quit

Page 6 of 16

for $(x, y) \in \Delta$, then

$$(2.8) \quad u(x, y) \leq k \exp(A(x, y)),$$

for $(x, y) \in \Delta$, where

$$(2.9) \quad A(x, y) = \int_{\alpha(x_0)}^{\alpha(x)} \int_{\beta(y_0)}^{\beta(y)} \left[a(s, t) + \int_{\alpha(x_0)}^s \int_{\beta(y_0)}^t b(s, t, \sigma, \eta) d\eta d\sigma \right] dt ds,$$

for $(x, y) \in \Delta$.

(b₂) Let g be as in Theorem 2.1, part (a₂). If

$$(2.10) \quad u(x, y) \leq k + \int_{\alpha(x_0)}^{\alpha(x)} \int_{\beta(y_0)}^{\beta(y)} \left[a(s, t) g(u(s, t)) + \int_{\alpha(x_0)}^s \int_{\beta(y_0)}^t b(s, t, \sigma, \eta) g(u(\sigma, \eta)) d\eta d\sigma \right] dt ds,$$

for $(x, y) \in \Delta$, then for $x_0 \leq x \leq x_1, y_0 \leq y \leq y_1$,

$$(2.11) \quad u(x, y) \leq G^{-1} [G(k) + A(x, y)],$$

where $A(x, y)$ is defined by (2.9), G, G^{-1} are as defined in Theorem 2.1, part (a₂) and $x_1 \in J_1, y_1 \in J_2$ are chosen so that

$$G(k) + A(x, y) \in \text{Dom}(G^{-1}),$$

for all x and y lying in $[x_0, x_1]$ and $[y_0, y_1]$ respectively.

3. Proofs of Theorems 2.1 and 2.2

From the hypotheses, we observe that $\alpha'(t) \geq 0$ for $t \in I$, $\alpha'(x) \geq 0$ for $x \in J_1$, $\beta'(y) \geq 0$ for $y \in J_2$.

(a₁) Let $k > 0$ and define a function $z(t)$ by the right hand side of (2.1). Then $z(t) > 0$, $z(t_0) = k$, $u(t) \leq z(t)$ and

$$(3.1) \quad z'(t) = \left[a(\alpha(t)) u(\alpha(t)) + \int_{\alpha(t_0)}^{\alpha(t)} [b(\alpha(t), \sigma) u(\sigma) d\sigma] \right] \alpha'(t) \\ \leq \left[a(\alpha(t)) z(\alpha(t)) + \int_{\alpha(t_0)}^{\alpha(t)} [b(\alpha(t), \sigma) z(\sigma) d\sigma] \right] \alpha'(t).$$


From (3.1) it is easy to observe that

$$(3.2) \quad \frac{z'(t)}{z(t)} \leq \left[a(\alpha(t)) + \int_{\alpha(t_0)}^{\alpha(t)} b(\alpha(t), \sigma) d\sigma \right] \alpha'(t).$$

Integrating (3.2) from t_0 to t , $t \in I$ and by making the change of variables yields

$$(3.3) \quad z(t) \leq k \exp(A(t)),$$

for $t \in I$. Using (3.3) in $u(t) \leq z(t)$ we get the inequality in (2.2). If $k \geq 0$, we carry out the above procedure with $k + \varepsilon$ instead of k , where $\varepsilon > 0$ is an arbitrary small constant, and subsequently pass to the limit as $\varepsilon \rightarrow 0$ to obtain (2.2).



On Some Retarded Integral
Inequalities and Applications

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 7 of 16

(a₂) Let $k > 0$ and define a function $z(t)$ by the right hand side of (2.4). Then $z(t) > 0$, $z(t_0) = k$, $u(t) \leq z(t)$ and as in the proof of (a₁) we get

$$(3.4) \quad \frac{z'(t)}{g(z(t))} \leq \left[a(\alpha(t)) + \int_{\alpha(t_0)}^{\alpha(t)} b(\alpha(t), \sigma) d\sigma \right] \alpha'(t).$$

From (2.6) and (3.4) we have

$$(3.5) \quad \begin{aligned} \frac{d}{dt} G(z(t)) \\ = \frac{z'(t)}{g(z(t))} \leq \left[a(\alpha(t)) + \int_{\alpha(t_0)}^{\alpha(t)} b(\alpha(t), \sigma) d\sigma \right] \alpha'(t). \end{aligned}$$


Integrating (3.5) from t_0 to t , $t \in I$ and by making the change of variables we have

$$(3.6) \quad G(z(t)) \leq G(k) + A(t).$$

Since $G^{-1}(z)$ is increasing, from (3.6) we have

$$(3.7) \quad z(t) \leq G^{-1}[G(k) + A(t)].$$

Using (3.7) in $u(t) \leq z(t)$ we get (2.5). The case $k \geq 0$ can be completed as mentioned in the proof of (a₁). The subinterval $t_0 \leq t \leq t_1$ for t is obvious.



On Some Retarded Integral Inequalities and Applications

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 8 of 16

(b₁) Let $k > 0$ and define a function $z(x, y)$ by the right hand side of (2.7). Then $z(x, y) > 0$, $z(x_0, y) = z(x, y_0) = k$, $u(x, y) \leq z(x, y)$ and

$$\begin{aligned}
 (3.8) D_1 z(x, y) &= \left[\int_{\beta(y_0)}^{\beta(y)} \left[a(\alpha(x), t) u(\alpha(x), t) \right. \right. \\
 &\quad \left. \left. + \int_{\alpha(x_0)}^{\alpha(x)} \int_{\beta(y_0)}^t b(\alpha(x), t, \sigma, \eta) u(\sigma, \eta) d\eta d\sigma \right] dt \right] \alpha'(x) \\
 &\leq \left[\int_{\beta(y_0)}^{\beta(y)} \left[a(\alpha(x), t) z(\alpha(x), t) \right. \right. \\
 &\quad \left. \left. + \int_{\alpha(x_0)}^{\alpha(x)} \int_{\beta(y_0)}^t b(\alpha(x), t, \sigma, \eta) z(\sigma, \eta) d\eta d\sigma \right] dt \right] \alpha'(x).
 \end{aligned}$$

From (3.8) it is easy to observe that

$$\begin{aligned}
 (3.9) \quad \frac{D_1 z(x, y)}{z(x, y)} &\leq \left[\int_{\beta(y_0)}^{\beta(y)} \left[a(\alpha(x), t) \right. \right. \\
 &\quad \left. \left. + \int_{\alpha(x_0)}^{\alpha(x)} \int_{\beta(y_0)}^t b(\alpha(x), t, \sigma, \eta) d\eta d\sigma \right] dt \right] \alpha'(x).
 \end{aligned}$$

Keeping y fixed in (3.9), setting $x = \xi$ and integrating it with respect to ξ from x_0 to x and making the change of variables we get

$$(3.10) \quad z(x, y) \leq k \exp(A(x, y)).$$



On Some Retarded Integral Inequalities and Applications

B.G. Pachpatte

Title Page

Contents



Go Back


Close

Quit

Page 9 of 16

Using (3.10) in $u(x, y) \leq z(x, y)$, we get the required inequality in (2.8).
The case $k \geq 0$ follows as mentioned in the proof of (a_1) .

(b_2) The proof can be completed by following the proof of (a_2) and closely looking at the proof of (b_1) . Here we omit the details.



On Some Retarded Integral Inequalities and Applications

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 10 of 16

4. Some Applications

In this section, we present some immediate applications of the inequality (a_1) in Theorem 2.1 to study certain properties of solutions of the integrodifferential equation

$$(P) \quad x'(t) = F \left(t, x(t-h(t)), \int_{t_0}^t f(t, \sigma, x(\sigma-h(\sigma))) d\sigma \right),$$

with the given initial condition

$$(P_0) \quad x(t_0) = x_0,$$

where $f \in C(I^2 \times \mathbb{R}, \mathbb{R})$, $F \in C(I \times \mathbb{R}^2, \mathbb{R})$, x_0 is a real constant and $h \in C^1(I, I)$ be nondecreasing with $t-h(t) \geq 0$, $h'(t) < 1$, $h(t_0) = 0$.

The following theorem deals with the estimate on the solution of $(P) - (P_0)$.


Theorem 4.1. *Suppose that*

$$(4.1) \quad |f(t, s, x)| \leq b(t, s) |x|,$$

$$(4.2) \quad |F(t, x, w)| \leq a(t) |x| + |w|,$$

where $a(t)$, $b(t, s)$ are as defined in Theorem 2.1 and let

$$(4.3) \quad M = \max_{t \in I} \frac{1}{1-h'(t)}.$$



On Some Retarded Integral
Inequalities and Applications

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 11 of 16

If $x(t)$ is any solution of $(P) - (P_0)$, then

$$(4.4) \quad |x(t)| \leq |x_0| \exp \left(\int_{t_0}^{t-h(t)} \left[Ma(s+h(\eta)) + \int_{t_0}^s M^2 b(s+h(\eta), \sigma+h(\tau)) d\sigma \right] ds \right),$$

for t, η, τ in I .

Proof. The solution $x(t)$ of $(P) - (P_0)$ can be written as


$$(4.5) \quad x(t) = x_0 + \int_{t_0}^t F \left(s, x(s-h(s)), \int_{t_0}^s f(s, \sigma, x(\sigma-h(\sigma))) d\sigma \right) ds.$$

Using (4.1) – (4.3) in (4.5) and making the change of variables we have

$$|x(t)| \leq |x_0| + \int_{t_0}^{t-h(t)} \left[Ma(s+h(\eta)) |x(s)| + \int_{t_0}^s M^2 b(s+h(\eta), \sigma+h(\tau)) |x(\sigma)| d\sigma \right] ds,$$

for t, η, τ in I . Now a suitable application of the inequality in (a_1) given in Theorem 2.1 yields the required estimate in (4.4). \square

Next, we shall prove the uniqueness of the solutions of $(P) - (P_0)$.



On Some Retarded Integral Inequalities and Applications

B.G. Pachpatte

Title Page

Contents




Go Back

Close

Quit

Page 12 of 16



Title Page

Contents



Go Back

Close

Quit

Page 13 of 16

Theorem 4.2. Suppose that the functions f, F in (P) satisfy the conditions

$$(4.6) \quad |f(t, s, x) - f(t, s, y)| \leq b(t, s) |x - y|,$$

$$(4.7) \quad |F(t, x, \bar{x}) - F(t, y, \bar{y})| \leq a(t) |x - y| + |\bar{x} - \bar{y}|,$$

where $a(t), b(t, s)$ are as defined in Theorem 2.1 and let M be as in (4.3). Then the problem $(P) - (P_0)$ has at most one solution on I .

Proof. Let $x(t)$ and $\bar{x}(t)$ be two solutions of $(P) - (P_0)$ on I , then we have

$$(4.8) \quad x(t) - \bar{x}(t) = \int_{t_0}^t \left\{ F\left(s, x(s - h(s)), \int_{t_0}^s f(s, \sigma, x(\sigma - h(\sigma))) d\sigma\right) - F\left(s, \bar{x}(s - h(s)), \int_{t_0}^s f(s, \sigma, \bar{x}(\sigma - h(\sigma))) d\sigma\right) \right\} ds.$$

Using (4.6), (4.7) in (4.8) and making the change of variables we have

$$(4.9) \quad |x(t) - \bar{x}(t)| \leq \int_{t_0}^{t-h(t)} \left[Ma(s + h(\eta)) |x(s) - \bar{x}(s)| + \int_{t_0}^s M^2 b(s + h(\eta), \sigma + h(\tau)) |x(\sigma) - \bar{x}(\sigma)| d\sigma \right] ds$$

for t, η, τ in I . A suitable application of the inequality in (a_1) given in Theorem 2.1 yields $|x(t) - \bar{x}(t)| \leq 0$. Therefore $x(t) = \bar{x}(t)$, i.e., there is at most one solution of $(P) - (P_0)$. \square

Our next result shows the dependency of solutions of $(P) - (P_0)$ on initial values.

Theorem 4.3. *Let $x_1(t)$ and $x_2(t)$ be the solutions of (P) with the given initial conditions*

$$(P_1) \quad x_1(t_0) = x_1,$$

and

$$(P_2) \quad x_2(t_0) = x_2,$$


respectively, where x_1, x_2 , are real constants. Suppose that the functions f and F in (P) satisfy the conditions (4.6) and (4.7) in Theorem 4.2 and let M be as in (4.3). Then

$$(4.10) \quad |x_1(t) - x_2(t)| \leq |x_1 - x_2| \exp \left(\int_{t_0}^{t-h(t)} \left[Ma(s+h(\eta)) + \int_{t_0}^s M^2 b(s+h(\eta), \sigma+h(\tau)) d\sigma \right] ds \right),$$

for t, η, τ in I .

Proof. By using the facts that $x_1(t)$ and $x_2(t)$ are the solutions of $(P) - (P_1)$ and $(P) - (P_2)$ respectively, we have

$$(4.11) \quad x_1(t) - x_2(t) = x_1 - x_2 + \int_{t_0}^t \left\{ F \left(s, x_1(s-h(s)), \int_{t_0}^s f(s, \sigma, x_1(\sigma-h(\sigma))) d\sigma \right) - F \left(s, x_2(s-h(s)), \int_{t_0}^s f(s, \sigma, x_2(\sigma-h(\sigma))) d\sigma \right) \right\} ds.$$



On Some Retarded Integral Inequalities and Applications

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 14 of 16

Using (4.6), (4.7) in (4.11) and by making the change of variables, we have

$$(4.12) \quad |x_1(t) - x_2(t)| \leq |x_1 - x_2| + \int_{t_0}^{t-h(t)} \left[Ma(s+h(\eta)) |x_1(s) - x_2(s)| + \int_{t_0}^s M^2 b(s+h(\eta), \sigma+h(\tau)) |x_1(\sigma) - x_2(\sigma)| d\sigma \right] ds,$$

for t, η, τ in I . Now a suitable application of the inequality in (a_1) given in Theorem 2.1 to (4.12) yields the required estimate in (4.10). \square

In concluding we note that the inequality in (b_1) given in Theorem 2.2 can be used to study the similar properties as in Theorems 4.1 – 4.3 for the hyperbolic partial integrodifferential equation

$$(4.13) \quad D_1 D_2 z(x, y) = F(x, y, z(x - h_1(x), y - h_2(y)), Tz(x, y)),$$


with the given initial boundary conditions

$$(4.14) \quad z(x, y_0) = a_1(x), \quad z(x_0, y) = a_2(y), \quad a_1(x_0) = a_2(y_0),$$

where

$$(4.15) \quad Tz(x, y) = \int_{x_0}^x \int_{y_0}^y K(x, y, s, t, z(s - h_1(s), t - h_2(t))) dt ds,$$

under some suitable conditions on the functions involved in (4.13) – (4.15). Since the formulations of these results are very close to those given above, we omit it here. Various other applications of the inequalities given here is left to another work.



Title Page

Contents



Go Back


Close

Quit

Page 15 of 16

References

- [1] D. BAINOV AND P. SIMEONOV, *Integral Inequalities and Applications*, Kluwer Academic Publishers, Dordrecht, 1992.
- [2] R. BELLMAN, The stability of solutions of linear differential equations, *Duke Math. J.*, **10** (1943), 643–647.
- [3] I. BIHARI, A generalization of a lemma of Bellman and its application to uniqueness problems of differential equations, *Acta. Math. Acad. Sci. Hungar.*, **7** (1965), 81–94.
- [4] T.H. GRONWALL, Note on the derivatives with respect to a parameter of the solutions of a system of differential equations, *Ann. Math.*, **20** (1919), 292–296.
- [5] O. LIPOVAN, A retarded Gronwall-like inequality and its applications, *J. Math. Anal. Appl.*, **252** (2000), 389–401.
- [6] B.G. PACHPATTE, *Inequalities for Differential and Integral Equations*, Academic Press, New York, 1998.



On Some Retarded Integral
Inequalities and Applications

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 16 of 16