


## MONOTONICITY RESULTS FOR A COMPOUND QUADRATURE METHOD FOR FINITE-PART INTEGRALS

KAI DIETHELM

Institut Computational Mathematics  
Technische Universität Braunschweig  
Pockelsstraße 14  
38106 Braunschweig, Germany.  
EMail: [k.diethelm@tu-bs.de](mailto:k.diethelm@tu-bs.de)



---

volume 5, issue 2, article 44,  
2004.

*Received 19 May, 2003;  
accepted 30 March, 2004.*

*Communicated by: G. Milovanović*

---

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

## Abstract

Given a function  $f \in C^3[0, 1]$  and some  $q \in (0, 1)$ , we look at the approximation for the Hadamard finite-part integral  $\int_0^1 x^{-q-1} f(x) dx$  based on a piecewise linear interpolant for  $f$  at  $n$  equispaced nodes (i.e., the product trapezoidal rule). The main purpose of this paper is to give sufficient conditions for the sequence of approximations to converge against the correct value of the integral in a monotonic way. An application of the results yields detailed information on the error term of a backward differentiation formula for a fractional differential equation.

**2000 Mathematics Subject Classification:** Primary 41A55; Secondary 65D30, 65L05.  
**Key words:** Hadamard finite-part integral; Quadrature formula; Product trapezoidal formula; Monotonicity; Fractional differential equation; Discrete Gronwall inequality; Backward differentiation formula.

The work described in this paper was partially supported by the U.S. Army Medical Research and Material Command Grant No. DAMD17-01-1-0673 to The Cleveland Clinic to which the author is a co-investigator.

## Contents

1	Introduction	3
2	Main Result	6
3	Further Remarks	16
	References	



---

### Monotonicity Results for a Compound Quadrature Method for Finite-Part Integrals

Kai Diethelm

---

Title Page

Contents



Go Back

Close

Quit

Page 2 of 21

# 1. Introduction


When discussing problems in numerical integration, it is often not sufficient to prove that a certain sequence of approximations is convergent. Frequently one additionally wants to know whether the sequence converges in a monotonic fashion, i.e. whether one can be certain that an approximation using more quadrature nodes is actually better than an approximation with fewer nodes. Such monotonicity results are closely related to the question of finding so-called stopping rules: One needs to determine the value of an integral with a certain prescribed accuracy and the smallest possible amount of work.

For the classical setting when the integral in question is a standard unweighted integral  $\int_a^b f(x)dx$ , this topic is well investigated; we refer to the comprehensive survey of Förster [9] and the references cited therein and to the more recent papers [6, 10, 12] for a description of the present state of the art. However, to the best of our knowledge nothing is known about such results when the functional to be approximated is a weighted strongly singular integral of the form

$$(1.1) \quad I_q[f] := \int_0^1 x^{-q-1} f(x) dx := \sum_{k=0}^{\lfloor q \rfloor} \frac{f^{(k)}(0)}{(k-q)k!} + \int_0^1 x^{-q-1} R_{\lfloor q \rfloor}(x) dx$$

interpreted in Hadamard's finite-part sense (see, e.g., [11] or [2, §1.6.1]). Here we assume  $q$  to be a positive non-integer number, and

$$R_\mu(x) := \frac{1}{\mu!} \int_0^x (x-y)^\mu f^{(\mu+1)}(y) dy$$



---

## Monotonicity Results for a Compound Quadrature Method for Finite-Part Integrals

Kai Diethelm

---

Title Page

Contents



Go Back

Close

Quit

Page 3 of 21


is the remainder of the Taylor polynomial of  $f$ , centered at 0. By  $\lfloor q \rfloor$ , we denote the largest integer not exceeding  $q$ . It is well known that a sufficient condition for the existence of  $I_q[f]$  is that  $f \in C^{\lfloor q \rfloor + 1}[0, 1]$ . Among the most important properties of these integral operators we mention here only that, in contrast to the classical Riemann or Lebesgue integral,  $I_q$  is not a positive functional, i. e. the inequality  $|I_q[f]| \leq I_q[|f|]$  is not true in general. Additional properties are described in [2, §1.6.1]. Since integrals of this type are known to have important applications in various methods for solving partial differential equations or ordinary differential equations of fractional (i.e., non-integer) order [3, 4, 7, 8, 11], we now aim to extend the classical theory to this setting.

Specifically we shall investigate what is probably the most important example of a quadrature formula for  $I_q$ , the product trapezoidal method. The construction of the method is simple: Given an integer  $n$ , we divide the fundamental integral  $[0, 1]$  into  $n$  subintervals of equal length with break points  $x_j = \frac{j}{n}$ ,  $j = 0, 1, \dots, n$ . We then replace the function  $f$  by its piecewise linear interpolant (linear interpolating spline) with knots and nodes at  $x_0, x_1, \dots, x_n$ . Denoting this interpolant by  $f_{n+1}$  (the subscript  $n + 1$  being the number of interpolation points), we then define our approximation  $I_{q,n+1}$  for  $I_q$  according to

$$I_{q,n+1}[f] := I_q[f_{n+1}],$$

where we note that the piecewise linear structure of  $f_{n+1}$  allows us to calculate the expression on the right-hand side effectively.

An explicit representation for  $I_{q,n+1}$  is available from [3, Lemma 2.1]:




---

**Monotonicity Results for a  
Compound Quadrature Method  
for Finite-Part Integrals**

Kai Diethelm

---

Title Page

Contents



Go Back

Close

Quit

Page 4 of 21

**Lemma 1.1.** *We have*


$$I_{q,n+1}[f] = \sum_{k=0}^n \alpha_{kn} f\left(\frac{k}{n}\right),$$

where

$$= \begin{cases} q(1-q)n^{-q}\alpha_{kn} & \\ -1 & \text{for } k = 0, \\ 2k^{1-q} - (k-1)^{1-q} - (k+1)^{1-q} & \text{for } k = 1, 2, \dots, n-1, \\ (q-1)k^{-q} - (k-1)^{1-q} + k^{1-q} & \text{for } k = n. \end{cases}$$

There are various reasons for choosing this formula as a first candidate for our investigations:

- It is a generalization of the classical trapezoidal formula, which is in turn the quadrature formula for standard integrals that was historically among the first and is very thoroughly investigated with respect to its monotonicity properties.
- Many other properties of this formula have been studied in great detail, see, e.g., [4, 5].
- It has been used very successfully as the basic ingredient for algorithms for the numerical solution of fractional differential equations [3].




---

**Monotonicity Results for a Compound Quadrature Method for Finite-Part Integrals**

Kai Diethelm

---

Title Page

Contents



Go Back

Close

Quit

Page 5 of 21

## 2. Main Result

The main result of this paper is the following monotonicity theorem that directly corresponds to an analogous result for standard integrals (see, e.g., [13] or [1, Thm. 105]).

**Theorem 2.1.** *Let  $0 < q < 1$  be fixed, and let  $f \in C^3[0, 1]$ . Moreover assume that  $f''$  is nonnegative on  $[0, 1]$  (i.e.  $f$  is convex) and  $f'''$  is nonpositive on  $[0, 1]$ . Then, the sequence  $(I_{q,n+1}[f])_{n=1}^{\infty}$  is monotonically decreasing, and its limit is  $I_q[f]$ .*

For the proof we shall use some properties of the quadrature rule  $(I_{q,n+1})_{n=1}^{\infty}$  that have been established previously. Here and in the following we will make use of the notation

$$R_{n+1} := I_q - I_{q,n+1}$$

to denote the remainder functional of  $I_{q,n+1}$ . For the sake of simplicity we have suppressed the dependence on  $q$  in our notation. (Remember that  $q$  is assumed to be fixed.)

In view of the above mentioned properties of the functional  $I_q$  and its approximation  $I_{q,n+1}$ , we may apply the classical Peano kernel theorem [16] to  $R_{n+1}$  and derive

**Lemma 2.2.** *Let  $0 < q < 1$  or  $1 < q < 2$ , and assume that  $f \in C^2[0, 1]$ . Then,*

$$R_{n+1}[f] = \int_0^1 K_2(R_{n+1}, x) f''(x) dx,$$

where  $K_2(R_{n+1}, \cdot)$  is the second Peano kernel of  $R_{n+1}$ , given by

$$K_2(R_{n+1}, x) := R_{n+1}[(\cdot - x)_+].$$



---

### Monotonicity Results for a Compound Quadrature Method for Finite-Part Integrals

Kai Diethelm

---

Title Page

Contents



Go Back

Close

Quit

Page 6 of 21

Here  $(\cdot)_+$  is the truncated power function defined by

$$(x)_+ := \begin{cases} x & \text{if } x \geq 0, \\ 0 & \text{otherwise.} \end{cases}$$

From Lemma 2.2 we can deduce the explicit representation

$$(2.1) \quad K_2(R_{n+1}, t) = \sum_{k=0}^j \alpha_{kn} \binom{k}{n} - t \quad - \frac{t^{1-q}}{q(1-q)} \quad \text{for } t \in \left[ \frac{j}{n}, \frac{j+1}{n} \right]$$

of the Peano kernel in a straightforward way (as is done, e.g., in [1, Thm. 16] for classical quadrature formulas).


In [4, p. 487] it has been stated that  $R_{n+1}$  is negative definite of order two whenever  $0 < q < 1$  or  $1 < q < 2$ . Unfortunately this result is incorrect; it should read as follows.

**Lemma 2.3.** *For any  $n \geq 2$ , the functional  $R_{n+1}$  is negative definite for  $0 < q < 1$  and indefinite for  $1 < q < 2$ .*

*Proof.* It is clear that

$$\begin{aligned} R_{n+1}[f] &= I_q[f - f_{n+1}] \\ &= \int_0^{n-1} u^{-q-1}(f(u) - f_{n+1}(u))du + \int_{n-1}^1 u^{-q-1}(f(u) - f_{n+1}(u))du. \end{aligned}$$

To prove the negative definiteness in the case  $0 < q < 1$  it is sufficient to show that  $R_{n+1}[f] \leq 0$  whenever  $f$  is convex. Thus we assume  $f$  to be convex.




---

**Monotonicity Results for a Compound Quadrature Method for Finite-Part Integrals**

Kai Diethelm

---

Title Page

Contents



Go Back

Close

Quit

Page 7 of 21

Then, as is well known,  $f(u) \leq f_{n+1}(u)$  for all  $u$ , and hence the second integral is nonpositive. Moreover, for the first integral we can explicitly calculate the Peano kernel representation

$$\int_0^{n^{-1}} u^{-q-1}(f(u) - f_{n+1}(u))du =: A[f] = \int_0^{n^{-1}} f''(u)K_2(A, u)du.$$

In view of the relation between the functionals  $A$  and  $R_n$ , it is evident that for  $u \in [0, n^{-1}]$  we have


$$(2.2) \quad K_2(A, u) = K_2(R_{n+1}, u) = -\frac{u}{q(1-q)}(u^{-q} - n^q) \leq 0$$

because of (2.1). Thus, the first integral is nonpositive too if  $f$  is convex, and the claim follows.

The indefiniteness in the case  $1 < q < 2$  follows by very similar arguments. We find the same expression for the Peano kernel  $K_2(A, \cdot)$  as above, but now the sign of  $(1 - q)$  and hence the sign of the complete expression has changed. Thus  $K_2(A, u) \geq 0$  for  $0 < u < n^{-1}$ . Since nothing changes in the integral over  $[n^{-1}, 1]$ , we deduce that now  $R_{n+1}$  is the sum of a positive definite functional on  $C^2[0, n^{-1}]$  and a negative definite functional on  $C^2[n^{-1}, 1]$ , and hence it must be indefinite.  $\square$

Formulated in terms of Peano kernels, the case  $0 < q < 1$  of Lemma 2.3 can be restated as:

**Lemma 2.4.** *Let  $0 < q < 1$ ,  $x \in [0, 1]$  and  $n \in \mathbb{N}$ . Then,  $K_2(R_{n+1}, x) \leq 0$ .*




---

**Monotonicity Results for a  
Compound Quadrature Method  
for Finite-Part Integrals**

Kai Diethelm

---

Title Page

Contents



Go Back

Close

Quit

Page 8 of 21



Standard methods from elementary Peano kernel theory give us additional fundamental results on the function  $K_2(R_{n+1}, \cdot)$  and its  $L_1$  norm

$$\rho_{n+1} := \|K_2(R_{n+1}, \cdot)\|_1 = \int_0^1 |K_2(R_{n+1}, x)| dx;$$

we omit the details of the proof.

**Lemma 2.5.** *Let  $0 < q < 1$ .*

(a) *For  $j = 0, 1, \dots, n$  we have  $K_2(R_{n+1}, x_j) = 0$ .*

(b) *The sequence  $(\rho_{n+1})_{n=1}^\infty$  is monotonically decreasing.*

Finally we quote another result on the sequence mentioned in Lemma 2.5 (b) from [5, Thm. 1.2]; more details are given there and in [4, Thm. 2.3].

**Lemma 2.6.** *For  $0 < q < 1$  there exists some constant  $c_q$  such that  $\rho_{n+1} = c_q n^{q-2} + O(n^{-2})$ .*

We are now in a position to prove our main result.


*Proof of Theorem 2.1.* First we note that, by Lemmas 2.2 and 2.4,

$$R_{n+1}[f] = \int_0^1 K_2(R_{n+1}, x) f''(x) dx \leq 0,$$

and hence by definition of  $R_{n+1}$  we find that  $I_{q,n+1}[f] \geq I_q[f]$ .

Moreover, by Lemma 2.2, Hölder's inequality and Lemma 2.6,

$$|R_{n+1}[f]| = \left| \int_0^1 K_2(R_{n+1}, x) f''(x) dx \right| \leq \|f''\|_\infty \cdot \rho_{n+1} \rightarrow 0,$$




---

**Monotonicity Results for a  
Compound Quadrature Method  
for Finite-Part Integrals**

Kai Diethelm

---

Title Page

Contents



Go Back

Close

Quit

Page 9 of 21

i.e. (again by definition of  $R_{n+1}$ ),  $I_{q,n+1}[f] \rightarrow I_q[f]$  as  $n \rightarrow \infty$ .

It remains to prove that the sequence  $(I_{q,n+1}[f])$  decreases monotonically or, equivalently, that the sequence  $(R_{n+1}[f])$  increases monotonically. To this end, we use the representation of  $R_{n+1}[f]$  from Lemma 2.2 and introduce the functions  $J_{n+1}$  and  $L_{n+1}$  according to

$$J_{n+1}(x) := K_2(R_{n+1}, x) + \rho_{n+1} \quad \text{and} \quad L_{n+1}(x) := \int_0^x J_{n+1}(t) dt.$$

Then, a partial integration yields


$$\begin{aligned} R_{n+1}[f] &= \int_0^1 (J_{n+1}(x) - \rho_{n+1}) f''(x) dx \\ &= f''(x) [L_{n+1}(x) - x\rho_{n+1}]_0^1 - \int_0^1 f'''(x) [L_{n+1}(x) - x\rho_{n+1}] dx \\ &= f''(1) [L_{n+1}(1) - \rho_{n+1}] - \int_0^1 f'''(x) [L_{n+1}(x) - x\rho_{n+1}] dx \end{aligned}$$

since obviously  $L_{n+1}(0) = 0$ . Moreover,

$$\begin{aligned} L_{n+1}(1) &= \int_0^1 J_{n+1}(t) dt \\ &= \int_0^1 (K_2(R_{n+1}, x) + \rho_{n+1}) dx = \int_0^1 K_2(R_{n+1}, x) dx + \rho_{n+1}. \end{aligned}$$

Recalling the definition of  $\rho_{n+1}$  and the nonpositivity of  $K_2(R_{n+1}, \cdot)$  (see Lemma 2.4), we find

$$\rho_{n+1} = \int_0^1 |K_2(R_{n+1}, x)| dx = - \int_0^1 K_2(R_{n+1}, x) dx,$$



## Monotonicity Results for a Compound Quadrature Method for Finite-Part Integrals

Kai Diethelm

Title Page

Contents



Go Back

Close

Quit

Page 10 of 21

and hence  $L_{n+1}(1) = 0$  too. Combining these results we find

$$R_{n+1}[f] = -\rho_{n+1}f''(1) - \int_0^1 f'''(x) [L_{n+1}(x) - x\rho_{n+1}] dx.$$

Under our assumptions on  $f$ , we know that  $f''(1) \geq 0$ , and hence by Lemma 2.5 (b) we see that the first expression on the right-hand side, viz. the quantity  $-\rho_{n+1}f''(1)$ , is indeed a monotonically increasing function of  $n$ . It thus remains to prove that the remaining term has got this property as well. Since  $f'''$  is assumed to be negative, it is sufficient for this purpose to show that, for every fixed  $x \in [0, 1]$ , the function  $\phi_x$  defined by

$$\phi_x(n+1) := L_{n+1}(x) - x\rho_{n+1}$$

is a non-decreasing function of  $n$ . Note that

$$\phi_x(n+1) = \int_0^x (K_2(R_{n+1}, t) + \rho_{n+1}) dt - x\rho_{n+1} = \int_0^x K_2(R_{n+1}, t) dt.$$


For the proof of the monotonicity of  $\phi_x$  we distinguish two cases.

First we look at  $0 \leq x \leq (n+1)^{-1}$ . An explicit representation for  $K_2(R_{n+1}, t)$  in the case  $0 < t < n^{-1}$  can be taken from eq. (2.1); it reads

$$K_2(R_{n+1}, t) = \frac{t}{q(1-q)} (n^q - t^{-q}).$$

Consequently,

$$\phi_x(n+1) = \frac{1}{q(1-q)} \left( \frac{1}{2}n^q x^2 - \frac{1}{2-q} x^{2-q} \right)$$



### Monotonicity Results for a Compound Quadrature Method for Finite-Part Integrals

Kai Diethelm

Title Page

Contents



Go Back

Close

Quit

Page 11 of 21

for  $0 \leq x \leq n^{-1}$ . In an analogous manner we find

$$\phi_x(n+2) = \frac{1}{q(1-q)} \left( \frac{1}{2}(n+1)^q x^2 - \frac{1}{2-q} x^{2-q} \right)$$

for  $0 \leq x \leq (n+1)^{-1}$ . From these identities we immediately see

$$\phi_x(n+1) \leq \phi_x(n+2)$$

for all  $n$  and  $0 \leq x \leq (n+1)^{-1}$  as required.

In the second case  $(n+1)^{-1} < x \leq n^{-1}$  we will prove the relation


$$\phi_x(n+1) \leq \phi_{(n+1)^{-1}}(n+1) \leq \phi_{n^{-1}}(n+2) \leq \phi_x(n+2).$$

Since  $K_2(R_{n+1}, \cdot)$  is a nonpositive function (see Lemma 2.4), we find that  $\phi_x(n+1)$  is a decreasing function of  $x$ . Thus the first and the last of the three inequalities above are evident. It remains to show the middle one. To this end we note that we still have, as above,

$$\phi_x(n+1) = \frac{1}{q(1-q)} \left( \frac{1}{2} n^q x^2 - \frac{1}{2-q} x^{2-q} \right),$$

and therefore

$$\begin{aligned} \phi_{(n+1)^{-1}}(n+1) &= \frac{1}{q(1-q)} \left( \frac{1}{2} n^q (n+1)^{-2} - \frac{1}{2-q} (n+1)^{q-2} \right) \\ &= \frac{1}{q(1-q)(n+1)^2} \left( \frac{1}{2} n^q - \frac{1}{2-q} (n+1)^q \right). \end{aligned}$$




---

**Monotonicity Results for a  
Compound Quadrature Method  
for Finite-Part Integrals**

Kai Diethelm

---

Title Page

Contents



Go Back

Close

Quit


Page 12 of 21

However we now pass a node of the formula  $I_{q,n+2}$ , namely the point  $(n+1)^{-1}$ , and hence the Peano kernel  $K_2$  of this formula becomes

$$\begin{aligned} K_2(R_{n+2}, t) &= -\alpha_{0,n+1}t + \alpha_{1,n+1} \left( \frac{1}{n+1} - t \right) - \frac{t^{1-q}}{q(1-q)} \\ &= \frac{(n+1)^q}{q(1-q)} \left( t + (2 - 2^{1-q}) \left( \frac{1}{n+1} - t \right) - (n+1)^{-q}t^{1-q} \right) \end{aligned}$$

according to eq. (2.1) and Lemma 1.1. Thus we have

$$\begin{aligned} \phi_{n-1}(n+2) &= \int_0^{n-1} K_2(R_{n+2}, t) dt \\ &= \int_0^{(n+1)^{-1}} K_2(R_{n+2}, t) dt + \int_{(n+1)^{-1}}^{n-1} K_2(R_{n+2}, t) dt \\ &= \phi_{(n+1)^{-1}}(n+2) + \int_{(n+1)^{-1}}^{n-1} K_2(R_{n+2}, t) dt \\ &= \frac{1}{q(1-q)} \left( \frac{1}{2} - \frac{1}{2-q} \right) (n+1)^{q-2} + \int_{(n+1)^{-1}}^{n-1} K_2(R_{n+2}, t) dt \\ &= -\frac{1}{2(1-q)(2-q)} (n+1)^{q-2} + \int_{(n+1)^{-1}}^{n-1} K_2(R_{n+2}, t) dt \\ &= -\frac{1}{2(1-q)(2-q)} (n+1)^{q-2} \\ &\quad + \frac{1}{2q(1-q)(2-q)} (2(n+1)^{q-2} - 2n^{q-2} \\ &\quad \quad + n^{-2}(n+1)^{q-2}(q-2)(1-2^{1-q}-2n)), \end{aligned}$$



**Monotonicity Results for a  
Compound Quadrature Method  
for Finite-Part Integrals**

Kai Diethelm

Title Page

Contents



Go Back

Close

Quit

Page 13 of 21

and after a rather long but simple calculation we obtain the required inequality.

On the remaining part of the interval  $[0, 1]$ , the required representation of the Peano kernel is also given in (2.1). For the purpose of illustration we have plotted the graphs for  $\phi_x(n+1)$  versus  $x$  in Figure 1 for the special case  $q = 0.3$  and  $n \in \{5, 6, 7\}$ . In a qualitative sense these graphs can be considered to be typical also for other values of  $q \in (0, 1)$ . Using these representations, we can deduce the required property after a lengthy but straightforward calculation on these intervals too.  $\square$



---

**Monotonicity Results for a  
Compound Quadrature Method  
for Finite-Part Integrals**

Kai Diethelm

---

Title Page

Contents




Go Back

Close

Quit

Page 14 of 21



---

**Monotonicity Results for a  
Compound Quadrature Method  
for Finite-Part Integrals**

Kai Diethelm

---

Title Page

Contents



Go Back

Close

Quit

Page 15 of 21




Figure 1: Plots of  $\phi_x(n+1)$  versus  $x$  for  $n = 5$  (dotted line; bottom),  $n = 6$  (dashed line; middle), and  $n = 7$  (solid line; top), over the entire interval  $x \in [0, 1]$  (upper graph) and zoom over the subinterval  $x \in [0, 0.1]$  (lower graph).

### 3. Further Remarks

In Lemma 2.3 we had pointed out a mistake in the discussion of the case  $1 < q < 2$  in our earlier paper [4]. This observation leads to some consequences.

To begin with, an error estimate for the quadrature rule considered above has been discussed in [4, Thm. 2.3]. The analysis there was partly based on the incorrect result and needs to be modified slightly. The correction due to this problem affects only the case  $1 < q < 2$  (the parameter  $p$  used in that paper corresponds to  $q+1$  here), but since the original proof regrettably also contained some typographical errors in the case  $0 < q < 1$ , we give the correct result in both cases here.

**Theorem 3.1.** *Let  $0 < q < 1$  or  $1 < q < 2$ . Then, for every  $n \in \mathbb{N}$  we have*

$$\begin{aligned} \frac{30 - q(q+1)}{360q \cdot |1 - q|(2 - q)} n^{q-2} + \left( \frac{q+1}{180} - \frac{1}{6q} \right) n^{-2} \\ < \rho_{n+1} < \left( \frac{1}{6q} + \frac{1}{2|1 - q|(2 - q)} \right) n^{q-2} - \frac{1}{6q} n^{-2}. \end{aligned}$$

*Proof.* In [4, Proof of Thm. 2.3], we have seen that


$$\rho_{n+1} = \sigma + \tau,$$

where

$$\frac{1 - n^{-q}}{q} \left( \frac{1}{6} - \frac{q(q+1)}{180} \right) n^{q-2} < \sigma < \frac{1 - n^{-q}}{6q} n^{q-2}$$

and

$$\tau = \sup_{\|f''\|_{\infty} \leq 1} |A[f]|,$$



Monotonicity Results for a  
Compound Quadrature Method  
for Finite-Part Integrals

Kai Diethelm

Title Page

Contents



Go Back

Close

Quit

Page 16 of 21



where  $A[f]$  is as in the proof of Lemma 2.3. Thus, standard Peano kernel theory reveals that

$$\tau = \int_0^{n-1} |K_2(A, u)| du.$$

From the explicit representation of  $K_2(A, \cdot)$  in eq. (2.2) we find that

$$\tau = \frac{1}{q \cdot |1 - q|} \int_0^{n-1} u(u^{-q} - n^q) du = \frac{n^{q-2}}{2|1 - q|(2 - q)},$$

and the claim follows.  $\square$


Another aspect of the results presented in §2 is related to the fact that finite-part integrals are a convenient means to represent derivatives of fractional order. To be precise, as is well known [7], we have that the Caputo-type fractional differential operator  $D_*^q$  can be rewritten as

$$D_*^q y(x) = \frac{1}{\Gamma(-q)} \int_0^x (y(u) - y(0))(x - u)^{-q-1} du$$

for  $0 < q < 1$  and as

$$D_*^q y(x) = \frac{1}{\Gamma(-q)} \int_0^x (y(u) - y(0) - uy'(0))(x - u)^{-q-1} du$$

for  $1 < q < 2$ . We refer to the books of Podlubny [14] or Samko et al. [15] for detailed information on fractional derivatives and fractional differential equations; here we only note that our results above can be applied in a direct way to derive monotonically convergent approximations for such derivatives. We recall




---

**Monotonicity Results for a  
Compound Quadrature Method  
for Finite-Part Integrals**

Kai Diethelm

---

Title Page

Contents



Go Back

Close

Quit

Page 17 of 21

that certain other important properties of the approximation method investigated here have been given in [5].

Differential equations involving such operators have proven to be an important tool in many applications in physics, engineering, finance, etc.; see, e.g., the examples mentioned in [14] and the references cited therein. It is an obvious consequence of the above considerations that we may also use the product trapezoidal method for finite-part integrals as a means to construct numerical solutions for fractional differential equations. First results on this topic have been given in [3, 5]. In the analysis of the algorithm, a discrete Gronwall inequality [3, Lemma 2.3] turned out to be helpful. In view of our new developments above we may now strengthen this result and bring it into the form of a two-sided inequality:

**Theorem 3.2.** For  $0 < q < 1$ , let the sequence  $(d_j)$  be given by  $d_1 = 1$  and


$$d_j = 1 + q(1 - q)j^{-q} \sum_{k=1}^{j-1} \alpha_{kj} d_{j-k}, \quad j = 2, 3, \dots,$$

where  $\alpha_{kj}$  is as in Lemma 1.1. Then,

$$j^q \leq d_j \leq \frac{\sin \pi q}{\pi q(1 - q)} j^q, \quad j = 1, 2, 3, \dots$$

We can thus see that the upper bound gives the correct rate of growth of the sequence  $(d_j)$ .

*Proof.* The upper bound is known [3, Lemma 2.3]. For the lower bound, we proceed inductively. The induction basis ( $j = 1$ ) is presupposed. For the induction step we use the fact that  $\alpha_{kj} > 0$  for all  $j$  and  $k$  under consideration and




---

**Monotonicity Results for a  
Compound Quadrature Method  
for Finite-Part Integrals**

Kai Diethelm

---

Title Page

Contents



Go Back

Close

Quit

Page 18 of 21

find, using the function  $\phi(x) = (1 - x)^q$ , that


$$\begin{aligned} d_{j+1} &= 1 + q(1 - q)(j + 1)^{-q} \sum_{k=1}^j \alpha_{k,j+1} d_{j+1-k} \\ &\geq 1 + q(1 - q) \sum_{k=1}^{j+1} \alpha_{k,j+1} \left( \frac{j + 1 - k}{j + 1} \right)^q \\ &= 1 + q(1 - q) (I_{q,j+2}[\phi] - \alpha_{0,j+1} \phi(0)) \\ &= 1 + q(1 - q) I_{q,j+2}[\phi] + (j + 1)^q. \end{aligned}$$

It thus remains to prove that  $q(1 - q)I_{q,j+2}[\phi] \geq -1$ . In view of the fact that  $\phi''(x) < 0$  and  $\phi'''(x) \geq 0$  for  $0 < x < 1$ , Theorem 2.1 allows us to conclude that it is sufficient for this purpose to show that  $q(1 - q)I_{q,2}[\phi] \geq -1$ . An explicit calculation reveals that indeed

$$q(1 - q)I_{q,2}[\phi] = q(1 - q) (\alpha_{02} + \alpha_{12}2^{-q}) = -2^q + 2 - 2^{1-q} \geq -1.$$

This completes the proof.  $\square$

It is our belief that the new lower bound may be useful in gaining an even better understanding of the properties of the differential equation solver; in particular we hope to prove that the error bound derived in [3, Thm. 1.1] is not improvable. But this will be the topic of a different paper.




---

**Monotonicity Results for a  
Compound Quadrature Method  
for Finite-Part Integrals**

Kai Diethelm

---

Title Page

Contents



Go Back

Close

Quit

Page 19 of 21

## References

- [1] H. BRASS, *Quadraturverfahren*, Vandenhoeck & Ruprecht, Göttingen, 1977.
- [2] P.J. DAVIS AND P. RABINOWITZ, *Methods of Numerical Integration*, Academic Press, Orlando, 2nd ed., 1984.
- [3] K. DIETHELM, An algorithm for the numerical solution of differential equations of fractional order, *Elec. Transact. Numer. Anal.*, **5** (1997), 1–6.
- [4] K. DIETHELM, Generalized compound quadrature formulae for finite-part integrals, *IMA J. Numer. Anal.*, **17** (1997), 479–493.
- [5] K. DIETHELM AND G. WALZ, Numerical solution of fractional order differential equations by extrapolation, *Numer. Algorithms*, **16** (1997), 231–253.
- [6] S. EHRICH, Stopping functionals for Gaussian quadrature formulas, *J. Comput. Appl. Math.*, **127** (2001), 153–171.
- [7] D. ELLIOTT, An asymptotic analysis of two algorithms for certain Hadamard finite-part integrals, *IMA J. Numer. Anal.*, **13** (1993), 445–462.
- [8] D. ELLIOTT, Three algorithms for Hadamard finite-part integrals and fractional derivatives, *J. Comput. Appl. Math.*, **62** (1995), 267–283.
- [9] K.-J. FÖRSTER, A survey of stopping rules in quadrature based on Peano kernel methods, *Rend. Circ. Mat. Palermo* (2) Suppl., **33** (1993), 311–330.



---

Monotonicity Results for a  
Compound Quadrature Method  
for Finite-Part Integrals

Kai Diethelm

---

Title Page

Contents




Go Back

Close

Quit

Page 20 of 21

- [10] K.-J. FÖRSTER, P. KÖHLER AND G. NIKOLOV, Monotonicity and stopping rules for compound Gauss-type quadrature formulae, *East J. Approx.*, **4** (1998), 55–74.
- [11] J. HADAMARD, *Lectures on Cauchy's Problem in Linear Partial Differential Equations*, Dover Publ., New York, 1952. Reprint.
- [12] P. KÖHLER, A note on definiteness and monotonicity of quadrature formulae, *Z. Angew. Math. Mech.*, **75** (1995), S645–S646.
- [13] D.J. NEWMAN, Monotonicity of quadrature approximations, *Proc. Amer. Math. Soc.*, **42** (1974), 251–257.
- [14] I. PODLUBNY, *Fractional Differential Equations*, Academic Press, San Diego, 1999.
- [15] S.G. SAMKO, A.A. KILBAS AND O.I. MARICHEV, *Fractional Integrals and Derivatives: Theory and Applications*, Gordon and Breach, Yverdon, 1993.
- [16] A. SARD, Integral representations of remainders, *Duke Math. J.*, **15** (1948), 333–345.




---

**Monotonicity Results for a  
Compound Quadrature Method  
for Finite-Part Integrals**

Kai Diethelm

---

Title Page

Contents



Go Back

Close

Quit

Page 21 of 21