


# Journal of Inequalities in Pure and Applied Mathematics

## ON GRÜSS TYPE INTEGRAL INEQUALITIES

B.G. PACHPATTE

57, Shri Niketen Colony  
Aurangabad - 431 001,  
(Maharashtra) India.  
EMail: [bgpachpatte@hotmail.com](mailto:bgpachpatte@hotmail.com)

©2000 Victoria University  
ISSN (electronic): 1443-5756  
070-01



---

volume 3, issue 1, article 11,  
2002.

*Received 29 June, 2001;  
accepted 18 October, 2001.*

*Communicated by: J. Sándor*

---

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

## Abstract


In this paper we establish two new integral inequalities similar to that of the Grüss inequality by using a fairly elementary analysis.

*2000 Mathematics Subject Classification:* 26D15, 26D20.

*Key words:* Grüss type, Integral Inequalities, Absolutely Continuous, Integral Identity.

## Contents

1	Introduction .....	3
2	Statement of Results .....	4
3	Proof of Theorem 2.1 .....	7
4	Proof of Theorem 2.2 .....	10
	References	



---

### On Grüss Type Integral Inequalities

B.G. Pachpatte

---

Title Page

Contents



Go Back

Close

Quit

Page 2 of 12

# 1. Introduction

In 1935 (see [4, p. 296]), G. Grüss proved the following integral inequality which gives an estimation for the integral of a product in terms of the product of integrals:


$$\left| \frac{1}{b-a} \int_a^b f(x) g(x) dx - \frac{1}{b-a} \int_a^b f(x) dx \cdot \frac{1}{b-a} \int_a^b g(x) dx \right| \leq \frac{1}{4} (M - m) (N - n),$$

provided that  $f$  and  $g$  are two integrable functions on  $[a, b]$  and satisfying the condition

$$m \leq f(x) \leq M, \quad n \leq g(x) \leq N,$$

for all  $x \in [a, b]$ , where  $m, M, n, N$  are given real constants.

A great deal of attention has been given to the above inequality and many papers dealing with various generalizations, extensions and variants have appeared in the literature, see [1] – [6] and the references cited therein. The main purpose of the present paper is to establish two new integral inequalities similar to that of the Grüss inequality involving functions and their higher order derivatives. The analysis used in the proof is elementary and our results provide new estimates on inequalities of this type.



On Grüss Type Integral Inequalities

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 3 of 12

## 2. Statement of Results

In this section, we state our results to be proved in this paper. In what follows, we denote by  $\mathbb{R}$ , the set of real numbers and  $[a, b] \subset \mathbb{R}$ ,  $a < b$ .

Our main results are given in the following theorems.

**Theorem 2.1.** *Let  $f, g : [a, b] \rightarrow \mathbb{R}$  be functions such that  $f^{(n-1)}, g^{(n-1)}$  are absolutely continuous on  $[a, b]$  and  $f^{(n)}, g^{(n)} \in L_\infty[a, b]$ . Then*


$$(2.1) \quad \left| \frac{1}{b-a} \int_a^b f(x)g(x)dx - \left( \frac{1}{b-a} \int_a^b f(x)dx \right) \left( \frac{1}{b-a} \int_a^b g(x)dx \right) - \frac{1}{2(b-a)^2} \int_a^b \left[ \left( \sum_{k=1}^{n-1} F_k(x) \right) g(x) + \left( \sum_{k=1}^{n-1} G_k(x) \right) f(x) \right] dx \right| \leq \frac{1}{2(b-a)^2} \int_a^b (|g(x)| \|f^{(n)}\|_\infty + |f(x)| \|g^{(n)}\|_\infty) A_n(x) dx,$$

where

$$(2.2) \quad F_k(x) = \left[ \frac{(b-x)^{k+1} + (-1)^k(x-a)^{k+1}}{(k+1)!} \right] f^{(k)}(x),$$

$$(2.3) \quad G_k(x) = \left[ \frac{(b-x)^{k+1} + (-1)^k(x-a)^{k+1}}{(k+1)!} \right] g^{(k)}(x),$$

$$(2.4) \quad A_n(x) = \int_a^b |K_n(x, t)| dt,$$



On Grüss Type Integral Inequalities

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 4 of 12

in which  $K_n : [a, b]^2 \rightarrow \mathbb{R}$  is given by

$$(2.5) \quad K_n(x, t) = \begin{cases} \frac{(t-a)^n}{n!} & \text{if } t \in [a, x] \\ \frac{(t-b)^n}{n!} & \text{if } t \in (x, b] \end{cases}$$

and


$$\|f^{(n)}\|_\infty = \sup_{t \in [a, b]} |f^{(n)}(t)| < \infty,$$

$$\|g^{(n)}\|_\infty = \sup_{t \in [a, b]} |g^{(n)}(t)| < \infty,$$

for  $x \in [a, b]$  and  $n \geq 1$  a natural number.

**Theorem 2.2.** Let  $p, q : [a, b] \rightarrow \mathbb{R}$  be functions such that  $p^{(n-1)}, q^{(n-1)}$  are absolutely continuous on  $[a, b]$  and  $p^{(n)}, q^{(n)} \in L_\infty[a, b]$ . Then

$$(2.6) \quad \left| \frac{1}{b-a} \int_a^b p(x) q(x) dx - n \left( \frac{1}{b-a} \int_a^b p(x) dx \right) \left( \frac{1}{b-a} \int_a^b q(x) dx \right) + \frac{1}{2(b-a)} \int_a^b \left[ \left( \sum_{k=1}^{n-1} P_k(x) \right) q(x) + \left( \sum_{k=1}^{n-1} Q_k(x) \right) p(x) \right] dx \right| \leq \frac{1}{2(n-1)!(b-a)^2} \int_a^b (|q(x)| \|p^{(n)}\|_\infty + |p(x)| \|q^{(n)}\|_\infty) B_n(x) dx,$$



On Grüss Type Integral Inequalities

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 5 of 12

where

$$(2.7) \quad P_k(x) = \frac{(n-k)}{k!} \cdot \frac{p^{(k-1)}(a)(x-a)^k - p^{(k-1)}(b)(x-b)^k}{b-a},$$

$$(2.8) \quad Q_k(x) = \frac{(n-k)}{k!} \cdot \frac{q^{(k-1)}(a)(x-a)^k - q^{(k-1)}(b)(x-b)^k}{b-a},$$

$$(2.9) \quad B_n(x) = \int_a^b |(x-t)^{n-1} r(t,x)| dt,$$

in which


$$r(t,x) = \begin{cases} t-a, & \text{if } a \leq t \leq x \leq b, \\ t-b, & \text{if } a \leq x < t \leq b, \end{cases}$$

and

$$\|p^{(n)}\|_\infty = \sup_{t \in [a,b]} |p^{(n)}(t)| < \infty,$$

$$\|q^{(n)}\|_\infty = \sup_{t \in [a,b]} |q^{(n)}(t)| < \infty,$$

for  $x \in [a, b]$  and  $n \geq 1$  is a natural number.



### On Grüss Type Integral Inequalities

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 6 of 12

### 3. Proof of Theorem 2.1

From the hypotheses on  $f$ , we have the following integral identity (see [1, p. 52]):


$$(3.1) \quad \int_a^b f(t) dt = \sum_{k=0}^{n-1} \left[ \frac{(b-x)^{k+1} + (-1)^k (x-a)^{k+1}}{(k+1)!} \right] f^{(k)}(x) \\ + (-1)^n \int_a^b K_n(x, t) f^{(n)}(t) dt,$$

for  $x \in [a, b]$ . In [1] the identity (3.1) is proved by mathematical induction. For a different proof, see [6]. The identity (3.1) can be rewritten as

$$(3.2) \quad f(x) = \frac{1}{b-a} \int_a^b f(t) dt - \frac{1}{b-a} \sum_{k=1}^{n-1} F_k(x) \\ - \frac{(-1)^n}{b-a} \int_a^b K_n(x, t) f^{(n)}(t) dt.$$

Similarly, from the hypotheses on  $g$  we have the identity

$$(3.3) \quad g(x) = \frac{1}{b-a} \int_a^b g(t) dt - \frac{1}{b-a} \sum_{k=1}^{n-1} G_k(x) \\ - \frac{(-1)^n}{b-a} \int_a^b K_n(x, t) g^{(n)}(t) dt.$$



On Grüss Type Integral Inequalities

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit


Page 7 of 12

Multiplying (3.2) by  $g(x)$  and (3.3) by  $f(x)$  and summing the resulting identities and integrating from  $a$  to  $b$  and rewriting we have

$$(3.4) \quad \frac{1}{b-a} \int_a^b f(x) g(x) dx = \left( \frac{1}{b-a} \int_a^b f(x) dx \right) \left( \frac{1}{b-a} \int_a^b g(x) dx \right) \\ - \frac{1}{2(b-a)^2} \int_a^b \left[ \left( \sum_{k=1}^{n-1} F_k(x) \right) g(x) + \left( \sum_{k=1}^{n-1} G_k(x) \right) f(x) \right] dx \\ - \frac{1}{2(b-a)^2} \left[ \int_a^b g(x) \left\{ \frac{(-1)^n}{b-a} \int_a^b K_n(x,t) f^{(n)}(t) dt \right\} dx \right. \\ \left. + \int_a^b f(x) \left\{ \frac{(-1)^n}{b-a} \int_a^b K_n(x,t) g^{(n)}(t) dt \right\} dx \right].$$

From (3.4) we observe that

$$\left| \frac{1}{b-a} \int_a^b f(x) g(x) dx - \left( \frac{1}{b-a} \int_a^b f(x) dx \right) \left( \frac{1}{b-a} \int_a^b g(x) dx \right) \right. \\ \left. - \frac{1}{2(b-a)^2} \int_a^b \left[ \left( \sum_{k=1}^{n-1} F_k(x) \right) g(x) + \left( \sum_{k=1}^{n-1} G_k(x) \right) f(x) \right] dx \right| \\ \leq \frac{1}{2(b-a)^2} \int_a^b \left( |g(x)| \left( \int_a^b |K_n(x,t)| |f^{(n)}(t)| dt \right) \right. \\ \left. + |f(x)| \left( \int_a^b |K_n(x,t)| |g^{(n)}(t)| dt \right) \right) dx$$



On Grüss Type Integral  
Inequalities

B.G. Pachpatte

Title Page

Contents



Go Back

Close


Quit

Page 8 of 12



$$\leq \frac{1}{2(b-a)^2} \int_a^b (|g(x)| \|f^{(n)}\|_\infty + |f(x)| \|g^{(n)}\|_\infty) A_n(x),$$

which is the required inequality in (2.1). The proof is complete.




---

**On Grüss Type Integral Inequalities**

B.G. Pachpatte

---

Title Page

Contents



Go Back

Close

Quit

Page 9 of 12

## 4. Proof of Theorem 2.2

From the hypotheses on  $p$  we have the following integral identity (see [2, p. 291]):


$$(4.1) \quad \frac{1}{n} \left( p(x) + \sum_{k=1}^{n-1} P_k(x) \right) - \frac{1}{b-a} \int_a^b p(y) dy \\ = \frac{1}{n!(b-a)} \int_a^b (x-t)^{n-1} r(t,x) p^{(n)}(t) dt,$$

for  $x \in [a, b]$ . The identity (4.1) can be rewritten as

$$(4.2) \quad p(x) = \frac{n}{b-a} \int_a^b p(x) dx - \sum_{k=1}^{n-1} P_k(x) \\ + \frac{1}{(n-1)!(b-a)} \int_a^b (x-t)^{n-1} r(t,x) p^{(n)}(t) dt.$$

Similarly, from the hypotheses on  $q$  we have the identity

$$(4.3) \quad q(x) = \frac{n}{b-a} \int_a^b q(x) dx - \sum_{k=1}^{n-1} Q_k(x) \\ + \frac{1}{(n-1)!(b-a)} \int_a^b (x-t)^{n-1} r(t,x) q^{(n)}(t) dt.$$



On Grüss Type Integral  
Inequalities

B.G. Pachpatte

Title Page

Contents



Go Back

Close


Quit

Page 10 of 12

Multiplying (4.2) by  $q(x)$  and (4.3) by  $p(x)$  and summing the resulting identities and integrating from  $a$  to  $b$  and rewriting we have

$$(4.4) \quad \begin{aligned} \frac{1}{b-a} \int_a^b p(x) q(x) dx &= n \left( \frac{1}{b-a} \int_a^b p(x) dx \right) \left( \frac{1}{b-a} \int_a^b q(x) dx \right) \\ &\quad - \frac{1}{2(b-a)} \int_a^b \left[ \left( \sum_{k=1}^{n-1} P_k(x) \right) q(x) + \left( \sum_{k=1}^{n-1} Q_k(x) \right) p(x) \right] dx \\ &\quad + \frac{1}{2(n-1)!(b-a)^2} \left[ \int_a^b q(x) \left\{ \int_a^b (x-t)^{n-1} r(t,x) p^{(n)}(t) dt \right\} dx \right. \\ &\quad \left. + \int_a^b p(x) \left\{ \int_a^b (x-t)^{n-1} r(t,x) q^{(n)}(t) dt \right\} dx \right]. \end{aligned}$$

From (4.4) and following the similar arguments as in the last part of the proof of Theorem 2.1, we get the desired inequality in (2.6). The proof is complete.



### On Grüss Type Integral Inequalities

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 11 of 12

## References

- [1] P. CERONE, S.S. DRAGOMIR AND J. ROUMELIOTIS, Some Ostrowski type inequalities for  $n$ -time differentiable mappings and applications, *Demonstratio Mathematica*, **32**(2) (1999), 679–712. *RGMIA Research Report Collection*, **1**(1) (1998), 51–66. ONLINE: <http://rgmia.vu.edu.au/v1n1.html>
- [2] A.M. FINK, Bounds on the derivation of a function from its averages, *Czechoslovak Math. Jour.*, **42** (1992), 289–310.
- [3] G.V. MILOVANOVIĆ AND J.E. PEČARIĆ, On generalization of the inequality of A. Ostrowski and some related applications, *Univ. Beograd. Publ. Elek. Fak. Ser. Mat. Fiz.*, **544–576** (1976), 155–158.
- [4] D.S. MITRINOVIĆ, J.E. PEČARIĆ AND A.M. FINK, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht, 1993.
- [5] B.G. PACHPATTE, A note on some inequalities analogous to Grüss inequality, *Octagon Math. Magazine*, **5** (1997), 62–66.
- [6] J. SÁNDOR, On the Jensen-Hadamard inequality, *Studia Univ. Babeş-Bolyai, Math.*, **36**(1) (1991), 9–15.



---

On Grüss Type Integral  
Inequalities

B.G. Pachpatte

---

Title Page

Contents



Go Back

Close

Quit

Page 12 of 12