

CERTAIN CLASSES OF ANALYTIC FUNCTIONS INVOLVING SÄLÄGEAN OPERATOR



Analytic Functions Involving
Sälägean Operator
Sevtap Sümer Eker and
Shigeyoshi Owa

vol. 10, iss. 1, art. 22, 2009

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 1 of 25

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.

SEVTAP SÜMER EKER

Department of Mathematics
Faculty of Science and Letters
Dicle University
21280 - Diyarbakır, Turkey
EMail: sevtaps@dicle.edu.tr

SHIGEYOSHI OWA

Department of Mathematics
Kinki University
Higashi-Osaka, Osaka 577 - 8502
Japan
EMail: owa@math.kindai.ac.jp

Received: 21 March, 2007

Accepted: 27 December, 2008

Communicated by: [G. Kohr](#)

2000 AMS Sub. Class.: 26D15

Key words: Sälägean operator, coefficient inequalities, distortion inequalities, extreme points, integral means, fractional derivative.

Abstract: Using Sälägean differential operator, we study new subclasses of analytic functions. Coefficient inequalities and distortion theorems and extreme points of these classes are studied. Furthermore, integral means inequalities are obtained for the fractional derivatives of these classes.

Contents

1	Introduction and Definitions	3
2	Coefficient Inequalities for classes $\mathcal{N}_{m,n}(\alpha, \beta)$ and $\mathcal{M}_{m,n}^s(\alpha, \beta)$	5
3	Relation for $\widetilde{\mathcal{N}}_{m,n}(\alpha, \beta)$ and $\widetilde{\mathcal{M}}_{m,n}^s(\alpha, \beta)$	9
4	Distortion Inequalities	10
5	Extreme Points	15
6	Integral Means Inequalities	18



Analytic Functions Involving
Sălăgean Operator
Sevtap Sümer Eker and
Shigeyoshi Owa
vol. 10, iss. 1, art. 22, 2009

Title Page

Contents



Page 2 of 25


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



1. Introduction and Definitions

Let \mathcal{A} denote the class of functions $f(z)$ of the form

$$(1.1) \quad f(z) = z + \sum_{j=2}^{\infty} a_j z^j$$

which are analytic in the open disc $\mathbb{U} = \{z : |z| < 1\}$. Let \mathcal{S} be the subclass of \mathcal{A} consisting of analytic and univalent functions $f(z)$ in \mathbb{U} . We denote by $\mathcal{S}^*(\alpha)$ and $\mathcal{K}(\alpha)$ the class of starlike functions of order α and the class of convex functions of order α , respectively, that is,

$$\mathcal{S}^*(\alpha) = \left\{ f \in \mathcal{A} : \operatorname{Re} \left(\frac{z f'(z)}{f(z)} \right) > \alpha, 0 \leq \alpha < 1, z \in \mathbb{U} \right\}$$

and

$$\mathcal{K}(\alpha) = \left\{ f \in \mathcal{A} : \operatorname{Re} \left(1 + \frac{z f''(z)}{f'(z)} \right) > \alpha, 0 \leq \alpha < 1, z \in \mathbb{U} \right\}.$$

For $f(z) \in \mathcal{A}$, Sălăgean [1] introduced the following operator which is called the Sălăgean operator:

$$D^0 f(z) = f(z)$$

$$D^1 f(z) = Df(z) = z f'(z)$$

$$D^n f(z) = D(D^{n-1} f(z)) \quad (n \in \mathbb{N} = 1, 2, 3, \dots).$$

We note that,

$$D^n f(z) = z + \sum_{j=2}^{\infty} j^n a_j z^j \quad (n \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}).$$

Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 3 of 25


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 4 of 25

Go Back

Full Screen

Close

Let $\mathcal{N}_{m,n}(\alpha, \beta)$ denote the subclass of \mathcal{A} consisting of functions $f(z)$ which satisfy the inequality

$$\operatorname{Re} \left\{ \frac{D^m f(z)}{D^n f(z)} \right\} > \beta \left| \frac{D^m f(z)}{D^n f(z)} - 1 \right| + \alpha$$

for some $0 \leq \alpha < 1$, $\beta \geq 0$, $m \in \mathbb{N}$, $n \in \mathbb{N}_0$ and all $z \in \mathbb{U}$. Also let $\mathcal{M}_{m,n}^s(\alpha, \beta)$ ($s = 0, 1, 2, \dots$) be the subclass of \mathcal{A} consisting of functions $f(z)$ which satisfy the condition:

$$f(z) \in \mathcal{M}_{m,n}^s(\alpha, \beta) \Leftrightarrow D^s f(z) \in \mathcal{N}_{m,n}(\alpha, \beta).$$

It is easy to see that if $s = 0$, then $\mathcal{M}_{m,n}^0(\alpha, \beta) \equiv \mathcal{N}_{m,n}(\alpha, \beta)$. Furthermore, special cases of our classes are the following:

- (i) $\mathcal{N}_{1,0}(\alpha, 0) = \mathcal{S}^*(\alpha)$ and $\mathcal{N}_{2,1}(\alpha, 0) = \mathcal{K}(\alpha)$ which were studied by Silverman [2].
- (ii) $\mathcal{N}_{1,0}(\alpha, \beta) = \mathcal{SD}(\alpha, \beta)$ and $\mathcal{M}_{1,0}^1(\alpha, \beta) = \mathcal{KD}(\alpha, \beta)$ which were studied by Shams at all [3].
- (iii) $\mathcal{N}_{m,n}(\alpha, 0) = \mathcal{K}_{m,n}(\alpha)$ and $\mathcal{M}_{m,n}^s(\alpha, 0) = \mathcal{M}_{m,n}^s(\alpha)$ which were studied by Eker and Owa [4].

Therefore, our present paper is a generalization of these papers. In view of the coefficient inequalities for $f(z)$ to be in the classes $\mathcal{N}_{m,n}(\alpha, \beta)$ and $\mathcal{M}_{m,n}^s(\alpha, \beta)$, we introduce two subclasses $\tilde{\mathcal{N}}_{m,n}(\alpha, \beta)$ and $\tilde{\mathcal{M}}_{m,n}^s(\alpha, \beta)$. Some distortion inequalities for $f(z)$ and some integral means inequalities for fractional calculus of $f(z)$ in the above classes are discussed in this paper.



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 5 of 25

Go Back

Full Screen

Close

2. Coefficient Inequalities for classes $\mathcal{N}_{m,n}(\alpha, \beta)$ and $\mathcal{M}_{m,n}^s(\alpha, \beta)$

Theorem 2.1. *If $f(z) \in \mathcal{A}$ satisfies*

$$(2.1) \quad \sum_{j=2}^{\infty} \Psi(m, n, j, \alpha, \beta) |a_j| \leq 2(1 - \alpha)$$

where

$$(2.2) \quad \Psi(m, n, j, \alpha, \beta) = |j^m - j^n - \alpha j^n| + (j^m + j^n - \alpha j^n) + 2\beta |j^m - j^n|$$

for some $\alpha(0 \leq \alpha < 1)$, $\beta \geq 0$, $m \in \mathbb{N}$ and $n \in \mathbb{N}_0$, then $f(z) \in \mathcal{N}_{m,n}(\alpha, \beta)$.

Proof. Suppose that (2.1) is true for $\alpha(0 \leq \alpha < 1)$, $\beta \geq 0$, $m \in \mathbb{N}$, $n \in \mathbb{N}_0$. For $f(z) \in \mathcal{A}$, let us define the function $F(z)$ by


$$F(z) = \frac{D^m f(z)}{D^n f(z)} - \beta \left| \frac{D^m f(z)}{D^n f(z)} - 1 \right| - \alpha.$$

It suffices to show that

$$\left| \frac{F(z) - 1}{F(z) + 1} \right| < 1 \quad (z \in \mathbb{U}).$$

We note that

$$\begin{aligned} & \left| \frac{F(z) - 1}{F(z) + 1} \right| \\ &= \left| \frac{D^m f(z) - \beta e^{i\theta} |D^m f(z) - D^n f(z)| - \alpha D^n f(z) - D^n f(z)}{D^m f(z) - \beta e^{i\theta} |D^m f(z) - D^n f(z)| - \alpha D^n f(z) + D^n f(z)} \right| \end{aligned}$$



$$\begin{aligned}
 &= \left| \frac{-\alpha + \sum_{j=2}^{\infty} (j^m - j^n - \alpha j^n) a_j z^{j-1} - \beta e^{i\theta} \left| \sum_{j=2}^{\infty} (j^m - j^n) a_j z^{j-1} \right|}{(2 - \alpha) + \sum_{j=2}^{\infty} (j^m + j^n - \alpha j^n) a_j z^{j-1} - \beta e^{i\theta} \left| \sum_{j=2}^{\infty} (j^m - j^n) a_j z^{j-1} \right|} \right| \\
 &\leq \frac{\alpha + \sum_{j=2}^{\infty} |j^m - j^n - \alpha j^n| |a_j| |z|^{j-1} + \beta |e^{i\theta}| \sum_{j=2}^{\infty} |j^m - j^n| |a_j| |z|^{j-1}}{(2 - \alpha) - \sum_{j=2}^{\infty} (j^m + j^n - \alpha j^n) |a_j| |z|^{j-1} - \beta |e^{i\theta}| \sum_{j=2}^{\infty} |j^m - j^n| |a_j| |z|^{j-1}} \\
 &\leq \frac{\alpha + \sum_{j=2}^{\infty} |j^m - j^n - \alpha j^n| |a_j| + \beta \sum_{j=2}^{\infty} |j^m - j^n| |a_j|}{(2 - \alpha) - \sum_{j=2}^{\infty} (j^m + j^n - \alpha j^n) |a_j| - \beta \sum_{j=2}^{\infty} |j^m - j^n| |a_j|}.
 \end{aligned}$$

The last expression is bounded above by 1, if

$$\begin{aligned}
 &\alpha + \sum_{j=2}^{\infty} |j^m - j^n - \alpha j^n| |a_j| + \beta \sum_{j=2}^{\infty} |j^m - j^n| |a_j| \\
 &\leq (2 - \alpha) - \sum_{j=2}^{\infty} (j^m + j^n - \alpha j^n) |a_j| - \beta \sum_{j=2}^{\infty} |j^m - j^n| |a_j|
 \end{aligned}$$

which is equivalent to our condition (2.1). This completes the proof of our theorem. \square

By using Theorem 2.1, we have:

Theorem 2.2. *If $f(z) \in \mathcal{A}$ satisfies*

$$\sum_{j=2}^{\infty} j^s \Psi(m, n, j, \alpha, \beta) |a_j| \leq 2(1 - \alpha),$$

where $\Psi(m, n, j, \alpha, \beta)$ is defined by (2.2) for some $\alpha(0 \leq \alpha < 1)$, $\beta \geq 0$, $m \in \mathbb{N}$ and $n \in \mathbb{N}_0$, then $f(z) \in \mathcal{M}_{m,n}^s(\alpha, \beta)$.

Title Page

Contents

◀◀ ▶▶


◀ ▶

Page 6 of 25

Go Back

Full Screen

Close



Title Page

Contents



Page 7 of 25

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Proof. From

$$f(z) \in \mathcal{M}_{m,n}^s(\alpha, \beta) \Leftrightarrow D^s f(z) \in \mathcal{N}_{m,n}(\alpha, \beta),$$

replacing a_j by $j^s a_j$ in Theorem 2.1, we have the theorem. □

Example 2.1. The function $f(z)$ given by

$$f(z) = z + \sum_{j=2}^{\infty} \frac{2(2+\delta)(1-\alpha)\epsilon_j}{(j+\delta)(j+1+\delta)\Psi(m, n, j, \alpha, \beta)} z^j = z + \sum_{j=2}^{\infty} A_j z^j$$

with

$$A_j = \frac{2(2+\delta)(1-\alpha)\epsilon_j}{(j+\delta)(j+1+\delta)\Psi(m, n, j, \alpha, \beta)}$$

belongs to the class $\mathcal{N}_{m,n}(\alpha, \beta)$ for $\delta > -2$, $0 \leq \alpha < 1$, $\beta \geq 0$, $\epsilon_j \in \mathbb{C}$ and $|\epsilon_j| = 1$.

Because, we know that

$$\begin{aligned} \sum_{j=2}^{\infty} \Psi(m, n, j, \alpha, \beta) |A_j| &\leq \sum_{j=2}^{\infty} \frac{2(2+\delta)(1-\alpha)}{(j+\delta)(j+1+\delta)} \\ &= \sum_{j=2}^{\infty} 2(2+\delta)(1-\alpha) \sum_{j=2}^{\infty} \left(\frac{1}{j+\delta} - \frac{1}{j+1+\delta} \right) \\ &= 2(1-\alpha). \end{aligned}$$

Example 2.2. The function $f(z)$ given by

$$f(z) = z + \sum_{j=2}^{\infty} \frac{2(2+\delta)(1-\alpha)\epsilon_j}{j^s(j+\delta)(j+1+\delta)\Psi(m, n, j, \alpha, \beta)} z^j = z + \sum_{j=2}^{\infty} B_j z^j$$

with

$$B_j = \frac{2(2+\delta)(1-\alpha)\epsilon_j}{j^s(j+\delta)(j+1+\delta)\Psi(m, n, j, \alpha, \beta)}$$

belongs to the class $\mathcal{M}_{m,n}^s(\alpha, \beta)$ for $\delta > -2$, $0 \leq \alpha < 1$, $\beta \geq 0$, $\epsilon_j \in \mathbb{C}$ and $|\epsilon_j| = 1$.
 Because, the function $f(z)$ gives us that

$$\sum_{j=2}^{\infty} j^s \Psi(m, n, j, \alpha, \beta) |B_j| \leq \sum_{j=2}^{\infty} \frac{2(2 + \delta)(1 - \alpha)}{(j + \delta)(j + 1 + \delta)} = 2(1 - \alpha).$$



**Analytic Functions Involving
 Sălăgean Operator**
 Sevtap Sümer Eker and
 Shigeyoshi Owa
 vol. 10, iss. 1, art. 22, 2009

Title Page

Contents



Page 8 of 25


Go Back

Full Screen

Close

journal of **inequalities**
 in pure and applied
 mathematics

issn: 1443-5756



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 9 of 25

Go Back

Full Screen

Close

3. Relation for $\tilde{\mathcal{N}}_{m,n}(\alpha, \beta)$ and $\tilde{\mathcal{M}}_{m,n}^s(\alpha, \beta)$

In view of Theorem 2.1 and Theorem 2.2, we now introduce the subclasses

$$\tilde{\mathcal{N}}_{m,n}(\alpha, \beta) \subset \mathcal{N}_{m,n}(\alpha, \beta) \quad \text{and} \quad \tilde{\mathcal{M}}_{m,n}^s(\alpha, \beta) \subset \mathcal{M}_{m,n}^s(\alpha, \beta)$$

which consist of functions

$$(3.1) \quad f(z) = z + \sum_{j=2}^{\infty} a_j z^j \quad (a_j \geq 0)$$

whose Taylor-Maclaurin coefficients satisfy the inequalities (2.1) and (2.2), respectively. By the coefficient inequalities for the classes $\tilde{\mathcal{N}}_{m,n}(\alpha, \beta)$ and $\tilde{\mathcal{M}}_{m,n}^s(\alpha, \beta)$, we see:

Theorem 3.1.

$$\tilde{\mathcal{N}}_{m,n}(\alpha, \beta_2) \subset \tilde{\mathcal{N}}_{m,n}(\alpha, \beta_1)$$

for some β_1 and β_2 , $0 \leq \beta_1 \leq \beta_2$.

Proof. For $0 \leq \beta_1 \leq \beta_2$ we obtain

$$\sum_{j=2}^{\infty} \Psi(m, n, j, \alpha, \beta_1) a_j \leq \sum_{j=2}^{\infty} \Psi(m, n, j, \alpha, \beta_2) a_j.$$


Therefore, if $f(z) \in \tilde{\mathcal{N}}_{m,n}(\alpha, \beta_2)$, then $f(z) \in \tilde{\mathcal{N}}_{m,n}(\alpha, \beta_1)$. Hence we get the required result. □

By using Theorem 3.1, we also have

Corollary 3.2.

$$\tilde{\mathcal{M}}_{m,n}^s(\alpha, \beta_2) \subset \tilde{\mathcal{M}}_{m,n}^s(\alpha, \beta_1)$$

for some β_1 and β_2 , $0 \leq \beta_1 \leq \beta_2$.



4. Distortion Inequalities

Lemma 4.1. *If $f(z) \in \tilde{\mathcal{N}}_{m,n}(\alpha, \beta)$, then we have*

$$\sum_{j=p+1}^{\infty} a_j \leq \frac{2(1-\alpha) - \sum_{j=2}^p \Psi(m, n, j, \alpha, \beta)a_j}{\Psi(m, n, p+1, \alpha, \beta)}.$$

Proof. In view of Theorem 2.1, we can write

$$(4.1) \quad \sum_{j=p+1}^{\infty} \Psi(m, n, j, \alpha, \beta)a_j \leq 2(1-\alpha) - \sum_{j=2}^p \Psi(m, n, j, \alpha, \beta)a_j.$$

Clearly $\Psi(m, n, j, \alpha, \beta)$ is an increasing function for j . Then from (2.2) and (4.1), we have

$$\Psi(m, n, p+1, \alpha, \beta) \sum_{j=p+1}^{\infty} a_j \leq 2(1-\alpha) - \sum_{j=2}^p \Psi(m, n, j, \alpha, \beta)a_j.$$

Thus, we obtain

$$\sum_{j=p+1}^{\infty} a_j \leq \frac{2(1-\alpha) - \sum_{j=2}^p \Psi(m, n, j, \alpha, \beta)a_j}{\Psi(m, n, p+1, \alpha, \beta)} = A_j.$$

□

Lemma 4.2. *If $f(z) \in \tilde{\mathcal{N}}_{m,n}(\alpha, \beta)$, then*

$$\sum_{j=p+1}^{\infty} ja_j \leq \frac{2(1-\alpha) - \sum_{j=2}^p \Psi(m, n, j, \alpha, \beta)a_j}{\Psi(m-1, n-1, p+1, \alpha, \beta)} = B_j.$$

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 10 of 25


[Go Back](#)

[Full Screen](#)

[Close](#)


journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents



Page 11 of 25

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Corollary 4.3. If $f(z) \in \widetilde{M}_{m,n}^s(\alpha)$, then

$$\sum_{j=p+1}^{\infty} a_j \leq \frac{2(1-\alpha) - \sum_{j=2}^p j^s \Psi(m, n, j, \alpha, \beta) a_j}{(p+1)^s \Psi(m, n, p+1, \alpha, \beta)} = C_j$$

and

$$\sum_{j=p+1}^{\infty} j a_j \leq \frac{2(1-\alpha) - \sum_{j=2}^p j^s \Psi(m, n, j, \alpha, \beta) a_j}{(p+1)^s \Psi(m-1, n-1, p+1, \alpha, \beta)} = D_j.$$

Theorem 4.4. Let $f(z) \in \widetilde{N}_{m,n}(\alpha, \beta)$. Then for $|z| = r < 1$

$$r - \sum_{j=2}^p a_j |z|^j - A_j r^{p+1} \leq |f(z)| \leq r + \sum_{j=2}^p a_j |z|^j + A_j r^{p+1}$$

and

$$1 - \sum_{j=2}^p j a_j |z|^{j-1} - B_j r^p \leq |f'(z)| \leq 1 + \sum_{j=2}^p j a_j |z|^j + B_j r^p$$

where A_j and B_j are given by Lemma 4.1 and Lemma 4.2.

Proof. Let $f(z)$ given by (1.1). For $|z| = r < 1$, using Lemma 4.1, we have

$$\begin{aligned} |f(z)| &\leq |z| + \sum_{j=2}^p a_j |z|^j + \sum_{j=p+1}^{\infty} a_j |z|^j \\ &\leq |z| + \sum_{j=2}^p a_j |z|^j + |z|^{p+1} \sum_{j=p+1}^{\infty} a_j \\ &\leq r + \sum_{j=2}^p a_j |z|^j + A_j r^{p+1} \end{aligned}$$

and


$$\begin{aligned}
 |f(z)| &\geq |z| - \sum_{j=2}^p a_j |z|^j - \sum_{j=p+1}^{\infty} a_j |z|^j \\
 &\geq |z| - \sum_{j=2}^p a_j |z|^j - |z|^{p+1} \sum_{j=p+1}^{\infty} a_j \\
 &\geq r - \sum_{j=2}^p a_j |z|^j - A_j r^{p+1}.
 \end{aligned}$$

Furthermore, for $|z| = r < 1$ using Lemma 4.2, we obtain

$$\begin{aligned}
 |f'(z)| &\leq 1 + \sum_{j=2}^p j a_j |z|^{j-1} + \sum_{j=p+1}^{\infty} j a_j |z|^{j-1} \\
 &\leq 1 + \sum_{j=2}^p j a_j |z|^{j-1} + |z|^p \sum_{j=p+1}^{\infty} j a_j \\
 &\leq 1 + \sum_{j=2}^p j a_j |z|^{j-1} + B_j r^p
 \end{aligned}$$



and

$$\begin{aligned}
 |f'(z)| &\geq 1 - \sum_{j=2}^p j a_j |z|^{j-1} - \sum_{j=p+1}^{\infty} j a_j |z|^{j-1} \\
 &\geq 1 - \sum_{j=2}^p j a_j |z|^{j-1} - |z|^p \sum_{j=p+1}^{\infty} j a_j
 \end{aligned}$$



Title Page

Contents



Page 12 of 25

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

$$\geq 1 - \sum_{j=2}^p j a_j |z|^{j-1} - B_j r^p.$$

This completes the assertion of Theorem 4.4. □

Theorem 4.5. Let $f(z) \in \widetilde{\mathcal{M}}_{m,n}^s(\alpha, \beta)$. Then

$$r - \sum_{j=2}^p a_j |z|^j - C_j r^{p+1} \leq |f(z)| \leq r + \sum_{j=2}^p a_j |z|^j + C_j r^{p+1}$$

and

$$1 - \sum_{j=2}^p j a_j |z|^{j-1} - D_j r^p \leq |f'(z)| \leq 1 + \sum_{j=2}^p j a_j |z|^j + D_j r^p$$

where C_j and D_j are given by Corollary 4.3.

Proof. Using a similar method to that in the proof of Theorem 4.4 and making use Corollary 4.3, we get our result. □

Taking $p = 1$ in Theorem 4.4 and Theorem 4.5, we have:

Corollary 4.6. Let $f(z) \in \widetilde{\mathcal{N}}_{m,n}(\alpha, \beta)$. Then for $|z| = r < 1$

$$r - \frac{2(1-\alpha)}{\Psi(m, n, 2, \alpha, \beta)} r^2 \leq |f(z)| \leq r + \frac{2(1-\alpha)}{\Psi(m, n, 2, \alpha, \beta)} r^2$$

and

$$1 - \frac{2(1-\alpha)}{\Psi(m-1, n-1, 2, \alpha, \beta)} r \leq |f'(z)| \leq 1 + \frac{2(1-\alpha)}{\Psi(m-1, n-1, 2, \alpha, \beta)} r.$$



Analytic Functions Involving

Sălăgean Operator



Sevtap Sümer Eker and

Shigeyoshi Owa

vol. 10, iss. 1, art. 22, 2009

Title Page

Contents



Page 13 of 25


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Corollary 4.7. Let $f(z) \in \widetilde{\mathcal{M}}_{m,n}^s(\alpha, \beta)$. Then for $|z| = r < 1$

$$r - \frac{2(1-\alpha)}{2^s \Psi(m, n, 2, \alpha, \beta)} r^2 \leq |f(z)| \leq r + \frac{2(1-\alpha)}{2^s \Psi(m, n, 2, \alpha, \beta)} r^2$$

and

$$1 - \frac{2(1-\alpha)}{2^s \Psi(m-1, n-1, 2, \alpha, \beta)} r \leq |f'(z)| \leq 1 + \frac{2(1-\alpha)}{2^s \Psi(m-1, n-1, 2, \alpha, \beta)} r.$$

Analytic Functions Involving
Sălăgean Operator
Sevtap Sümer Eker and
Shigeyoshi Owa

vol. 10, iss. 1, art. 22, 2009

Title Page

Contents



Page 14 of 25

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 15 of 25

Go Back

Full Screen

Close

5. Extreme Points

The determination of the extreme points of a family F of univalent functions enables us to solve many extremal problems for F . Now, let us determine extreme points of the classes $\tilde{\mathcal{N}}_{m,n}(\alpha, \beta)$ and $\tilde{\mathcal{M}}_{m,n}^s(\alpha, \beta)$.

Theorem 5.1. Let $f_1(z) = z$ and

$$f_j(z) = z + \frac{2(1-\alpha)}{\Psi(m, n, j, \alpha, \beta)} z^j \quad (j = 2, 3, \dots).$$

where $\Psi(m, n, j, \alpha, \beta)$ is defined by (2.2). Then $f \in \tilde{\mathcal{N}}_{m,n}(\alpha, \beta)$ if and only if it can be expressed in the form

$$f(z) = \sum_{j=1}^{\infty} \lambda_j f_j(z),$$


where $\lambda_j > 0$ and $\sum_{j=1}^{\infty} \lambda_j = 1$.

Proof. Suppose that

$$f(z) = \sum_{j=1}^{\infty} \lambda_j f_j(z) = z + \sum_{j=2}^{\infty} \lambda_j \frac{2(1-\alpha)}{\Psi(m, n, j, \alpha, \beta)} z^j.$$

Then

$$\begin{aligned} & \sum_{j=2}^{\infty} \Psi(m, n, j, \alpha, \beta) \frac{2(1-\alpha)}{\Psi(m, n, j, \alpha, \beta)} \lambda_j \\ &= \sum_{j=2}^{\infty} 2(1-\alpha) \lambda_j = 2(1-\alpha) \sum_{j=2}^{\infty} \lambda_j = 2(1-\alpha)(1-\lambda_1) < 2(1-\alpha) \end{aligned}$$



[Title Page](#)

[Contents](#)

◀◀ ▶▶

◀ ▶

Page 16 of 25

[Go Back](#)

[Full Screen](#)

[Close](#)

Thus, $f(z) \in \tilde{\mathcal{N}}_{m,n}(\alpha, \beta)$ from the definition of the class of $\tilde{\mathcal{N}}_{m,n}(\alpha, \beta)$.

Conversely, suppose that $f \in \tilde{\mathcal{N}}_{m,n}(\alpha, \beta)$. Since

$$a_j \leq \frac{2(1-\alpha)}{\Psi(m, n, j, \alpha, \beta)} \quad (j = 2, 3, \dots),$$

we may set

$$\lambda_j = \frac{\Psi(m, n, j, \alpha, \beta)}{2(1-\alpha)} a_j$$

and

$$\lambda_1 = 1 - \sum_{j=2}^{\infty} \lambda_j.$$

Then,

$$f(z) = \sum_{j=1}^{\infty} \lambda_j f_j(z).$$

This completes the proof of the theorem. □

Corollary 5.2. Let $g_1(z) = z$ and

$$g_j(z) = z + \frac{2(1-\alpha)}{j^s \Psi(m, n, j, \alpha, \beta)} z^j \quad (j = 2, 3, \dots).$$

Then $g \in \tilde{\mathcal{M}}_{m,n}^s(\alpha, \beta)$ if and only if it can be expressed in the form

$$g(z) = \sum_{j=1}^{\infty} \lambda_j g_j(z),$$

where $\lambda_j > 0$ and $\sum_{j=1}^{\infty} \lambda_j = 1$.



Corollary 5.3. *The extreme points of $\tilde{\mathcal{N}}_{m,n}(\alpha, \beta)$ are the functions $f_1(z) = z$ and*

$$f_j(z) = z + \frac{2(1 - \alpha)}{\Psi(m, n, j, \alpha, \beta)} z^j \quad (j = 2, 3, \dots).$$


Corollary 5.4. *The extreme points of $\tilde{\mathcal{M}}_{m,n}^s(\alpha, \beta)$ are given by $g_1(z) = z$ and*

$$g_j(z) = z + \frac{2(1 - \alpha)}{j^s \Psi(m, n, j, \alpha, \beta)} z^j \quad (j = 2, 3, \dots).$$

Analytic Functions Involving
Sălăgean Operator
Sevtap Sümer Eker and
Shigeyoshi Owa
vol. 10, iss. 1, art. 22, 2009

Title Page

Contents



Page 17 of 25

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



6. Integral Means Inequalities

We shall use the following definitions for fractional derivatives by Owa [6] (also Srivastava and Owa [7]).

Definition 6.1. *The fractional derivative of order λ is defined, for a function $f(z)$, by*

$$(6.1) \quad D_z^\lambda f(z) = \frac{1}{\Gamma(1-\lambda)} \frac{d}{dz} \int_0^z \frac{f(\xi)}{(z-\xi)^\lambda} d\xi \quad (0 \leq \lambda < 1),$$

where the function $f(z)$ is analytic in a simply-connected region of the complex z -plane containing the origin, and the multiplicity of $(z-\xi)^{-\lambda}$ is removed by requiring $\log(z-\xi)$ to be real when $(z-\xi) > 0$.

Definition 6.2. *Under the hypotheses of Definition 6.1, the fractional derivative of order $(p+\lambda)$ is defined, for a function $f(z)$, by*

$$D_z^{p+\lambda} f(z) = \frac{d^p}{dz^p} D_z^\lambda f(z)$$

where $0 \leq \lambda < 1$ and $p \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}$.

It readily follows from (6.1) in Definition 6.1 that

$$(6.2) \quad D_z^\lambda z^k = \frac{\Gamma(k+1)}{\Gamma(k-\lambda+1)} z^{k-\lambda} \quad (0 \leq \lambda < 1).$$

Further, we need the concept of subordination between analytic functions and a subordination theorem by Littlewood [5] in our investigation.

Let us consider two functions $f(z)$ and $g(z)$, which are analytic in \mathbb{U} . The function $f(z)$ is said to be *subordinate* to $g(z)$ in \mathbb{U} if there exists a function $w(z)$ analytic

Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 18 of 25


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 19 of 25

[Go Back](#)

[Full Screen](#)

[Close](#)

in \mathbb{U} with

$$w(0) = 0 \quad \text{and} \quad |w(z)| < 1 \quad (z \in \mathbb{U}),$$

such that

$$f(z) = g(w(z)) \quad (z \in \mathbb{U}).$$

We denote this subordination by

$$f(z) \prec g(z).$$

Theorem 6.3 (Littlewood [5]). *If $f(z)$ and $g(z)$ are analytic in \mathbb{U} with $f(z) \prec g(z)$, then for $\mu > 0$ and $z = re^{i\theta}$ ($0 < r < 1$)*


$$\int_0^{2\pi} |f(z)|^\mu d\theta \leq \int_0^{2\pi} |g(z)|^\mu d\theta.$$

Theorem 6.4. *Let $f(z) \in \mathcal{A}$ given by (3.1) be in the class $\tilde{\mathcal{N}}_{m,n}(\alpha, \beta)$ and suppose that*

$$\sum_{j=2}^{\infty} (j-p)_{p+1} a_j \leq \frac{2(1-\alpha)\Gamma(k+1)\Gamma(3-\lambda-p)}{\Psi(m, n, k, \alpha, \beta)\Gamma(k+1-\lambda-p)\Gamma(2-p)}$$

for some $0 \leq p \leq 2$, $0 \leq \lambda < 1$ where $(j-p)_{p+1}$ denotes the Pochhammer symbol defined by $(j-p)_{p+1} = (j-p)(j-p+1) \cdots j$. Also given is the function $f_k(z)$ by

$$(6.3) \quad f_k(z) = z + \frac{2(1-\alpha)}{\Psi(m, n, k, \alpha, \beta)} z^k \quad (k \geq 2).$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 20 of 25

Go Back

Full Screen

Close

If there exists an analytic function $w(z)$ given by

$$\{w(z)\}^{k-1} = \frac{\Psi(m, n, k, \alpha, \beta)\Gamma(k+1-\lambda-p)}{2(1-\alpha)\Gamma(k+1)} \\ \times \sum_{j=2}^{\infty} (j-p)_{p+1} \frac{\Gamma(j-p)}{\Gamma(j+1-\lambda-p)} a_j z^{j-1},$$

then for $z = re^{i\theta}$ ($0 < r < 1$) and $\mu > 0$,

$$\int_0^{2\pi} |D_z^{p+\lambda} f(z)|^\mu d\theta \leq \int_0^{2\pi} |D_z^{p+\lambda} f_k(z)|^\mu d\theta.$$

Proof. By virtue of the fractional derivative formula (6.2) and Definition 6.2, we find from (1.1) that


$$D_z^{p+\lambda} f(z) = \frac{z^{1-\lambda-p}}{\Gamma(2-\lambda-p)} \left\{ 1 + \sum_{j=2}^{\infty} \frac{\Gamma(2-\lambda-p)\Gamma(j+1)}{\Gamma(j+1-\lambda-p)} a_j z^{j-1} \right\} \\ = \frac{z^{1-\lambda-p}}{\Gamma(2-\lambda-p)} \left\{ 1 + \sum_{j=2}^{\infty} \Gamma(2-\lambda-p)(j-p)_{p+1} \Phi(j) a_j z^{j-1} \right\}$$

where

$$\Phi(j) = \frac{\Gamma(j-p)}{\Gamma(j+1-\lambda-p)}.$$


Since $\Phi(j)$ is a decreasing function of j , we have

$$0 < \Phi(j) \leq \Phi(2) = \frac{\Gamma(2-p)}{\Gamma(3-\lambda-p)} \quad (0 \leq \lambda < 1 \quad ; \quad 0 \leq p \leq 2 \leq j).$$



Title Page

Contents



Page 21 of 25

Go Back

Full Screen

Close

Similarly, from (6.2), (6.3) and Definition 6.2, we obtain

$$D_z^{p+\lambda} f_k(z) = \frac{z^{1-\lambda-p}}{\Gamma(2-\lambda-p)} \left\{ 1 + \frac{2(1-\alpha)\Gamma(2-\lambda-p)\Gamma(k+1)}{\Psi(m, n, k, \alpha, \beta)\Gamma(k+1-\lambda-p)} z^{k-1} \right\}.$$

For $z = re^{i\theta}$, $0 < r < 1$, we must show that

$$\begin{aligned} & \int_0^{2\pi} \left| 1 + \sum_{j=2}^{\infty} \Gamma(2-\lambda-p)(j-p)_{p+1} \Phi(j) a_j z^{j-1} \right|^\mu d\theta \\ & \leq \int_0^{2\pi} \left| 1 + \frac{2(1-\alpha)\Gamma(2-\lambda-p)\Gamma(k+1)}{\Psi(m, n, k, \alpha, \beta)\Gamma(k+1-\lambda-p)} z^{k-1} \right|^\mu d\theta \quad (\mu > 0). \end{aligned}$$

Thus by applying Littlewood's subordination theorem, it would suffice to show that


$$(6.4) \quad 1 + \sum_{j=2}^{\infty} \Gamma(2-\lambda-p)(j-p)_{p+1} \Phi(j) a_j z^{j-1} < 1 + \frac{2(1-\alpha)\Gamma(2-\lambda-p)\Gamma(k+1)}{\Psi(m, n, k, \alpha, \beta)\Gamma(k+1-\lambda-p)} z^{k-1}.$$

By setting

$$\begin{aligned} & 1 + \sum_{j=2}^{\infty} \Gamma(2-\lambda-p)(j-p)_{p+1} \Phi(j) a_j z^{j-1} \\ & = 1 + \frac{2(1-\alpha)\Gamma(2-\lambda-p)\Gamma(k+1)}{\Psi(m, n, k, \alpha, \beta)\Gamma(k+1-\lambda-p)} \{w(z)\}^{k-1} \end{aligned}$$

we find that

$$\{w(z)\}^{k-1} = \frac{\Psi(m, n, k, \alpha, \beta)\Gamma(k+1-\lambda-p)}{2(1-\alpha)\Gamma(k+1)} \sum_{j=2}^{\infty} (j-p)_{p+1} \Phi(j) a_j z^{j-1}$$



Title Page

Contents



Page 22 of 25

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.

which readily yields $w(0) = 0$.

Therefore, we have

$$\begin{aligned}
 |w(z)|^{k-1} &= \left| \frac{\Psi(m, n, k, \alpha, \beta) \Gamma(k+1-\lambda-p)}{2(1-\alpha) \Gamma(k+1)} \sum_{j=2}^{\infty} (j-p)_{p+1} \Phi(j) a_j z^{j-1} \right| \\
 &\leq \frac{\Psi(m, n, k, \alpha, \beta) \Gamma(k+1-\lambda-p)}{2(1-\alpha) \Gamma(k+1)} \sum_{j=2}^{\infty} (j-p)_{p+1} \Phi(j) a_j |z|^{j-1} \\
 &\leq |z| \frac{\Psi(m, n, k, \alpha, \beta) \Gamma(k+1-\lambda-p)}{2(1-\alpha) \Gamma(k+1)} \Phi(2) \sum_{j=2}^{\infty} (j-p)_{p+1} a_j \\
 &= |z| \frac{\Psi(m, n, k, \alpha, \beta) \Gamma(k+1-\lambda-p)}{2(1-\alpha) \Gamma(k+1)} \frac{\Gamma(2-p)}{\Gamma(3-\lambda-p)} \sum_{j=2}^{\infty} (j-p)_{p+1} a_j \\
 &\leq |z| < 1
 \end{aligned}$$

by means of the hypothesis of Theorem 6.4. □


For the special case $p = 0$, Theorem 6.4 readily yields the following result.

Corollary 6.5. Let $f(z) \in \mathcal{A}$ given by (3.1) be in the class $\tilde{\mathcal{N}}_{m,n}(\alpha, \beta)$ and suppose that

$$\sum_{j=2}^{\infty} j a_j \leq \frac{2(1-\alpha) \Gamma(k+1) \Gamma(3-\lambda)}{\Psi(m, n, k, \alpha, \beta) \Gamma(k+1-\lambda)}$$

for $0 \leq \lambda < 1$. Also let the function $f_k(z)$ be given by (6.3). If there exists an analytic function $w(z)$ given by

$$\{w(z)\}^{k-1} = \frac{\Psi(m, n, k, \alpha, \beta) \Gamma(k+1-\lambda)}{2(1-\alpha) \Gamma(k+1)} \sum_{j=2}^{\infty} \frac{\Gamma(j+1)}{\Gamma(j+1-\lambda)} a_j z^{j-1},$$



[Title Page](#)

[Contents](#)

◀◀ ▶▶

◀ ▶

Page 23 of 25

[Go Back](#)

[Full Screen](#)

[Close](#)

then for $z = re^{i\theta}$ and $0 < r < 1$,

$$\int_0^{2\pi} |D_z^\lambda f(z)|^\mu d\theta \leq \int_0^{2\pi} |D_z^\lambda f_k(z)|^\mu d\theta \quad (0 \leq \lambda < 1, \mu > 0).$$

Corollary 6.6. Let $f(z) \in \mathcal{A}$ given by (3.1) be in the class $\widetilde{\mathcal{M}}_{m,n}^s(\alpha, \beta)$ and suppose that

$$\sum_{j=2}^{\infty} (j-p)_{p+1} a_j \leq \frac{2(1-\alpha)\Gamma(k+1)\Gamma(3-\lambda-p)}{k^s \Psi(m, n, k, \alpha, \beta) \Gamma(k+1-\lambda-p) \Gamma(2-p)}$$

for some $0 \leq p \leq 2$, $0 \leq \lambda < 1$. Also let the function

$$(6.5) \quad g_k(z) = z + \frac{2(1-\alpha)}{k^s \Psi(m, n, k, \alpha, \beta)} z^k, \quad (k \geq 2).$$


If there exists an analytic function $w(z)$ given by

$$\begin{aligned} \{w(z)\}^{k-1} &= \frac{k^s \Psi(m, n, k, \alpha, \beta) \Gamma(k+1-\lambda-p)}{2(1-\alpha)\Gamma(k+1)} \\ &\quad \times \sum_{j=2}^{\infty} (j-p)_{p+1} \frac{\Gamma(j-p)}{\Gamma(j+1-\lambda-p)} a_j z^{j-1}, \end{aligned}$$

then for $z = re^{i\theta}$ ($0 < r < 1$) and $\mu > 0$,



$$\int_0^{2\pi} |D_z^{p+\lambda} f(z)|^\mu d\theta \leq \int_0^{2\pi} |D_z^{p+\lambda} g_k(z)|^\mu d\theta.$$

For the special case $p = 0$, Corollary 6.6 readily yields,



Title Page

Contents



Page 24 of 25

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Corollary 6.7. Let $f(z) \in \mathcal{A}$ given by (3.1) be in the class $\widetilde{\mathcal{M}}_{m,n}^s(\alpha, \beta)$ and suppose that

$$\sum_{j=2}^{\infty} j a_j \leq \frac{2(1-\alpha)\Gamma(k+1)\Gamma(3-\lambda)}{k^s \Psi(m, n, k, \alpha, \beta)\Gamma(k+1-\lambda)}$$

for $0 \leq \lambda < 1$. Also let the function $g_k(z)$ be given by (6.5). If there exists an analytic function $w(z)$ given by

$$\{w(z)\}^{k-1} = \frac{k^s \Psi(m, n, k, \alpha, \beta)\Gamma(k+1-\lambda)}{2(1-\alpha)\Gamma(k+1)} \sum_{j=2}^{\infty} \frac{\Gamma(j+1)}{\Gamma(j+1-\lambda)} a_j z^{j-1},$$

then for $z = re^{i\theta}$ ($0 < r < 1$) and $\mu > 0$,

$$\int_0^{2\pi} |D_z^\lambda f(z)|^\mu d\theta \leq \int_0^{2\pi} |D_z^\lambda g_k(z)|^\mu d\theta.$$

References

- [1] G.S. SĂLĂGEAN , Subclasses of univalent functions, *Complex analysis - Proc. 5th Rom.-Finn. Semin.*, Bucharest 1981, Part 1, *Lect. Notes Math.*, **1013** (1983), 362–372.
- [2] H. SILVERMAN, Univalent functions with negative coefficients, *Proc. Amer. Math. Soc.*, **51**(1) (1975), 109–116.
- [3] S. SHAMS, S.R. KULKARNI AND J.M. JAHANGIRI, Classes of uniformly starlike and convex functions *Internat. J. Math. Math. Sci.*, **2004** (2004), Issue 55, 2959–2961.
- [4] S. SÜMER EKER AND S. OWA, New applications of classes of analytic functions involving the Sălăgean Operator, *Proceedings of the International Symposium on Complex Function Theory and Applications*, Transilvania University of Braşov Printing House, Braşov, Romania, 2006, 21–34.
- [5] J.E. LITTLEWOOD, On inequalities in the theory of functions, *Proc. London Math. Soc.*, **23** (1925), 481–519.
- [6] S. OWA, On the distortion theorems I, *Kyungpook Math. J.*, **18** (1978), 53–59.
- [7] H.M. SRIVASTAVA AND S. OWA (Eds.), *Univalent Functions, Fractional Calculus, and Their Applications*, Halsted Press (Ellis Horwood Limited, Chichester) John Wiley and Sons, New York, Chichester, Brisbane and Toronto, 1989



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 25 of 25

[Go Back](#)

[Full Screen](#)

[Close](#)