



ON THE CYCLIC HOMOGENEOUS POLYNOMIAL INEQUALITIES OF DEGREE FOUR

VASILE CIRTOAJE

Department of Automatic Control and Computers
University of Ploiesti
Romania
EMail: vcirtoaje@upg-ploiesti.ro

Received: 20 April, 2009

Accepted: 01 July, 2009

Communicated by: N.K. Govil

2000 AMS Sub. Class.: 26D05.

Key words: Cyclic inequality, Symmetric inequality, Necessity and sufficiency, Homogeneous polynomial of degree four.

Abstract: Let $f(x, y, z)$ be a cyclic homogeneous polynomial of degree four with three variables which satisfies $f(1, 1, 1) = 0$. In this paper, we give the necessary and sufficient conditions to have $f(x, y, z) \geq 0$ for any real numbers x, y, z . We also give the necessary and sufficient conditions to have $f(x, y, z) \geq 0$ for the case when f is symmetric and x, y, z are nonnegative real numbers. Finally, some new inequalities with cyclic homogeneous polynomials of degree four are presented.



Cyclic Homogeneous Polynomial
Inequalities

Vasile Cirtoaje

vol. 10, iss. 3, art. 67, 2009

[Title Page](#)

[Contents](#)



Page 1 of 22

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Contents

| | | |
|----------|-----------------------------------|-----------|
| 1 | Introduction | 3 |
| 2 | Main Results | 5 |
| 3 | Proof of Theorem 2.1 | 12 |
| 4 | Proof of Theorem 2.6 | 17 |
| 5 | Other Related Inequalities | 19 |



**Cyclic Homogeneous Polynomial
Inequalities**

Vasile Cirtoaje

vol. 10, iss. 3, art. 67, 2009

Title Page

Contents



Page 2 of 22


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 3 of 22

Go Back

Full Screen

Close

1. Introduction

Let x, y, z be real numbers. The fourth degree Schur's inequality ([3], [5], [7]) is a well-known symmetric homogeneous polynomial inequality which states that

$$(1.1) \quad \sum x^4 + xyz \sum x \geq \sum xy(x^2 + y^2),$$

where \sum denotes a cyclic sum over x, y and z . Equality holds for $x = y = z$, and for $x = 0$ and $y = z$, or $y = 0$ and $z = x$, or $z = 0$ and $x = y$.

In [3], the following symmetric homogeneous polynomial inequality was proved

$$(1.2) \quad \sum x^4 + 8 \sum x^2 y^2 \geq 3 \left(\sum xy \right) \left(\sum x^2 \right),$$

with equality for $x = y = z$, and for $x/2 = y = z$, or $y/2 = z = x$, or $z/2 = x = y$. In addition, a more general inequality was proved in [3] for any real k ,

$$(1.3) \quad \sum (x - y)(x - ky)(x - z)(x - kz) \geq 0,$$

with equality for $x = y = z$, and again for $x/k = y = z$, or $y/k = z = x$, or $z/k = x = y$. Notice that this inequality is a consequence of the identity

$$\sum (x - y)(x - ky)(x - z)(x - kz) = \frac{1}{2} \sum (y - z)^2 (y + z - x - kx)^2.$$

In 1992, we established the following cyclic homogeneous inequality [1]:

$$(1.4) \quad \left(\sum x^2 \right)^2 \geq 3 \sum x^3 y,$$

which holds for any real numbers x, y, z , with equality for $x = y = z$, and for

$$\frac{x}{\sin^2 \frac{4\pi}{7}} = \frac{y}{\sin^2 \frac{2\pi}{7}} = \frac{z}{\sin^2 \frac{\pi}{7}}$$



Title Page

Contents



Page 4 of 22

Go Back

Full Screen

Close

or any cyclic permutation thereof.

Six years later, we established a similar cyclic homogeneous inequality [2],

$$(1.5) \quad \sum x^4 + \sum xy^3 \geq 2 \sum x^3y,$$

which holds for any real numbers x, y, z , with equality for $x = y = z$, and for

$$x \sin \frac{\pi}{9} = y \sin \frac{7\pi}{9} = z \sin \frac{13\pi}{9}$$

or any cyclic permutation thereof.

As shown in [3], substituting $y = x + p$ and $z = x + q$, the inequalities (1.4) and (1.5) can be rewritten in the form

$$(p^2 - pq + q^2)x^2 + f(p, q)x + g(p, q) \geq 0,$$

where the quadratic polynomial of x has the discriminant

$$\delta_1 = -3(p^3 - p^2q - 2pq^2 + q^3)^2 \leq 0,$$


and, respectively,

$$\delta_2 = -3(p^3 - 3pq^2 + q^3)^2 \leq 0.$$

The symmetric inequalities (1.1), (1.2) and (1.3), as well as the cyclic inequalities (1.4) and (1.5), are particular cases of the inequality $f(x, y, z) \geq 0$, where $f(x, y, z)$ is a cyclic homogeneous polynomial of degree four satisfying $f(1, 1, 1) = 0$. This polynomial has the general form

$$(1.6) \quad f(x, y, z) = w \sum x^4 + r \sum x^2y^2 + (p + q - r - w)xyz \sum x - p \sum x^3y - q \sum xy^3,$$

where p, q, r, w are real numbers. Since the inequality $f(x, y, z) \geq 0$ with $w \leq 0$ does not hold for all real numbers x, y, z , except the trivial case where $w = p = q = 0$ and $r \geq 0$, we will consider $w = 1$ throughout this paper.



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 5 of 22

[Go Back](#)

[Full Screen](#)

[Close](#)

2. Main Results

In 2008, we posted, without proof, the following theorem in the Mathlinks Forum [4].

Theorem 2.1. *Let p, q, r be real numbers. The cyclic inequality*

$$(2.1) \quad \sum x^4 + r \sum x^2 y^2 + (p + q - r - 1)xyz \sum x \geq p \sum x^3 y + q \sum xy^3$$

holds for any real numbers x, y, z if and only if

$$(2.2) \quad 3(1 + r) \geq p^2 + pq + q^2.$$

For $p = q = 1$ and $r = 0$, we obtain the fourth degree Schur's inequality (1.1). For $p = q = 3$ and $r = 8$ one gets (1.2), while for $p = q = k + 1$ and $r = k(k + 2)$ one obtains (1.3). In addition, for $p = 3, q = 0$ and $r = 2$ one gets (1.4), while for $p = 2, q = -1$ and $r = 0$ one obtains (1.5).

In the particular cases $r = 0, r = p + q - 1, q = 0$ and $p = q$, by Theorem 2.1, we have the following corollaries, respectively.

Corollary 2.2. *Let p and q be real numbers. The cyclic inequality*

$$(2.3) \quad \sum x^4 + (p + q - 1)xyz \sum x \geq p \sum x^3 y + q \sum xy^3$$

holds for any real numbers x, y, z if and only if

$$(2.4) \quad p^2 + pq + q^2 \leq 3.$$

Corollary 2.3. *Let p and q be real numbers. The cyclic inequality*

$$(2.5) \quad \sum x^4 + (p + q - 1) \sum x^2 y^2 \geq p \sum x^3 y + q \sum xy^3$$



Title Page

Contents



Page 6 of 22

Go Back

Full Screen

Close

holds for any real numbers x, y, z if and only if

$$(2.6) \quad 3(p + q) \geq p^2 + pq + q^2.$$

Corollary 2.4. Let p and q be real numbers. The cyclic inequality

$$(2.7) \quad \sum x^4 + r \sum x^2 y^2 + (p - r - 1)xyz \sum x \geq p \sum x^3 y$$

holds for any real numbers x, y, z if and only if

$$(2.8) \quad 3(1 + r) \geq p^2.$$

Corollary 2.5. Let p and q be real numbers. The symmetric inequality

$$(2.9) \quad \sum x^4 + r \sum x^2 y^2 + (2p - r - 1)xyz \sum x \geq p \sum xy(x^2 + y^2)$$


holds for any real numbers x, y, z if and only if

$$(2.10) \quad r \geq p^2 - 1.$$

Finding necessary and sufficient conditions such that the cyclic inequality (2.1) holds for any nonnegative real numbers x, y, z is a very difficult problem. On the other hand, the approach for nonnegative real numbers is less difficult in the case when the cyclic inequality (2.1) is symmetric. Thus, in 2008, Le Huu Dien Khue posted, without proof, the following theorem on the Mathlinks Forum [4].



Theorem 2.6. Let p and r be real numbers. The symmetric inequality (2.9) holds for any nonnegative real numbers x, y, z if and only if

$$(2.11) \quad r \geq (p - 1) \max\{2, p + 1\}.$$



Title Page

Contents



Page 7 of 22

Go Back

Full Screen

Close

From Theorem 2.1, setting $p = 1 + \sqrt{6}$, $q = 1 - \sqrt{6}$ and $r = 2$, and then $p = 3$, $q = -3$ and $r = 2$, we obtain the inequalities:

$$(2.12) \quad \left(\sum x^2\right) \left(\sum x^2 - \sum xy\right) \geq \sqrt{6} \left(\sum x^3y - \sum xy^3\right),$$

with equality for $x = y = z$, and for $x = y/\alpha = z/\beta$ or any cyclic permutation, where $\alpha \approx 0.4493$ and $\beta \approx -0.1009$ were found using a computer;

$$(2.13) \quad (x^2 + y^2 + z^2)^2 \geq 3 \sum xy(x^2 - y^2 + z^2),$$

with equality for $x = y = z$, and for $x = y/\alpha = z/\beta$ or any cyclic permutation, where $\alpha \approx 0.2469$ and $\beta \approx -0.3570$.

From Corollary 2.2, setting $p = \sqrt{3}$ and $q = -\sqrt{3}$ yields


$$(2.14) \quad \sum x^4 - xyz \sum x \geq \sqrt{3} \left(\sum x^3y - \sum xy^3\right),$$

with equality for $x = y = z$, and for $x = y/\alpha = z/\beta$ or any cyclic permutation, where $\alpha \approx 0.3767$ and $\beta \approx -0.5327$. Notice that if x, y, z are nonnegative real numbers, then the best constant in inequality (2.14) is $2\sqrt{2}$ (Problem 19, Section 2.3 in [3], by Pham Kim Hung):

$$(2.15) \quad \sum x^4 - xyz \sum x \geq 2\sqrt{2} \left(\sum x^3y - \sum xy^3\right).$$



From Corollary 2.3, setting $p = 1 + \sqrt{3}$ and $q = 1$, and then $p = 1 - \sqrt{3}$ and $q = 1$, we obtain the inequalities:

$$(2.16) \quad \sum x^4 - \sum xy^3 \geq (1 + \sqrt{3}) \left(\sum x^3y - \sum x^2y^2\right),$$



Title Page

Contents



Page 8 of 22

Go Back

Full Screen

Close

with equality for $x = y = z$, and for $x = y/\alpha = z/\beta$ or any cyclic permutation, where $\alpha \approx 0.7760$ and $\beta \approx 0.5274$;

$$(2.17) \quad \sum x^4 - \sum xy^3 \geq (\sqrt{3} - 1) \left(\sum x^2y^2 - \sum x^3y \right),$$

with equality for $x = y = z$, and for $x = y/\alpha = z/\beta$ or any cyclic permutation, where $\alpha \approx 1.631$ and $\beta \approx -1.065$.

From Corollary 2.4, setting in succession $p = \sqrt{3}$ and $r = 0$, $p = -\sqrt{3}$ and $r = 0$, $p = 6$ and $r = 11$, $p = 2$ and $r = 1/3$, $p = -1$ and $r = -2/3$, $p = r = (3 + \sqrt{21})/2$, $p = 1$ and $r = -2/3$, $p = r = (3 - \sqrt{21})/2$, $p = \sqrt{6}$ and $r = 1$, we obtain the inequalities below, respectively:

$$(2.18) \quad \sum x^4 + (\sqrt{3} - 1)xyz \sum x \geq \sqrt{3} \sum x^3y,$$

with equality for $x = y = z$, and for $x = y/\alpha = z/\beta$ or any cyclic permutation, where $\alpha \approx 0.7349$ and $\beta \approx -0.1336$ (Problem 5.3.10 in [6]);

$$(2.19) \quad \sum x^4 + \sqrt{3} \sum x^3y \geq (1 + \sqrt{3})xyz \sum x,$$

with equality for $x = y = z$, and for $x = y/\alpha = z/\beta$ or any cyclic permutation, where $\alpha \approx 7.915$ and $\beta \approx -6.668$;

$$(2.20) \quad \sum x^4 + 11 \sum x^2y^2 \geq 6 \left(\sum x^3y + xyz \sum x \right),$$

with equality for $x = y = z$, and for $x = y/\alpha = z/\beta$ or any cyclic permutation, where $\alpha \approx 0.5330$ and $\beta \approx 2.637$;

$$(2.21) \quad 3 \sum x^4 + \left(\sum xy \right)^2 \geq 6 \sum x^3y,$$



[Title Page](#)

[Contents](#)



Page 9 of 22

[Go Back](#)

[Full Screen](#)

[Close](#)

with equality for $x = y = z$, and for $x = y/\alpha = z/\beta$ or any cyclic permutation, where $\alpha \approx 0.7156$ and $\beta \approx -0.0390$;

$$(2.22) \quad \sum x^4 + \sum x^3y \geq \frac{2}{3} \left(\sum xy \right)^2,$$

with equality for $x = y = z$, and for $x = y/\alpha = z/\beta$ or any cyclic permutation, where $\alpha \approx 1.871$ and $\beta \approx -2.053$;

$$(2.23) \quad \sum x^4 - xyz \sum x \geq \frac{3 + \sqrt{21}}{2} \left(\sum x^3y - \sum x^2y^2 \right),$$

with equality for $x = y = z$, and for $x = y/\alpha = z/\beta$ or any cyclic permutation, where $\alpha \approx 0.570$ and $\beta \approx 0.255$;

$$(2.24) \quad \sum x^4 - \sum x^3y \geq \frac{2}{3} \left(\sum x^2y^2 - xyz \sum x \right),$$

with equality for $x = y = z$, and for $x = y/\alpha = z/\beta$ or any cyclic permutation, where $\alpha \approx 0.8020$ and $\beta \approx -0.4446$;

$$(2.25) \quad \sum x^4 - xyz \sum x \geq \frac{\sqrt{21} - 3}{2} \left(\sum x^2y^2 - \sum x^3y \right),$$

with equality for $x = y = z$, and for $x = y/\alpha = z/\beta$ or any cyclic permutation, where $\alpha \approx 1.528$ and $\beta \approx -1.718$;

$$(2.26) \quad \sum (x^2 - yz)^2 \geq \sqrt{6} \sum xy(x - z)^2,$$

with equality for $x = y = z$, and for $x = y/\alpha = z/\beta$ or any cyclic permutation, where $\alpha \approx 0.6845$ and $\beta \approx 0.0918$ (Problem 21, Section 2.3 in [3]).



Title Page

Contents



Page 10 of 22

Go Back

Full Screen

Close

From either Corollary 2.5 or Theorem 2.6, setting $r = p^2 - 1$ yields

$$(2.27) \quad \sum x^4 + (p^2 - 1) \sum x^2 y^2 + p(2 - p)xyz \sum x \geq p \sum xy(x^2 + y^2),$$

which holds for any real numbers p and x, y, z . For $p = k + 1$, the inequality (2.27) turns into (1.3).

Corollary 2.7. *Let x, y, z be real numbers. If p, q, r, s are real numbers such that*

$$(2.28) \quad p + q - r - 1 \leq s \leq 2(r + 1) + p + q - p^2 - pq - q^2,$$

then

$$(2.29) \quad \sum x^4 + r \sum x^2 y^2 + sxyz \sum x \geq p \sum x^3 y + q \sum xy^3.$$

Let

$$\alpha = \frac{r + s + 1 - p - q}{3} \geq 0.$$

Since

$$3(1 + r - \alpha) \geq p^2 + pq + q^2,$$

by Theorem 2.1 we have

$$\sum x^4 + (r - \alpha) \sum x^2 y^2 + (\alpha + p + q - r - 1)xyz \sum x \geq p \sum x^3 y + q \sum xy^3.$$

Adding this inequality to the obvious inequality

$$\alpha \left(\sum xy \right)^2 \geq 0,$$

we get (2.29).



Title Page

Contents



Page 11 of 22

Go Back

Full Screen

Close

From Corollary 2.7, setting $p = 1$, $q = r = 0$ and $s = 2$, we get

$$(2.30) \quad \sum x^4 + 2xyz \sum x \geq \sum x^3y,$$

with equality for $x = y/\alpha = z/\beta$ or any cyclic permutation, where $\alpha \approx 0.8020$ and $\beta \approx -0.4451$. Notice that (2.30) is equivalent to

$$(2.31) \quad \sum (2x^2 - y^2 - z^2 - xy + yz)^2 + 4 \left(\sum xy \right)^2 \geq 0.$$



Title Page

Contents



Page 12 of 22

Go Back

Full Screen

Close

3. Proof of Theorem 2.1

Proof of the Sufficiency. Since

$$\sum x^2y^2 - xyx \sum x = \frac{1}{2} \sum x^2(y-z)^2 \geq 0,$$

it suffices to prove the inequality (2.1) for the least value of r , that is

$$r = \frac{p^2 + pq + q^2}{3} - 1.$$

On this assumption, (2.1) is equivalent to each of the following inequalities:

$$(3.1) \quad \sum [2x^2 - y^2 - z^2 - pxy + (p+q)yz - qzx]^2 \geq 0,$$

$$(3.2) \quad \sum [3y^2 - 3z^2 - (p+2q)xy - (p-q)yz + (2p+q)zx]^2 \geq 0,$$

$$(3.3) \quad 3[2x^2 - y^2 - z^2 - pxy + (p+q)yz - qzx]^2 \\ + [3y^2 - 3z^2 - (p+2q)xy - (p-q)yz + (2p+q)zx]^2 \geq 0.$$

Thus, the conclusion follows. \square

Proof of the Necessity. For $p = q = 2$, we need to show that the condition $r \geq 3$ is necessary to have

$$\sum x^4 + r \sum x^2y^2 + (3-r)xyz \sum x \geq 2 \sum x^3y + 2 \sum xy^3$$



Title Page

Contents



Page 13 of 22

Go Back

Full Screen

Close

for any real numbers x, y, z . Indeed, setting $y = z = 1$ reduces this inequality to

$$(x - 1)^4 + (r - 3)(x - 1)^2 \geq 0,$$

which holds for any real x if and only if $r \geq 3$.

In the other cases (different from $p = q = 2$), by Lemma 3.1 below it follows that there is a triple $(a, b, c) = (1, b, c) \neq (1, 1, 1)$ such that

$$\sum [2a^2 - b^2 - c^2 - pab + (p + q)bc - qca]^2 = 0.$$

Since

$$\sum a^2b^2 - abc \sum a = \frac{1}{2} \sum a^2(b - c)^2 > 0,$$

we may write this relation as

$$\frac{p \sum a^3b + q \sum ab^3 - \sum a^4 - (p + q - 1)abc \sum a}{\sum a^2b^2 - abc \sum a} = \frac{p^2 + pq + q^2}{3} - 1.$$

On the other hand, since (2.1) holds for (a, b, c) (by hypothesis), we get

$$r \geq \frac{p \sum a^3b + q \sum ab^3 - \sum a^4 - (p + q - 1)abc \sum a}{\sum a^2b^2 - abc \sum a}.$$

Therefore,

$$r \geq \frac{p^2 + pq + q^2}{3} - 1,$$

which is the desired necessary condition. \square

Lemma 3.1. *Let p and q be real numbers. Excepting the case $p = q = 2$, there is a real triple $(x, y, z) = (1, y, z) \neq (1, 1, 1)$ such that*

$$(3.4) \quad \sum [2x^2 - y^2 - z^2 - pxy + (p + q)yz - qzx]^2 = 0.$$



Title Page

Contents



Page 14 of 22

Go Back

Full Screen

Close

Proof. We consider two cases: $p = q \neq 2$ and $p \neq q$.

Case 1. $p = q \neq 2$.

It is easy to prove that $(x, y, z) = (1, p - 1, 1) \neq (1, 1, 1)$ is a solution of the equation (3.4).

Case 2. $p \neq q$.

The equation (3.4) is equivalent to

$$\begin{cases} 2y^2 - z^2 - x^2 - pyz + (p + q)zx - qxy = 0 \\ 2z^2 - x^2 - y^2 - pzx + (p + q)xy - qyz = 0. \end{cases}$$

For $x = 1$, we get

$$(3.5) \quad \begin{cases} 2y^2 - z^2 - 1 - pyz + (p + q)z - qy = 0 \\ 2z^2 - 1 - y^2 - pz + (p + q)y - qyz = 0. \end{cases}$$

Adding the first equation multiplied by 2 to the second equation yields

$$(3.6) \quad z[(2p + q)y - p - 2q] = 3y^2 + (p - q)y - 3.$$

Under the assumption that $(2p + q)y - p - 2q \neq 0$, substituting z from (3.6) into the first equation, (3.5) yields

$$(3.7) \quad (y - 1)(ay^3 + by^2 + cy - a) = 0,$$

where

$$\begin{aligned} a &= 9 - 2p^2 - 5pq - 2q^2, \\ b &= 9 + 6p - 6q - 3p^2 + 3q^2 + 2p^3 + 3p^2q + 3pq^2 + q^3, \\ c &= -9 + 6p - 6q - 3p^2 + 3q^2 - p^3 - 3p^2q - 3pq^2 - 2q^3. \end{aligned}$$



Title Page

Contents



Page 15 of 22

Go Back

Full Screen

Close

The equation (3.7) has a real root $y_1 \neq 1$. To prove this claim, it suffices to show that the equation $ay^3 + by^2 + cy - a = 0$ does not have a root of 1; that is to show that $b + c \neq 0$. This is true because

$$\begin{aligned} b + c &= 12(p - q) - 6(p^2 - q^2) + p^3 - q^3 \\ &= (p - q)(12 - 6p - 6q + p^2 + q^2 + pq), \end{aligned}$$

and

$$p - q \neq 0,$$

$$\begin{aligned} 4(12 - 6p - 6q + p^2 + q^2 + pq) &> 48 - 24(p + q) + 3(p + q)^2 \\ &= 3(p + q - 4)^2 \\ &\geq 0. \end{aligned}$$

For $y = y_1$ and $(2p + q)y_1 - p - 2q \neq 0$, from (3.6) we get

$$z_1 = \frac{3y_1^2 + (p - q)y_1 - 3}{(2p + q)y_1 - p - 2q},$$

and the conclusion follows. Thus, it remains to consider that $(2p + q)y_1 - p - 2q = 0$. In this case, we have $2p + q \neq 0$ (since $2p + q = 0$ provides $p + 2q = 0$, which contradicts the hypothesis $p \neq q$), and hence

$$y_1 = \frac{p + 2q}{2p + q}.$$

For $y = y_1$, from (3.6) we get $3(y_1^2 - 1) + (p - q)y_1 = 0$, which yields

$$(3.8) \quad (2p + q)(p + 2q) = 9(p + q).$$



Substituting y_1 into the first equation (3.5), we get

$$(2p + q)z^2 - (p^2 + q^2 + pq)z + p + 2q = 0.$$

To complete the proof, it suffices to show that this quadratic equation has real roots. Due to (3.8), we need to prove that

$$(p^2 + q^2 + pq)^2 \geq 36(p + q).$$

For the nontrivial case $p + q > 0$, let us denote $s = p + q$, $s > 0$, and write the condition (3.8) as $9s - 2s^2 = pq$. Since $4pq \leq s^2$, we find that $s \geq 4$. Therefore,

$$(p^2 + q^2 + pq)^2 - 36(p + q) = 9(s^2 - 3s)^2 - 36s = 9s(s - 1)^2(s - 4) \geq 0.$$

□

Title Page

Contents

◀◀ ▶▶


◀ ▶

Page 16 of 22

Go Back

Full Screen

Close



Title Page

Contents



Page 17 of 22

Go Back

Full Screen

Close

4. Proof of Theorem 2.6

The condition $r \geq (p - 1) \max\{2, p + 1\}$ is equivalent to $r \geq p^2 - 1$ for $p \geq 1$, and $r \geq 2(p - 1)$ for $p \leq 1$.

Proof of the Sufficiency. By Theorem 2.1, if $r \geq p^2 - 1$, then the inequality (2.9) is true for any real numbers x, y, z . Thus, it only remains to consider the case when $p \leq 1$ and $r \geq 2(p - 1)$. Writing (2.9) as

$$\begin{aligned} \sum x^4 + xyz \sum x - \sum xy(x^2 + y^2) \\ + (1 - p) \left[\sum xy(x^2 + y^2) - 2 \sum x^2 y^2 \right] \\ + (r - 2p + 2) \left(\sum x^2 y^2 - xyz \sum x \right) \geq 0, \end{aligned}$$

we see that it is true because

$$\sum x^4 + xyz \sum x - \sum xy(x^2 + y^2) \geq 0$$

(Schur's inequality of fourth degree),

$$\sum xy(x^2 + y^2) - 2 \sum x^2 y^2 = \sum xy(x - y)^2 \geq 0$$

and

$$\sum x^2 y^2 - xyz \sum x = \frac{1}{2} \sum x^2 (y - z)^2 \geq 0.$$

□

Proof of the Necessity. We need to prove that the conditions $r \geq 2(p - 1)$ and $r \geq p^2 - 1$ are necessary such that the inequality (2.9) holds for any nonnegative real



Title Page

Contents



Page 18 of 22

Go Back

Full Screen

Close

numbers x, y, z . Setting $y = z = 1$, (2.9) becomes

$$(x - 1)^2[x^2 + 2(1 - p)x + 2 + r - 2p] \geq 0.$$


For $x = 0$, we get the necessary condition $r \geq 2(p - 1)$, while for $x = p - 1$, we get

$$(p - 2)^2(r + 1 - p^2) \geq 0.$$

If $p \neq 2$, then this inequality provides the necessary condition $r \geq p^2 - 1$. Thus, it remains to show that for $p = 2$, we have the necessary condition $r \geq 3$. Indeed, setting $p = 2$ and $y = z = 1$ reduces the inequality (2.9) to

$$(x - 1)^2[(x - 1)^2 + r - 3] \geq 0.$$

Clearly, this inequality holds for any nonnegative x if and only if $r \geq 3$. □



5. Other Related Inequalities

The following theorem establishes other interesting related inequalities with symmetric homogeneous polynomials of degree four.

Theorem 5.1. *Let x, y, z be real numbers, and let*

$$A = \sum x^4 - \sum x^2y^2, \quad B = \sum x^2y^2 - xyz \sum x,$$
$$C = \sum x^3y - xyz \sum x, \quad D = \sum xy^3 - xyz \sum x.$$

Then,

$$(5.1) \quad AB = C^2 - CD + D^2 \geq \frac{C^2 + D^2}{2} \geq \left(\frac{C + D}{2}\right)^2 \geq CD.$$

Moreover, if x, y, z are nonnegative real numbers, then

$$(5.2) \quad CD \geq B^2.$$

The equality $AB = CD$ holds for $x + y + z = 0$, and for $x = y$, or $y = z$, or $z = x$, while the equality $CD = B^2$ holds for $x = y = z$, and for $x = 0$, or $y = 0$, or $z = 0$.

Proof. The inequalities in Theorem 5.1 follow from the identities:

$$D - C = (x + y + z)(x - y)(y - z)(z - x),$$
$$AB - CD = (x + y + z)^2(x - y)^2(y - z)^2(z - x)^2,$$
$$AB - \left(\frac{C + D}{2}\right)^2 = \frac{3}{4}(x + y + z)^2(x - y)^2(y - z)^2(z - x)^2,$$

Title Page

Contents

◀◀ ▶▶


◀ ▶

Page 19 of 22

Go Back

Full Screen

Close



Title Page

Contents



Page 20 of 22

Go Back

Full Screen

Close

$$AB - \frac{C^2 + D^2}{2} = \frac{1}{2}(x + y + z)^2(x - y)^2(y - z)^2(z - x)^2,$$
$$CD - B^2 = xyz(x + y + z)(x^2 + y^2 + z^2 - xy - yz - zx)^2.$$

□

Remark 1. We obtained the identity $AB = C^2 - CD + D^2$ in the following way. For $3(r + 1) = p^2 + pq + q^2$, by Theorem 2.1 we have

$$A + (1 + r)B - pC - qD \geq 0,$$

which is equivalent to

$$Bp^2 + (Bq - 3C)p + Bq^2 - 3Dq + 3A \geq 0.$$

Since this inequality holds for any real p and $B \geq 0$, the discriminant of the quadratic of p is non-positive; that is

$$(Bq - 3C)^2 - 4B(Bq^2 - 3Dq + 3A) \leq 0,$$

which is equivalent to

$$B^2q^2 + 2B(C - 2D)q + 4AB - 3C^2 \geq 0.$$

Similarly, the discriminant of the quadratic of q is non-positive; that is

$$B^2(C - 2D)^2 - B^2(4AB - 3C^2) \leq 0,$$

which yields $AB \geq C^2 - CD + D^2$. Actually, this inequality is an identity.

Remark 2. The inequality $CD \geq B^2$ is true if

$$k^2C - 2kB + D \geq 0$$



for any real k . This inequality is equivalent to

$$\sum yz(x - ky)^2 \geq (k - 1)^2xyz \sum x,$$

which follows immediately from the Cauchy-Schwarz inequality

$$\left(\sum x\right) \left[\sum yz(x - ky)^2\right] \geq (k - 1)^2xyz \left(\sum x\right)^2.$$

On the other hand, assuming that $x = \min\{x, y, z\}$ and substituting $y = x + p$ and $z = x + q$, where $p, q \geq 0$, the inequality $CD \geq B^2$ can be rewritten as

$$A_1x^4 + B_1x^3 + C_1x^2 + D_1x \geq 0,$$

with

$$A_1 = 3(p^2 - pq + q^2)^2 \geq 0,$$

$$B_1 = 4(p + q)(p^2 - pq + q^2)^2 \geq 0,$$

$$C_1 = 2pq(p^2 - pq + q^2)^2 + pq(p^2 - q^2)^2 + (p^3 + q^3)^2 - 2p^2q^2(p^2 + q^2) + 5p^3q^3 \geq 0,$$

$$D_1 = pq[p^5 + q^5 - pq(p^3 + q^3) + p^2q^2(p + q)] \geq 0.$$

Title Page

Contents



Page 21 of 22

Go Back

Full Screen

Close

References

- [1] V. CIRTOAJE, Problem 22694, *Gazeta Matematica*, **7-8** (1992), 287.
- [2] V. CIRTOAJE, Problem O:887, *Gazeta Matematica*, **10** (1998), 434.
- [3] V. CIRTOAJE, *Algebraic Inequalities-Old and New Methods*, GIL Publishing House, 2006.
- [4] V. CIRTOAJE AND LE HUU DIEN KHUE, *Mathlinks Forum*, February 2008, [ONLINE: <http://www.mathlinks.ro/Forum/viewtopic.php?t=186179>].
- [5] G.H. HARDY, J.E. LITTLEWOOD AND G. POLYA, *Inequalities*, Cambridge University Press, 1952.
- [6] P.K. HUNG, *Secrets in Inequalities*, Vol. 2, GIL Publishing House, 2008.
- [7] D.S. MITRINOVIĆ, J. PEČARIĆ AND A.M. FINK, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht/Boston/London, 1993.



Cyclic Homogeneous Polynomial
Inequalities

Vasile Cirtoaje

vol. 10, iss. 3, art. 67, 2009

Title Page

Contents



Page 22 of 22

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756