



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 1 of 17

[Go Back](#)

[Full Screen](#)

[Close](#)

A NOTE ON ESTIMATES OF DIAGONAL ELEMENTS OF THE INVERSE OF DIAGONALLY DOMINANT TRIDIAGONAL MATRICES

TIZIANO POLITI AND MARINA POPOLIZIO

Dipartimento di Matematica

Politecnico di Bari

Via Orabona 4, I-70125, Bari (Italy)

EEmail: politi@poliba.it popolizio@dm.uniba.it

URL: <http://www.poliba.it> <http://www.dm.uniba.it/~popolizio>

Received: 06 May, 2007

Accepted: 15 March, 2008

Communicated by: S.S. Dragomir


2000 AMS Sub. Class.: 65F50, 15A39.

Key words: Tridiagonal matrices, Bounds of entries of the inverse.

Abstract: In this note we show how to improve some recent upper and lower bounds for the elements of the inverse of diagonally dominant tridiagonal matrices. In particular, a technique described by [R. Peluso, and T. Politi, Some improvements on two-sided bounds on the inverse of diagonally dominant tridiagonal matrices, Lin. Alg. Appl. Vol. 330 (2001) 1-14], is used to obtain better bounds for the diagonal elements.

Contents

1	Introduction	3
2	Some Known Results	5
2.1	Special cases	8
3	Lower and Upper Bounds for the Diagonal Elements	10
4	Numerical Examples	14



**Estimates of Diagonal
Elements**

Tiziano Politi and
Marina Popolizio

vol. 9, iss. 2, art. 31, 2008

[Title Page](#)

[Contents](#)



Page 2 of 17


[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 4 of 17

[Go Back](#)

[Full Screen](#)

[Close](#)

with $a_i \neq 0$, for $i = 1, \dots, n$; assume that A is *row diagonally dominant*, i.e.

$$|a_i| \geq |b_i| + |c_{i-1}|, \quad i = 1, \dots, n, \quad c_0 = b_n = 0.$$

It is natural to suppose that $c_i, b_i \neq 0$, for $i = 1, \dots, n-1$, as if one of these elements is 0 the problem can be reduced to smaller subproblems, see e.g. [1].

Let $C = A^{-1} = \{c_{i,j}\}$ be the inverse of A .

In the following we will use some relations derived from the identity $AC = I$; indeed, for the diagonal elements of C it results that


$$(1.1) \quad c_{j-1}c_{j-1,j} + a_j c_{j,j} + b_j c_{j+1,j} = 1, \quad j = 2, \dots, n-1$$

while for $j \geq 3$ and $i = 1, \dots, j$ the j -th equation of the system $AC = I$ reads

$$(1.2) \quad c_{i,j} = -\frac{a_{i-1}c_{i-1,j} + c_{i-2}c_{i-2,j}}{b_{i-1}}.$$

If $b_i < 0$, $c_i < 0$ and $a_i > 0$, and $a_1 > -b_1$ or $a_n > -c_{n-1}$, then A is an M -matrix. In this case $c_{i,j} > 0$, $\forall i, j$.

The paper is organized as follows: in Section 2 we recall a collection of known results, while in Section 3 we exploit some results described in Section 2 to improve the bounds for diagonal entries, and in Section 4 some numerical examples are shown.



2. Some Known Results

The technique common to [6], [7] and [4] is bounding $|c_{i,j}|$, $i \neq j$, as a function of diagonal elements $|c_{j,j}|$ and providing suitable bounds for these.

For a review of the most recent resulting bounds presented in [4], we define the quantities

$$\xi_i = \frac{|b_i|}{|a_i| - |\alpha_{i-1}| |c_{i-1}|}; \quad m_i = \frac{|b_i|}{|a_i| + |\alpha_{i-1}| |c_{i-1}|};$$

$$\lambda_i = \frac{|c_{i-1}|}{|a_i| - |\beta_{i+1}| |b_i|}; \quad n_i = \frac{|c_{i-1}|}{|a_i| + |\beta_{i+1}| |b_i|},$$

where

$$\alpha_i = \frac{b_i}{p_i}, \quad i = 1, \dots, n-1$$

$$p_i = a_i - \alpha_{i-1} c_{i-1}, \quad q_i = \beta_{i+1} b_i, \quad i = 1, \dots, n$$

$$\beta_i = \frac{c_{i-1}}{a_i - q_i}, \quad i = n, \dots, 2,$$


provided that $\alpha_0 = \beta_{n+1} = 0$.

Using these quantities the following bounds were derived in [4].

Theorem 2.1 ([4]). *If $|a_i| - |\beta_{i+1}| |b_i| > 0$ for $i = 2, \dots, n$, and $|a_i| - |\alpha_{i-1}| |c_{i-1}| > 0$ for $i = 1, \dots, n-1$, then the following bounds hold for the elements of C :*

$$(2.1) \quad n_i |c_{i-1,j}| \leq |c_{i,j}| \leq \lambda_i |c_{i-1,j}|, \quad i = j+1, \dots, n, \quad j = 1, \dots, n-1,$$

$$(2.2) \quad m_i |c_{i+1,j}| \leq |c_{i,j}| \leq \xi_i |c_{i+1,j}|, \quad i = 1, \dots, j-1, \quad j = 2, \dots, n,$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 6 of 17

Go Back

Full Screen

Close

$$(2.3) \quad |c_{j,j}| \prod_{k=j+1}^i n_k \leq |c_{i,j}| \leq |c_{j,j}| \prod_{k=j+1}^i \lambda_k, \quad \text{for } i > j,$$

$$(2.4) \quad |c_{j,j}| \prod_{k=i}^{j-1} m_k \leq |c_{i,j}| \leq |c_{j,j}| \prod_{k=i}^{j-1} \xi_k, \quad \text{for } i < j,$$

while for the diagonal entries

$$(2.5) \quad \frac{1}{|a_j| + \xi_{j-1}|c_{j-1}| + \lambda_{j+1}|b_j|} \leq |c_{j,j}| \leq \frac{1}{|a_j| - \xi_{j-1}|c_{j-1}| - \lambda_{j+1}|b_j|}.$$

Remark 1. The hypotheses of Theorem 2.1 hold when A is a strictly diagonally dominant or an irreducibly diagonally dominant matrix [4].

We recall the results presented in [7], for which we need the following definitions:

$$\tau_i = \frac{|b_i|}{|a_i| - |c_{i-1}|} \quad \delta_i = \frac{|b_i|}{|a_i| + |c_{i-1}|} \quad i = 1, \dots, n-1,$$

$$\omega_i = \frac{|c_{i-1}|}{|a_i| - |b_i|} \quad \gamma_i = \frac{|c_{i-1}|}{|a_i| + |b_i|} \quad i = 2, \dots, n,$$

$$s_j = \begin{cases} \tau_{j-1} & \text{if } a_{j-1}a_j b_{j-1}c_{j-1} < 0; \\ -\delta_{j-1} & \text{if } a_{j-1}a_j b_{j-1}c_{j-1} > 0, \end{cases}$$

$$t_j = \begin{cases} \omega_{j+1} & \text{if } a_{j+1}a_j b_j c_j < 0; \\ -\gamma_{j+1} & \text{if } a_{j+1}a_j b_j c_j > 0, \end{cases}$$

$$f_j = \begin{cases} -\tau_{j-1} & \text{if } a_{j-1}a_jb_{j-1}c_{j-1} > 0; \\ \delta_{j-1} & \text{if } a_{j-1}a_jb_{j-1}c_{j-1} < 0, \end{cases}$$

$$g_j = \begin{cases} -\omega_{j+1} & \text{if } a_{j+1}a_jb_jc_j > 0; \\ \gamma_{j+1} & \text{if } a_{j+1}a_jb_jc_j < 0. \end{cases}$$

Theorem 2.2 ([7]). *Let A be a nonsingular tridiagonal matrix. If A is row diagonally dominant, then the following bounds hold for the elements of C :*

$$(2.6) \quad |c_{j,j}| \prod_{k=j+1}^i \delta_k \leq |c_{i,j}| \leq |c_{j,j}| \prod_{k=j+1}^i \tau_k, \quad \text{for } i > j;$$

$$(2.7) \quad |c_{j,j}| \prod_{k=i}^{j-1} \gamma_k \leq |c_{i,j}| \leq |c_{j,j}| \prod_{k=i}^{j-1} \omega_k, \quad \text{for } i < j,$$

while for the diagonal entries

$$(2.8) \quad \frac{1}{|a_j| + s_j|c_{j-1}| + t_j|b_j|} \leq |c_{j,j}| \leq \frac{1}{|a_j| + f_j|c_{j-1}| + g_j|b_j|}.$$

Remark 2. There are few remarks for both the theorems recalled above, which have not been stressed so far.

From both theorems it is clear that for the off-diagonal elements we have $|c_{i,j}| \leq |c_{i-1,j}|$, since all the coefficients involved are less than 1; this leads to the well known result about the decreasing pattern of the elements of the inverse of a banded matrix, see e.g. [3].



Title Page

Contents

◀◀ ▶▶


◀ ▶

Page 7 of 17

Go Back

Full Screen

Close



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 8 of 17

[Go Back](#)

[Full Screen](#)

[Close](#)

From the lower bounds for the diagonal entries it is clear that $c_{i,i} \geq \frac{1}{a_{i,i}}$, which was a lower bound presented by Meurant [5] in the special case of diagonally dominant M -matrices, and so the estimates in (2.8) are more accurate than those in [5], although they are more costly to evaluate.

Remark 3. For $i < j$, the following inequalities hold (see [4]):

$$(2.9) \quad \begin{aligned} m_k &\geq \delta_k, & k &= i, \dots, j-1; \\ \xi_k &\leq \tau_k, & k &= i, \dots, j-1, \end{aligned}$$

and, for $i > j$:

$$(2.10) \quad \begin{aligned} n_k &\geq \gamma_k, & k &= j+1, \dots, i; \\ \lambda_k &\leq \omega_k, & k &= j+1, \dots, i, \end{aligned}$$

hence the bounds for the extradiagonal elements reported in Theorem 2.1 improve those given in 2.2.

2.1. Special cases

In the extensive survey about the inverse of tridiagonal matrices [5], Meurant considered also the special case in which one is interested in the inverse C of a *Toeplitz* matrix T_a , having a on the main diagonal and -1 on the other two diagonals. The provided upper bounds share the common idea of [4] and [7], that is $|c_{i,j}|$, $i \neq j$, is bounded as a function of $|c_{j,j}|$ and suitable bounds are presented for the diagonal entries.

We now recall that result:

Theorem 2.3. [5] *If $a > 2$ then*

$$(2.11) \quad (T_a^{-1})_{i,j} < r_-^{j-i} (T_a^{-1})_{i,i}, \quad \forall i, \forall j \geq i$$

$$(2.12) \quad (T_a^{-1})_{i,j} < \frac{r_-^{j-i+1}}{1-r}, \quad \forall i, \forall j \geq i+1$$

$$\text{for } r_{\pm} = \frac{a \pm \sqrt{(a^2-4)}}{2}; r = \frac{r_-}{r_+}.$$



**Estimates of Diagonal
Elements**

Tiziano Politi and
Marina Popolizio

vol. 9, iss. 2, art. 31, 2008

Title Page

Contents



Page 9 of 17


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



3. Lower and Upper Bounds for the Diagonal Elements

The following example motivated our investigation. We consider the tridiagonal matrix

$$A = \begin{bmatrix} 4 & 2 & & & \\ -2 & 4 & 2 & & \\ & 2 & 4 & 2 & \\ & & -2 & -4 & -2 \\ & & & -2 & 4 \end{bmatrix}$$


for which the hypotheses of Theorem 2.1 are true, and compare the bounds given by [7] and by [4]: to consider the bounds of [4], we construct the matrices U and L whose (i, j) entry represents the upper, respectively the lower, bound in (2.3) and in (2.4), whereas the (j, j) entry is the upper, respectively the lower, bound in (2.5). We use the same notation for representing the bounds given in [7].

In Table 1 the maximum errors on the upper and lower bounds are represented. From Table 1 we observe that the bounds by Liu et al. have a maximum error greater

	$\max_{i,j}\{U_{i,j} - c_{i,j} \}$	$\max_{i,j}\{ c_{i,j} - L_{i,j}\}$
(2.6),(2.7),(2.8)	0.4706	0.0846
(2.3),(2.4),(2.5)	0.6366	0.1213
	$\max_j\{U_{j,j} - c_{j,j} \}$	$\max_j\{ c_{j,j} - L_{j,j}\}$
(2.8)	0.1912	0.0353
(2.5)	0.6366	0.1213

Table 1:

than that by Peluso et al.; the reason is that the bounds derived in [4] do not improve those on the diagonal elements hence the global bounds are not better for all the elements of the matrix. Looking inside the bound matrices we observe that the



Title Page

Contents



Page 11 of 17

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

bounds improve for 40% of the entries and for 20% of diagonal elements when using (2.3), (2.4) and (2.5) in place of (2.3), (2.4), (3.1).

Our aim is to obtain sharper two-sided bounds for the diagonal elements of C , exploiting the signs of its entries, using exactly the same technique as in [7]. We now give the main result of the present paper.

Theorem 3.1. *Let A be a nonsingular tridiagonal matrix and $C = A^{-1}$. If A is row diagonally dominant then*

$$(3.1) \quad \frac{1}{|a_j| + s_j|c_{j-1}| + t_j|b_j|} \leq |c_{j,j}| \leq \frac{1}{|a_j| + f_j|c_{j-1}| + g_j|b_j|}, \quad j = 1, \dots, n,$$


where

$$s_j = \begin{cases} \xi_{j-1} & \text{if } a_{j-1}a_jb_{j-1}c_{j-1} < 0; \\ -m_{j-1} & \text{if } a_{j-1}a_jb_{j-1}c_{j-1} > 0, \end{cases}$$

$$t_j = \begin{cases} \lambda_{j+1} & \text{if } a_{j+1}a_jb_jc_j < 0; \\ -n_{j+1} & \text{if } a_{j+1}a_jb_jc_j > 0, \end{cases}$$

$$f_j = \begin{cases} -\xi_{j-1} & \text{if } a_{j-1}a_jb_{j-1}c_{j-1} > 0; \\ m_{j-1} & \text{if } a_{j-1}a_jb_{j-1}c_{j-1} < 0, \end{cases}$$

$$g_j = \begin{cases} -\lambda_{j+1} & \text{if } a_{j+1}a_jb_jc_j > 0; \\ n_{j+1} & \text{if } a_{j+1}a_jb_jc_j < 0, \end{cases}$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 12 of 17

Go Back

Full Screen

Close

Proof. We fix j and define $\mu_j = c_{j-1,j}/c_{j,j}$ and $\rho_j = c_{j+1,j}/c_{j,j}$; if we use these definitions in (1.1) we get

$$c_{j,j} = \frac{1}{a_j + \mu_j c_{j-1} + \rho_j b_j} = \frac{1}{a_j \left(1 + \mu_j \frac{c_{j-1}}{a_j} + \rho_j \frac{b_j}{a_j}\right)}$$

and

$$(3.2) \quad |c_{j,j}| = \frac{1}{|a_j| \left(1 + \varepsilon_1 |\mu_j| \frac{|c_{j-1}|}{|a_j|} + \varepsilon_2 |\rho_j| \frac{|b_j|}{|a_j|}\right)}$$

with $\varepsilon_1 = \text{sign}(\mu_j c_{j-1} a_j)$ and $\varepsilon_2 = \text{sign}(\rho_j b_j a_j)$.

For $i = j - 1$, (2.2) gives

$$m_{j-1} |c_{j,j}| \leq |c_{j-1,j}| \leq \xi_{j-1} |c_{j,j}|,$$

that is

$$(3.3) \quad m_{j-1} \leq |\mu_j| \leq \xi_{j-1}.$$

If $\varepsilon_1 > 0$, then $m_{j-1} |c_{j-1}| \leq \varepsilon_1 |\mu_j| |c_{j-1}| \leq \xi_{j-1} |c_{j-1}|$, while if $\varepsilon_1 < 0$, then $-\xi_{j-1} |c_{j-1}| \leq \varepsilon_1 |\mu_j| |c_{j-1}| \leq -m_{j-1} |c_{j-1}|$ and from these relations we get the expression for s_j .


It remains to show that $\varepsilon_1 = \text{sign}(a_j a_{j-1} b_{j-1} c_{j-1})$.

The equation (1.2) for $i = j$ gives

$$\frac{c_{j,j}}{c_{j-1,j}} = -\frac{a_{j-1}}{b_{j-1}} - \frac{c_{j-2}}{b_{j-1}} \frac{c_{j-2,j}}{c_{j-1,j}}.$$

Since $|c_{j-2,j}| \leq |c_{j-1,j}|$ and $|a_{j-1}| > |c_{j-2}|$, then

$$-\text{sign}(a_{j-1} b_{j-1}) = \text{sign}(c_{j,j} c_{j-1,j}) = \text{sign}(\mu_j)$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 13 of 17

Go Back

Full Screen

Close

from which we have the expression for ε_1 .

Using exactly the same technique, we may work on the upper bounds and discuss the influence of ε_2 , as in [7]. □


Again, as in [7], we can write explicit bounds when A is an M -matrix.

Remark 4. If A is an M -matrix, then the bounds proved in Theorem 3.1 become

$$\frac{1}{a_j - m_{j-1}|c_{j-1}| - n_{j+1}|b_j|} \leq c_{j,j} \leq \frac{1}{a_j - \xi_{j-1}|c_{j-1}| - \lambda_{j+1}|b_j|}, \quad j = 1, \dots, n.$$

We note that the upper bound is the same as that in [4] but the lower bound is sharper.

Remark 5. As a consequence of (2.9) and (2.10), the bounds proved in Theorem 3.1 are sharper (or not worse) than those shown in [4].



	$\max_{i,j}\{U_{i,j} - c_{i,j} \}$	$\max_{i,j}\{ c_{i,j} - L_{i,j}\}$
(2.3), (2.4), (3.1)	$2.4119e - 003$	$4.0993e - 004$
(2.3), (2.4), (2.5)	$2.1600e - 001$	$1.0290e - 001$
	$\max_j\{U_{j,j} - c_{j,j} \}$	$\max_j\{ c_{j,j} - L_{j,j}\}$
(2.3), (2.4), (3.1)	$1.7895e - 004$	$2.4253e - 005$
(2.3), (2.4), (2.5)	$2.1600e - 001$	$4.9282e - 003$

Table 2: Comparison with the Liu et al.'s upper bounds


trices is

$$T_4 = \begin{bmatrix} 4 & -1 & & & \\ -1 & 4 & -1 & & \\ & -1 & 4 & -1 & \\ & & \ddots & \ddots & \\ & & & -1 & 4 \end{bmatrix};$$

it arises when discretizing the Laplace operator on a rectangular domain with Dirichlet boundary conditions: it actually represents the structure of the diagonal blocks of the discretization of the Laplacian obtained by applying a five-point finite difference.

In [2], Benzi and Golub used this matrix to test an approximate inverse; they stress that an approximate inverse of T_4 is needed in the initial step of an incomplete Cholesky factorization for the two-dimensional model problem. By applying the equations (2.3), (2.4), (3.1) we are able to obtain good bounds for the generic entry of T_4^{-1} . We define the matrices U and L as in the previous example and, with a very naïve idea, we consider the matrix $M := \frac{1}{2}(U + L)$ as an approximant to T_4^{-1} ; we use a matrix T_4 of dimension 100×100 .

As in [2], we use the condition number cond of the matrix MT_4 to test the quality



of the approximation. We notice that $\text{cond}(MT_4) = 1.0736$, from which we may state that M would be an effective preconditioner.

As mentioned in [2], it is well known that, due to the decay of the elements of the inverse, considering only the tridiagonal approximations provides good results. We considered only the tridiagonal part of M , say M_1 , to obtain $\text{cond}(M_1T_4) = 1.2185$.

With the same matrix T_4 , the upper bounds in (2.3) and (2.4) are sharp when we use (3.1) for the diagonal entries. We in fact observed that

$$\max_{i,j} \{U_{i,j} - |c_{i,j}|\} < 10^{-16},$$

where U_{ij} and c_{ij} represent, respectively, the (i, j) entry of matrices U , of the upper bounds (2.3), (2.4) and (3.1), and T_4^{-1} .

Example 4.3. We consider the Toeplitz matrix T_4 of dimension 8×8 ; we compare the upper bounds presented in this paper with those proposed by Meurant [5].

	$\max_{i,j \geq i} \{U_{i,j} - c_{i,j} \}$
(2.3), (2.4), (3.1)	$8.6736e - 019$
(2.11)	$1.2780e - 007$
	$\max_{i,j \geq i+1} \{U_{i,j} - c_{i,j} \}$
(2.3), (2.4), (3.1)	$2.8709e - 017$
(2.12)	$5.7654e - 003$

Table 3: Comparison with the Meurant's upper bounds

The results in Table 3 show an impressive improvement when using the bounds (2.3), (2.4), (3.1).

References

- [1] W.W. BARRETT, A theorem on inverses of tridiagonal matrices, *Lin. Alg. Appl.*, **27** (1979), 211–217.
- [2] M. BENZI AND G.H. GOLUB. Bounds for the entries of matrix functions with applications to preconditioning, *BIT*, **39**(3) (1999), 417–438.
- [3] P. CONCUS, G.H. GOLUB AND G. MEURANT, Block preconditioning for the conjugate gradient methods, *SIAM J. Sci. Stat. Comput.*, **6** (1985), 220–252.
- [4] X.Q. LIU, T.Z. HUANG AND Y.D. FU, Estimates for the inverse elements of tridiagonal matrices, *Appl. Math. Lett.*, **19** (2006), 590–598.
- [5] G. MEURANT, A review on the inverse of symmetric tridiagonal and block tridiagonal matrices, *SIAM J. Matrix Anal. Appl.*, **13**(3) (1992), 707–728.
- [6] R. NABBEN, Two-sided bounds on the inverse of diagonally dominant tridiagonal matrices, *Lin. Alg. Appl.*, **287** (1999), 289–305.
- [7] R. PELUSO AND T. POLITI, Some improvements on two-sided bounds on the inverse of diagonally dominant tridiagonal matrices, *Lin. Alg. Appl.*, **330** (2001), 1–14.
- [8] P.N. SHIVAKUMAR AND CHUANXIANG JI, Upper and lower bounds for inverse elements of finite and infinite tridiagonal matrices, *Lin. Alg. Appl.*, **247** (1996), 297–316.





Estimates of Diagonal
Elements

Tiziano Politi and
Marina Popolizio

vol. 9, iss. 2, art. 31, 2008

Title Page

Contents



Page 17 of 17

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756