


# Journal of Inequalities in Pure and Applied Mathematics

## ON THE HYERS-ULAM STABILITY OF QUADRATIC FUNCTIONAL EQUATIONS

ICK-SOON CHANG AND HARK-MAHN KIM

Department of Mathematics,  
Chungnam National University,  
Taejeon 305-764, Korea  
*EMail:* [ischang@math.cnu.ac.kr](mailto:ischang@math.cnu.ac.kr)  
*EMail:* [hmkim@math.cnu.ac.kr](mailto:hmkim@math.cnu.ac.kr)



---

volume 3, issue 3, article 33,  
2002.

*Received 11 January, 2002;*  
*accepted 19 February, 2002.*

*Communicated by: Th. M. Rassias*

---

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

## Abstract

In this paper, we obtain the general solution and the generalized Hyers-Ulam stability for quadratic functional equations  $f(2x + y) + f(2x - y) = f(x + y) + f(x - y) + 6f(x)$  and  $f(2x + y) + f(x + 2y) = 4f(x + y) + f(x) + f(y)$ .

*2000 Mathematics Subject Classification:* 39B22, 39B52, 39B72.

*Key words:* Hyers-Ulam-Rassias stability; Quadratic function

## Contents

1	Introduction .....	3
2	Solution of (1.3), (1.4) .....	6
3	Stability of (1.3) .....	9
4	Stability of (1.4) .....	19

References



---

### An Inequality which Arises in the Absence of the Mountain Pass Geometry

Ick-Soon Chang and  
Hark-Mahn Kim

---

Title Page

Contents



Go Back

Close

Quit

Page 2 of 28

# 1. Introduction

In 1940, S.M. Ulam [20] gave a wide ranging talk before the mathematics club of the University of Wisconsin in which he discussed a number of important unsolved problems. Among those was the question concerning the stability of homomorphisms:


*Let  $G_1$  be a group and let  $G_2$  be a metric group with the metric  $d(\cdot, \cdot)$ . Given  $\epsilon > 0$ , does there exist a  $\delta > 0$  such that if a function  $h : G_1 \rightarrow G_2$  satisfies the inequality  $d(h(xy), h(x)h(y)) < \delta$  for all  $x, y \in G_1$ , then there exists a homomorphism  $H : G_1 \rightarrow G_2$  with  $d(h(x), H(x)) < \epsilon$  for all  $x \in G_1$ ?*

In other words, we are looking for situations when the homomorphisms are stable, i.e., if a mapping is almost a homomorphism, then there exists a true homomorphism near it. If we turn our attention to the case of functional equations, we can ask the question: When the solutions of an equation differing slightly from a given one must be close to the true solution of the given equation.

The case of approximately additive functions was solved by D. H. Hyers [9] under the assumption that  $G_1$  and  $G_2$  are Banach spaces. In 1978, a generalized version of the theorem of Hyers for approximately linear mappings was given by Th. M. Rassias [17]. During the last decades, the stability problems of several functional equations have been extensively investigated by a number of authors [2, 6, 11, 15]. The terminology generalized Hyers-Ulam stability originates from these historical backgrounds. These terminologies are also applied to the case of other functional equations. For more detailed definitions of such terminologies, we can refer to [10, 12, 18].

The functional equation

$$(1.1) \quad f(x + y) + f(x - y) = 2f(x) + 2f(y)$$



---

**An Inequality which Arises in  
the Absence of the Mountain  
Pass Geometry**

Ick-Soon Chang and  
Hark-Mahn Kim

---

Title Page

Contents



Go Back

Close

Quit

Page 3 of 28

is related to a symmetric biadditive function ([1], [16]). It is natural that each equation is called a quadratic functional equation. In particular, every solution of the quadratic equation (1.1) is said to be a quadratic function. It is well known that a function  $f$  between real vector spaces is quadratic if and only if there exists a unique symmetric biadditive function  $B$  such that  $f(x) = B(x, x)$  for all  $x$  (see [1], [16]). The biadditive function  $B$  is given by

$$(1.2) \quad B(x, y) = \frac{1}{4}(f(x + y) - f(x - y)).$$


A Hyers-Ulam stability problem for the quadratic functional equation (1.1) was proved by F. Skof for functions  $f : E_1 \rightarrow E_2$ , where  $E_1$  is a normed space and  $E_2$  a Banach space (see [19]). P. W. Cholewa [3] noticed that the theorem of Skof is still true if the relevant domain  $E_1$  is replaced by an abelian group. In the paper [4], S. Czerwik proved the Hyers-Ulam-Rassias stability of the quadratic functional equation (1.1). A. Grabiec [8] has generalized these results mentioned above. K. W. Jun and Y. H. Lee [13] proved the Hyers-Ulam-Rassias stability of the pexiderized quadratic equation (1.1).

Now, we introduce the following functional equations, which are somewhat different from (1.1),

$$(1.3) \quad f(2x + y) + f(2x - y) = f(x + y) + f(x - y) + 6f(x),$$

$$(1.4) \quad f(2x + y) + f(x + 2y) = 4f(x + y) + f(x) + f(y).$$

In this paper, we establish the general solution and the generalized Hyers-Ulam stability problem for the equations (1.3), (1.4), which are equivalent to (1.1). It is significant for us to decrease the possible estimator of the stability




---

**An Inequality which Arises in the Absence of the Mountain Pass Geometry**

Ick-Soon Chang and  
Hark-Mahn Kim

---

Title Page

Contents




Go Back

Close

Quit

Page 4 of 28

problem for the functional equations. This work is possible if we consider the stability problem in the sense of Hyers-Ulam-Rassias for the functional equations (1.3), (1.4). As a result, we have much better possible upper bounds for the equations (1.3), (1.4) than those of Czerwik [4] and Skof-Cholewa [3].



---

**An Inequality which Arises in  
the Absence of the Mountain  
Pass Geometry**

Ick-Soon Chang and  
Hark-Mahn Kim

---

Title Page

Contents



Go Back

Close

Quit

Page 5 of 28

## 2. Solution of (1.3), (1.4)

Let  $\mathbb{R}^+$  denote the set of all nonnegative real numbers and let both  $E_1$  and  $E_2$  be real vector spaces. We here present the general solution of (1.3), (1.4).


**Theorem 2.1.** *A function  $f : E_1 \rightarrow E_2$  satisfies the functional equation (1.1) if and only if  $f : E_1 \rightarrow E_2$  satisfies the functional equation (1.4) if and only if  $f : E_1 \rightarrow E_2$  satisfies the functional equation (1.3). Therefore, every solution of functional equations (1.3) and (1.4) is also a quadratic function.*

*Proof.* Let  $f : E_1 \rightarrow E_2$  satisfy the functional equation (1.1). Putting  $x = 0 = y$  in (1.1), we get  $f(0) = 0$ . Set  $x = 0$  in (1.1) to get  $f(y) = f(-y)$ . Letting  $y = x$  and  $y = 2x$  in (1.1), respectively, we obtain that  $f(2x) = 4f(x)$  and  $f(3x) = 9f(x)$  for all  $x \in E_1$ . By induction, we lead to  $f(kx) = k^2 f(x)$  for all positive integer  $k$ . Replacing  $x$  and  $y$  by  $2x + y$  and  $x + 2y$  in (1.1), respectively, we have

$$\begin{aligned}(2.1) \quad f(2x + y) + f(x + 2y) &= \frac{1}{2}[f(3x + 3y) + f(x - y)] \\ &= 4f(x + y) + \frac{1}{2}[f(x + y) + f(x - y)] \\ &= 4f(x + y) + f(x) + f(y)\end{aligned}$$

for all  $x, y \in E_1$ .

Let  $f : E_1 \rightarrow E_2$  satisfy the functional equation (1.4). Putting  $x = 0 = y$  in (1.4), we get  $f(0) = 0$ . Set  $y = 0$  in (1.4) to get  $f(2x) = 4f(x)$ . Letting  $y = x$  and  $y = -2x$  in (1.4), we obtain that  $f(3x) = 9f(x)$  and  $f(x) = f(-x)$  for all



---

An Inequality which Arises in  
the Absence of the Mountain  
Pass Geometry

Ick-Soon Chang and  
Hark-Mahn Kim

---

Title Page

Contents



Go Back

Close

Quit

Page 6 of 28

$x \in E_1$ . Putting  $x$  and  $y$  by  $x + y$  and  $x + y$  in (1.4), respectively, we obtain

$$(2.2) \quad f(2x + 3y) + f(x + 3y) = 4f(x + 2y) + f(x + y) + f(y),$$

$$(2.3) \quad f(3x + y) + f(3x + 2y) = 4f(2x + y) + f(x) + f(x + y).$$

Adding (2.2) to (2.3) and using (1.4), we obtain

$$(2.4) \quad f(2x + 3y) + f(3x + 2y) + f(x + 3y) + f(3x + y) \\ = 18f(x + y) + 5f(x) + 5f(y)$$

for all  $x, y \in E_1$ . Replacing  $y$  by  $2y$  and  $x$  by  $2x$  in (1.4), respectively, we have

$$(2.5) \quad 4f(x + y) + f(x + 4y) = 4f(x + 2y) + f(x) + 4f(y),$$

$$(2.6) \quad 4f(x + y) + f(4x + y) = 4f(2x + y) + 4f(x) + f(y)$$

for all  $x, y \in E_1$ . Adding (2.5) to (2.6) and using (1.4), we get

$$(2.7) \quad f(x + 4y) + f(4x + y) = 8f(x + y) + 9f(x) + 9f(y)$$


for all  $x, y \in E_1$ .

On the other hand, using (1.4), we get

$$(2.8) \quad f(x + 4y) + f(4x + y) \\ = f(6x + 9y) + f(9x + 6y) - 4f(5x + 5y) \\ = 9f(2x + 3y) + 9f(3x + 2y) - 100f(x + y),$$

which yields the relation by virtue of (2.7)

$$(2.9) \quad f(2x + 3y) + f(3x + 2y) = 12f(x + y) + f(x) + f(y)$$



An Inequality which Arises in  
the Absence of the Mountain  
Pass Geometry

Ick-Soon Chang and  
Hark-Mahn Kim

Title Page

Contents



Go Back

Close

Quit

Page 7 of 28

for all  $x, y \in E_1$ . Combining the last equation with (2.4), we get

$$(2.10) \quad f(x + 3y) + f(3x + y) = 6f(x + y) + 4f(x) + 4f(y).$$

Replacing  $x$  and  $y$  by  $\frac{x+y}{2}$  and  $\frac{x-y}{2}$  in (2.10), respectively, we have the desired result (1.3).

Now, let  $f : E_1 \rightarrow E_2$  satisfy the functional equation (1.3). Putting  $x = 0 = y$  in (1.3), we get  $f(0) = 0$ . Letting  $y = 0$  and  $y = x$  in (1.3), respectively, we obtain that  $f(2x) = 4f(x)$  and  $f(3x) = 9f(x)$  for all  $x \in E_1$ . Putting  $y = 2x$  in (1.3), we get  $f(x) = f(-x)$ . Replacing  $x$  and  $y$  by  $x + y$  and  $x - y$ , respectively, in (1.3), we have

$$(2.11) \quad f(3x + y) + f(x + 3y) = 6f(x + y) + 4f(x) + 4f(y)$$

for all  $x, y \in E_1$ . Replacing  $y$  by  $x + y$  in (1.3), we obtain

$$(2.12) \quad f(3x + y) + f(x - y) = 6f(x) + f(2x + y) + f(y).$$

Interchange  $x$  with  $y$  in (2.12) to get the relation

$$(2.13) \quad f(3y + x) + f(x - y) = 6f(y) + f(2y + x) + f(x).$$


Adding (2.12) to (2.13), we obtain

$$(2.14) \quad 6f(x + y) + 2f(x - y) = f(2x + y) + f(x + 2y) + 3f(x) + 3f(y)$$

for all  $x, y \in E_1$ . Setting  $-y$  instead of  $y$  in (2.14) and using the evenness of  $f$ , we get the relation

$$(2.15) \quad 6f(x - y) + 2f(x + y) = f(2x - y) + f(2y - x) + 3f(x) + 3f(y).$$

Adding (2.14) to (2.15), we obtain the result (1.1). □



**An Inequality which Arises in  
the Absence of the Mountain  
Pass Geometry**

Ick-Soon Chang and  
Hark-Mahn Kim

Title Page

Contents



Go Back

Close

Quit

Page 8 of 28



### 3. Stability of (1.3)

From now on, let  $X$  be a real vector space and let  $Y$  be a Banach space unless we give any specific reference. We will investigate the Hyers-Ulam-Rassias stability problem for the functional equation (1.3). Thus we find the condition that there exists a true quadratic function near an approximately quadratic function.

**Theorem 3.1.** *Let  $\phi : X^2 \rightarrow \mathbb{R}^+$  be a function such that*

$$(3.1) \quad \sum_{i=0}^{\infty} \frac{\phi(2^i x, 0)}{4^i} \quad \left( \sum_{i=1}^{\infty} 4^i \phi\left(\frac{x}{2^i}, 0\right), \text{ respectively} \right)$$

*converges and*

$$(3.2) \quad \lim_{n \rightarrow \infty} \frac{\phi(2^n x, 2^n y)}{4^n} = 0 \quad \left( \lim_{n \rightarrow \infty} 4^n \phi\left(\frac{x}{2^n}, \frac{y}{2^n}\right) = 0 \right)$$


*for all  $x, y \in X$ . Suppose that a function  $f : X \rightarrow Y$  satisfies*

$$(3.3) \quad \|f(2x + y) + f(2x - y) - f(x + y) - f(x - y) - 6f(x)\| \leq \phi(x, y)$$

*for all  $x, y \in X$ . Then there exists a unique quadratic function  $T : X \rightarrow Y$  which satisfies the equation (1.3) and the inequality*

$$(3.4) \quad \|f(x) - T(x)\| \leq \frac{1}{8} \sum_{i=0}^{\infty} \frac{\phi(2^i x, 0)}{4^i}$$

$$\left( \|f(x) - T(x)\| \leq \frac{1}{8} \sum_{i=1}^{\infty} 4^i \phi\left(\frac{x}{2^i}, 0\right) \right)$$



An Inequality which Arises in the Absence of the Mountain Pass Geometry

Ick-Soon Chang and  
Hark-Mahn Kim

Title Page

Contents



Go Back

Close

Quit

Page 9 of 28

for all  $x \in X$ . The function  $T$  is given by

$$(3.5) \quad T(x) = \lim_{n \rightarrow \infty} \frac{f(2^n x)}{4^n} \quad \left( T(x) = \lim_{n \rightarrow \infty} 4^n f\left(\frac{x}{2^n}\right) \right)$$

for all  $x \in X$ .

*Proof.* Putting  $y = 0$  in (3.3) and dividing by 8, we have

$$(3.6) \quad \left\| \frac{f(2x)}{4} - f(x) \right\| \leq \frac{1}{8} \phi(x, 0)$$


for all  $x \in X$ . Replacing  $x$  by  $2x$  in (3.6) and dividing by 4 and summing the resulting inequality with (3.6), we get

$$(3.7) \quad \left\| \frac{f(2^2 x)}{4^2} - f(x) \right\| \leq \frac{1}{8} \left[ \phi(x, 0) + \frac{\phi(2x, 0)}{4} \right]$$

for all  $x \in X$ . Using the induction on a positive integer  $n$ , we obtain that

$$(3.8) \quad \begin{aligned} \left\| \frac{f(2^n x)}{4^n} - f(x) \right\| &\leq \frac{1}{8} \sum_{i=0}^{n-1} \frac{\phi(2^i x, 0)}{4^i} \\ &\leq \frac{1}{8} \sum_{i=0}^{\infty} \frac{\phi(2^i x, 0)}{4^i} \end{aligned}$$

for all  $x \in X$ . In order to prove convergence of the sequence  $\left\{ \frac{f(2^n x)}{4^n} \right\}$ , we



**An Inequality which Arises in  
the Absence of the Mountain  
Pass Geometry**

Ick-Soon Chang and  
Hark-Mahn Kim

Title Page

Contents



Go Back

Close

Quit

Page 10 of 28

divide inequality (3.8) by  $4^m$  and also replace  $x$  by  $2^m x$  to find that for  $n, m > 0$ ,

$$(3.9) \quad \left\| \frac{f(2^n 2^m x)}{4^{n+m}} - \frac{f(2^m x)}{4^m} \right\| = \frac{1}{4^m} \left\| \frac{f(2^n 2^m x)}{4^n} - f(2^m x) \right\|$$

$$\leq \frac{1}{8 \cdot 4^m} \sum_{i=0}^{n-1} \frac{\phi(2^i 2^m x, 0)}{4^i}$$

$$\leq \frac{1}{8} \sum_{i=0}^{\infty} \frac{\phi(2^i 2^m x, 0)}{4^{m+i}}.$$


Since the right hand side of the inequality tends to 0 as  $m$  tends to infinity, the sequence  $\left\{ \frac{f(2^n x)}{4^n} \right\}$  is a Cauchy sequence. Therefore, we may define  $T(x) = \lim_{n \rightarrow \infty} 2^{-2n} f(2^n x)$  for all  $x \in X$ . By letting  $n \rightarrow \infty$  in (3.8), we arrive at the formula (3.4). To show that  $T$  satisfies the equation (1.3), replace  $x, y$  by  $2^n x, 2^n y$ , respectively, in (3.3) and divide by  $4^n$ , then it follows that

$$4^{-n} \|f(2^n(2x + y)) + f(2^n(2x - y)) - f(2^n(x + y)) - f(2^n(x - y)) - 6f(2^n x)\| \leq 4^{-n} \phi(2^n x, 2^n y).$$

Taking the limit as  $n \rightarrow \infty$ , we find that  $T$  satisfies (1.3) for all  $x, y \in X$ .

To prove the uniqueness of the quadratic function  $T$  subject to (3.4), let us assume that there exists a quadratic function  $S : X \rightarrow Y$  which satisfies (1.3) and the inequality (3.4). Obviously, we have  $S(2^n x) = 4^n S(x)$  and  $T(2^n x) = 4^n T(x)$  for all  $x \in X$  and  $n \in \mathbb{N}$ . Hence it follows from (3.4) that

$$\|S(x) - T(x)\| = 4^{-n} \|S(2^n x) - T(2^n x)\|$$



An Inequality which Arises in  
the Absence of the Mountain  
Pass Geometry

Ick-Soon Chang and  
Hark-Mahn Kim

Title Page

Contents



Go Back

Close

Quit

Page 11 of 28

$$\begin{aligned} &\leq 4^{-n}(\|S(2^n x) - f(2^n x)\| + \|f(2^n x) - T(2^n x)\|) \\ &\leq \frac{1}{4} \sum_{i=0}^{\infty} \frac{\phi(2^i 2^n x, 0)}{4^{n+i}} \end{aligned}$$

for all  $x \in X$ . By letting  $n \rightarrow \infty$  in the preceding inequality, we immediately find the uniqueness of  $T$ . This completes the proof of the theorem.  $\square$

Throughout this paper, let  $B$  be a unital Banach algebra with norm  $|\cdot|$ , and let  ${}_B\mathbb{B}_1$  and  ${}_B\mathbb{B}_2$  be left Banach  $B$ -modules with norms  $\|\cdot\|$  and  $\|\cdot\|$ , respectively. A quadratic mapping  $Q : {}_B\mathbb{B}_1 \rightarrow {}_B\mathbb{B}_2$  is called  $B$ -quadratic if


$$Q(ax) = a^2 Q(x), \quad \forall a \in B, \forall x \in {}_B\mathbb{B}_1.$$

**Corollary 3.2.** *Let  $\phi : {}_B\mathbb{B}_1 \times {}_B\mathbb{B}_1 \rightarrow \mathbb{R}^+$  be a function satisfying (3.1) and (3.2) for all  $x, y \in {}_B\mathbb{B}_1$ . Suppose that a mapping  $f : {}_B\mathbb{B}_1 \rightarrow {}_B\mathbb{B}_2$  satisfies*

$$\begin{aligned} &\|f(2\alpha x + \alpha y) + f(2\alpha x - \alpha y) - \alpha^2 f(x + y) - \alpha^2 f(x - y) - 6\alpha^2 f(x)\| \\ &\leq \phi(x, y) \end{aligned}$$

for all  $\alpha \in B$  ( $|\alpha| = 1$ ) and for all  $x, y \in {}_B\mathbb{B}_1$ , and  $f$  is measurable or  $f(tx)$  is continuous in  $t \in \mathbb{R}$  for each fixed  $x \in {}_B\mathbb{B}_1$ . Then there exists a unique  $B$ -quadratic mapping  $T : {}_B\mathbb{B}_1 \rightarrow {}_B\mathbb{B}_2$ , defined by (3.5), which satisfies the equation (1.3) and the inequality (3.4) for all  $x \in {}_B\mathbb{B}_1$ .

*Proof.* By Theorem 3.1, it follows from the inequality of the statement for  $\alpha = 1$  that there exists a unique quadratic mapping  $T : {}_B\mathbb{B}_1 \rightarrow {}_B\mathbb{B}_2$  satisfying the inequality (3.4) for all  $x \in {}_B\mathbb{B}_1$ . Under the assumption that  $f$  is measurable or



An Inequality which Arises in the Absence of the Mountain Pass Geometry

Ick-Soon Chang and  
Hark-Mahn Kim

Title Page

Contents



Go Back

Close

Quit

Page 12 of 28

$f(tx)$  is continuous in  $t \in \mathbb{R}$  for each fixed  $x \in {}_B\mathbb{B}_1$ , by the same reasoning as the proof of [5], the quadratic mapping  $T : {}_B\mathbb{B}_1 \rightarrow {}_B\mathbb{B}_2$  satisfies

$$T(tx) = t^2T(x), \quad \forall x \in {}_B\mathbb{B}_1, \forall t \in \mathbb{R}.$$

That is,  $T$  is  $\mathbb{R}$ -quadratic. For each fixed  $\alpha \in B$  ( $|\alpha| = 1$ ), replacing  $f$  by  $T$  and setting  $y = 0$  in (1.3), we have  $T(\alpha x) = \alpha^2T(x)$  for all  $x \in {}_B\mathbb{B}_1$ . The last relation is also true for  $\alpha = 0$ . For each element  $a \in B$  ( $a \neq 0$ ),  $a = |a| \cdot \frac{a}{|a|}$ . Since  $T$  is  $\mathbb{R}$ -quadratic and  $T(\alpha x) = \alpha^2T(x)$  for each element  $\alpha \in B$  ( $|\alpha| = 1$ ),


$$\begin{aligned} T(ax) &= T\left(|a| \cdot \frac{a}{|a|}x\right) \\ &= |a|^2 \cdot T\left(\frac{a}{|a|}x\right) \\ &= |a|^2 \cdot \frac{a^2}{|a|^2} \cdot T(x) \\ &= a^2T(x), \quad \forall a \in B(a \neq 0), \quad \forall x \in {}_B\mathbb{B}_1. \end{aligned}$$

So the unique  $\mathbb{R}$ -quadratic mapping  $T : {}_B\mathbb{B}_1 \rightarrow {}_B\mathbb{B}_2$  is also  $B$ -quadratic, as desired. This completes the proof of the corollary.  $\square$

Since  $\mathbb{C}$  is a Banach algebra, the Banach spaces  $E_1$  and  $E_2$  are considered as Banach modules over  $\mathbb{C}$ . Thus we have the following corollary.

**Corollary 3.3.** *Let  $E_1$  and  $E_2$  be Banach spaces over the complex field  $\mathbb{C}$ , and let  $\varepsilon \geq 0$  be a real number. Suppose that a mapping  $f : E_1 \rightarrow E_2$  satisfies*

$$\|f(2\alpha x + \alpha y) + f(2\alpha x - \alpha y) - \alpha^2 f(x + y) - \alpha^2 f(x - y) - 6\alpha^2 f(x)\| \leq \varepsilon$$




---

**An Inequality which Arises in the Absence of the Mountain Pass Geometry**

Ick-Soon Chang and  
Hark-Mahn Kim

---

Title Page

Contents



Go Back

Close

Quit

Page 13 of 28

for all  $\alpha \in \mathbb{C}$  ( $|\alpha| = 1$ ) and for all  $x, y \in E_1$ , and  $f$  is measurable or  $f(tx)$  is continuous in  $t \in \mathbb{R}$  for each fixed  $x \in E_1$ . Then there exists a unique  $\mathbb{C}$ -quadratic mapping  $T : E_1 \rightarrow E_2$  which satisfies the equation (1.3) and the inequality

$$\|f(x) - T(x)\| \leq \frac{\varepsilon}{6}$$

for all  $x \in E_1$ .


The S. Czerwik [4] theorem for the functional equation (1.1) states that if a function  $f : G \rightarrow Y$ , where  $G$  is an abelian group and  $Y$  a Banach space, satisfies the inequality  $\|f(x+y) + f(x-y) - 2f(x) - 2f(y)\| \leq \varepsilon(\|x\|^p + \|y\|^p)$  for  $p \neq 2$  and for all  $x, y \in G$ , then there exists a unique quadratic function  $q$  such that  $\|f(x) - q(x)\| \leq \frac{\varepsilon\|x\|^p}{|4-2^p|} + \frac{\|f(0)\|}{3}$  for all  $x \in G$ , and for all  $x \in G - \{0\}$  and  $\|f(0)\| = 0$  if  $p < 0$ . From the main theorem 3.1, we obtain the following corollary concerning the stability of the equation (1.3). We note that  $p$  need not be equal to  $q$ .

**Corollary 3.4.** *Let  $X$  and  $Y$  be a real normed space and a Banach space, respectively, and let  $\varepsilon, p, q$  be real numbers such that  $\varepsilon \geq 0, q > 0$  and either  $p, q < 2$  or  $p, q > 2$ . Suppose that a function  $f : X \rightarrow Y$  satisfies*

$$\|f(2x + y) + f(2x - y) - f(x + y) - f(x - y) - 6f(x)\| \leq \varepsilon(\|x\|^p + \|y\|^q)$$

for all  $x, y \in X$ . Then there exists a unique quadratic function  $T : X \rightarrow Y$  which satisfies the equation (1.3) and the inequality

$$\|f(x) - T(x)\| \leq \frac{\varepsilon}{2|4 - 2^p|} \|x\|^p$$



An Inequality which Arises in the Absence of the Mountain Pass Geometry

Ick-Soon Chang and  
Hark-Mahn Kim

Title Page

Contents




Go Back

Close

Quit

Page 14 of 28



**An Inequality which Arises in the Absence of the Mountain Pass Geometry**

Ick-Soon Chang and  
Hark-Mahn Kim

Title Page

Contents



Go Back

Close

Quit

Page 15 of 28

for all  $x \in X$  and for all  $x \in X - \{0\}$  if  $p < 0$ . The function  $T$  is given by

$$T(x) = \lim_{n \rightarrow \infty} \frac{f(2^n x)}{4^n} \text{ if } p, q < 2 \quad \left( T(x) = \lim_{n \rightarrow \infty} 4^n f\left(\frac{x}{2^n}\right) \text{ if } p, q > 2 \right)$$

for all  $x \in X$ . Further, if for each fixed  $x \in X$  the mapping  $t \mapsto f(tx)$  from  $\mathbb{R}$  to  $Y$  is continuous, then  $T(rx) = r^2 T(x)$  for all  $r \in \mathbb{R}$ .

The proof of the last assertion in the above corollary goes through in the same way as that of [4].

The Skof-Cholewa [3] theorem for the functional equation (1.1) states that if a function  $f : G \rightarrow Y$ , where  $G$  is an abelian group and  $Y$  a Banach space, satisfies the inequality  $\|f(x+y) + f(x-y) - 2f(x) - 2f(y)\| \leq \varepsilon$  for all  $x, y \in G$ , then there exists a unique quadratic function  $q$  such that  $\|f(x) - q(x)\| \leq \frac{\varepsilon}{2}$  for all  $x \in G$ . But we have a much better possible upper bound concerning the stability theorem for the functional equation (1.3) as follows. The following corollary is an immediate consequence of Theorem 3.1.

**Corollary 3.5.** *Let  $X$  and  $Y$  be a real normed space and a Banach space, respectively, and let  $\varepsilon \geq 0$  be a real number. Suppose that a function  $f : X \rightarrow Y$  satisfies*

$$(3.10) \quad \|f(2x+y) + f(2x-y) - f(x+y) - f(x-y) - 6f(x)\| \leq \varepsilon$$

for all  $x, y \in X$ . Then there exists a unique quadratic function  $T : X \rightarrow Y$  defined by  $T(x) = \lim_{n \rightarrow \infty} \frac{f(2^n x)}{4^n}$  which satisfies the equation (1.3) and the inequality

$$(3.11) \quad \|f(x) - T(x)\| \leq \frac{\varepsilon}{6}$$

for all  $x \in X$ . Further, if for each fixed  $x \in X$  the mapping  $t \mapsto f(tx)$  from  $\mathbb{R}$  to  $Y$  is continuous, then  $T(rx) = r^2T(x)$  for all  $r \in \mathbb{R}$ .

**Remark 3.1.** If we write  $y = x$  in the inequality of (3.3), we get

$$(3.12) \quad \|f(3x) - 5f(x) - f(2x)\| \leq \phi(x, x) + \|f(0)\|.$$

Combining (3.12) with (3.6), we have

$$(3.13) \quad \|f(3x) - 9f(x)\| \leq \phi(x, x) + \frac{\phi(x, 0)}{2} + \|f(0)\|.$$

We can easily show the following relation by induction on  $n$  together with (3.13)

$$\left\| \frac{f(3^n x)}{9^n} - f(x) \right\| \leq \frac{1}{9} \sum_{i=0}^{n-1} \frac{1}{9^i} \left[ \phi(3^i x, 3^i x) + \frac{\phi(3^i x, 0)}{2} + \|f(0)\| \right]$$

for all  $x \in X$ .


In Theorem 3.1, let  $\phi : X^2 \rightarrow \mathbb{R}^+$  be a function such that

$$\sum_{i=0}^{\infty} \frac{\phi(3^i x, 3^i x) + \phi(3^i x, 0)}{9^i} \quad \left( \sum_{i=1}^{\infty} 9^i \left[ \phi\left(\frac{x}{3^i}, \frac{x}{3^i}\right) + \phi\left(\frac{x}{3^i}, 0\right) \right], \text{ respectively} \right)$$

converges and

$$\lim_{n \rightarrow \infty} \frac{\phi(3^n x, 3^n y)}{9^n} = 0 \quad \left( \lim_{n \rightarrow \infty} 9^n \phi\left(\frac{x}{3^n}, \frac{y}{3^n}\right) = 0 \right)$$

for all  $x, y \in X$ . Note that in the second case  $f(0) = 0$  since  $\phi(0, 0) = 0$ . Then, using the last inequality and the same argument of Theorem 3.1, we can



An Inequality which Arises in  
the Absence of the Mountain  
Pass Geometry

Ick-Soon Chang and  
Hark-Mahn Kim

Title Page

Contents



Go Back

Close

Quit

Page 16 of 28



find the unique quadratic function  $T$  defined by  $T(x) = \lim_{n \rightarrow \infty} 3^{-2n} f(3^n x)$  which satisfies (1.3) and the inequality

$$(3.14) \quad \|f(x) - T(x)\| \leq \frac{1}{9} \sum_{i=0}^{\infty} \frac{1}{9^i} \left[ \phi(3^i x, 3^i x) + \frac{\phi(3^i x, 0)}{2} \right] + \frac{\|f(0)\|}{8}$$

$$\left( \|f(x) - T(x)\| \leq \frac{1}{9} \sum_{i=1}^{\infty} 9^i \left[ \phi\left(\frac{x}{3^i}, \frac{x}{3^i}\right) + \frac{\phi\left(\frac{x}{3^i}, 0\right)}{2} \right] \right)$$

for all  $x \in X$ . Thus we obtain an alternative result of Theorem 3.1. In Theorem 3.1, we have a simpler possible upper bound (3.4) than that of (3.14). The advantage of the inequality (3.4) compared to (3.14) is that the right hand side of (3.4) has no term for  $\|f(0)\|$ .


As a consequence of the above Remark 3.1, we have the following corollary. Because of the restricted condition  $0 < p$ , we have  $f(0) = 0$ .

**Corollary 3.6.** *Let  $X$  and  $Y$  be a real normed space and a Banach space, respectively, and let  $\varepsilon \geq 0$ ,  $0 < p \neq 2$  be real numbers. Suppose that a function  $f : X \rightarrow Y$  satisfies*

$$\|f(2x + y) + f(2x - y) - f(x + y) - f(x - y) - 6f(x)\| \leq \varepsilon(\|x\|^p + \|y\|^p)$$

for all  $x, y \in X$ . Then there exists a unique quadratic function  $T : X \rightarrow Y$  which satisfies the equation (1.3) and the inequality

$$\|f(x) - T(x)\| \leq \frac{5\varepsilon}{2|9 - 3^p|} \|x\|^p$$



An Inequality which Arises in the Absence of the Mountain Pass Geometry

Ick-Soon Chang and  
Hark-Mahn Kim

Title Page

Contents



Go Back

Close

Quit

Page 17 of 28

for all  $x \in X$ . The function  $T$  is given by


$$T(x) = \lim_{n \rightarrow \infty} \frac{f(3^n x)}{9^n} \quad \text{if } 0 < p < 2 \quad \left( T(x) = \lim_{n \rightarrow \infty} 9^n f\left(\frac{x}{3^n}\right) \quad \text{if } p > 2 \right)$$

for all  $x \in X$ . Further, if for each fixed  $x \in X$  the mapping  $t \mapsto f(tx)$  from  $\mathbb{R}$  to  $Y$  is continuous, then  $T(rx) = r^2 T(x)$  for all  $r \in \mathbb{R}$ .

**Remark 3.2.** If we put  $y = x = 0$  in the inequality of (3.10), we get  $6\|f(0)\| \leq \varepsilon$ . Applying Remark 3.1 to (3.10), we know that there exists a unique quadratic function  $T : X \rightarrow Y$  defined by  $T(x) = \lim_{n \rightarrow \infty} \frac{f(3^n x)}{9^n}$  which satisfies the equation (1.3) and the inequality

$$\|f(x) - T(x)\| \leq \frac{3\varepsilon}{16} + \frac{\|f(0)\|}{8} \leq \frac{5\varepsilon}{24}$$

for all  $x \in X$ . But we have a better possible upper bound (3.11) than that of the last inequality.




---

An Inequality which Arises in the Absence of the Mountain Pass Geometry

Ick-Soon Chang and  
Hark-Mahn Kim

---

Title Page

Contents



Go Back

Close

Quit

Page 18 of 28

## 4. Stability of (1.4)

We will investigate the Hyers-Ulam-Rassias stability problem for the functional equation (1.4). Thus we find the condition that there exists a true quadratic function near an approximately quadratic function.

**Theorem 4.1.** *Let  $\phi : X^2 \rightarrow \mathbb{R}^+$  be a function such that*

$$(4.1) \quad \sum_{i=0}^{\infty} \frac{1}{9^i} \left[ \frac{\phi(3^i x, 3^i x)}{2} + 2\phi(3^i x, 0) \right]$$

$$\left( \sum_{i=1}^{\infty} 9^i \left[ \frac{1}{2}\phi\left(\frac{x}{3^i}, \frac{x}{3^i}\right) + 2\phi\left(\frac{x}{3^i}, 0\right) \right], \text{ respectively} \right)$$

converges and

$$(4.2) \quad \lim_{n \rightarrow \infty} \frac{\phi(3^n x, 3^n y)}{9^n} = 0 \quad \left( \lim_{n \rightarrow \infty} 9^n \phi\left(\frac{x}{3^n}, \frac{y}{3^n}\right) = 0 \right)$$

for all  $x, y \in X$ . Suppose that a function  $f : X \rightarrow Y$  satisfies

$$(4.3) \quad \|f(2x + y) + f(x + 2y) - 4f(x + y) - f(x) - f(y)\| \leq \phi(x, y)$$

for all  $x, y \in X$ . Then there exists a unique quadratic function  $T : X \rightarrow Y$  which satisfies the equation (1.4) and the inequality

$$(4.4) \quad \|f(x) - T(x)\| \leq \frac{1}{9} \sum_{i=0}^{\infty} \frac{1}{9^i} \left[ \frac{\phi(3^i x, 3^i x)}{2} + 2\phi(3^i x, 0) \right] + \frac{\|f(0)\|}{4}$$

$$\left( \|f(x) - T(x)\| \leq \frac{1}{9} \sum_{i=1}^{\infty} 9^i \left[ \frac{1}{2}\phi\left(\frac{x}{3^i}, \frac{x}{3^i}\right) + 2\phi\left(\frac{x}{3^i}, 0\right) \right], \right)$$



An Inequality which Arises in  
the Absence of the Mountain  
Pass Geometry

Ick-Soon Chang and  
Hark-Mahn Kim

Title Page

Contents



Go Back

Close

Quit

Page 19 of 28

for all  $x \in X$ . The function  $T$  is given by

$$(4.5) \quad T(x) = \lim_{n \rightarrow \infty} \frac{f(3^n x)}{9^n} \quad \left( T(x) = \lim_{n \rightarrow \infty} 9^n f\left(\frac{x}{3^n}\right) \right)$$

for all  $x \in X$ .

*Proof.* If we write  $y = x$  in the inequality of (4.3), we get

$$(4.6) \quad \|f(3x) - 2f(2x) - f(x)\| \leq \frac{1}{2}\phi(x, x).$$

Putting  $y = 0$  in (4.3) and multiplying by 2, we have


$$(4.7) \quad \|2f(2x) - 8f(x)\| \leq 2\phi(x, 0) + 2\|f(0)\|$$

for all  $x \in X$ . Adding the inequality (4.6) with (4.7) and then dividing by 9, we get

$$(4.8) \quad \left\| \frac{f(3x)}{9} - f(x) \right\| \leq \frac{1}{9} \left[ \frac{\phi(x, x)}{2} + 2\phi(x, 0) + 2\|f(0)\| \right]$$

for all  $x \in X$ . Using the induction on  $n$ , we obtain that

$$(4.9) \quad \begin{aligned} \left\| \frac{f(3^n x)}{9^n} - f(x) \right\| &\leq \frac{1}{9} \sum_{i=0}^{n-1} \frac{1}{9^i} \left[ \frac{\phi(3^i x, 3^i x)}{2} + 2\phi(3^i x, 0) + 2\|f(0)\| \right] \\ &\leq \frac{1}{9} \sum_{i=0}^{\infty} \frac{1}{9^i} \left[ \frac{\phi(3^i x, 3^i x)}{2} + 2\phi(3^i x, 0) \right] + \frac{\|f(0)\|}{4} \end{aligned}$$



**An Inequality which Arises in  
the Absence of the Mountain  
Pass Geometry**

Ick-Soon Chang and  
Hark-Mahn Kim

Title Page

Contents



Go Back

Close

Quit

Page 20 of 28

for all  $x \in X$ .

Repeating the similar argument of Theorem 3.1, we obtain the desired result. The proof of assertion indicated by parentheses in the theorem is similarly proved and we omit it. In this case,  $f(0) = 0$  since  $\phi(0, 0) = 0$  by assumption. This completes the proof of the theorem.  $\square$

The proof of the following corollary is similar to that of Corollary 3.2.

**Corollary 4.2.** *Let  $\phi : {}_B\mathbb{B}_1 \times {}_B\mathbb{B}_1 \rightarrow \mathbb{R}^+$  be a function satisfying (4.1) and (4.2) for all  $x, y \in {}_B\mathbb{B}_1$ . Suppose that a mapping  $f : {}_B\mathbb{B}_1 \rightarrow {}_B\mathbb{B}_2$  satisfies*

$$(4.10) \quad \|f(2\alpha x + \alpha y) + f(\alpha x + 2\alpha y) - 4\alpha^2 f(x + y) - \alpha^2 f(x) - \alpha^2 f(y)\| \leq \phi(x, y)$$


for all  $\alpha \in B$  ( $|\alpha| = 1$ ) and for all  $x, y \in {}_B\mathbb{B}_1$ , and  $f$  is measurable or  $f(tx)$  is continuous in  $t \in \mathbb{R}$  for each fixed  $x \in {}_B\mathbb{B}_1$ . Then there exists a unique  $B$ -quadratic mapping  $T : {}_B\mathbb{B}_1 \rightarrow {}_B\mathbb{B}_2$ , defined by (4.5), which satisfies the equation (1.4) and the inequality (4.4) for all  $x \in {}_B\mathbb{B}_1$ .

**Corollary 4.3.** *Let  $E_1$  and  $E_2$  be Banach spaces over the complex field  $\mathbb{C}$ , and let  $\varepsilon \geq 0$  be a real number. Suppose that a mapping  $f : E_1 \rightarrow E_2$  satisfies*

$$\|f(2\alpha x + \alpha y) + f(\alpha x + 2\alpha y) - 4\alpha^2 f(x + y) - \alpha^2 f(x) - \alpha^2 f(y)\| \leq \varepsilon$$

for all  $\alpha \in \mathbb{C}$  ( $|\alpha| = 1$ ) and for all  $x, y \in E_1$ , and  $f$  is measurable or  $f(tx)$  is continuous in  $t \in \mathbb{R}$  for each fixed  $x \in E_1$ . Then there exists a unique  $\mathbb{C}$ -quadratic mapping  $T : E_1 \rightarrow E_2$  which satisfies the equation (1.3) and the inequality

$$\|f(x) - T(x)\| \leq \frac{5\varepsilon}{16}$$



An Inequality which Arises in the Absence of the Mountain Pass Geometry

Ick-Soon Chang and Hark-Mahn Kim

Title Page

Contents



Go Back

Close

Quit

Page 21 of 28

for all  $x \in E_1$ .

In Theorem 4.1, we obtain the alternative result if the conditions of  $\phi$  are replaced by the following.

**Remark 4.1.** Let  $\phi : X^2 \rightarrow \mathbb{R}^+$  be a function such that

$$\sum_{i=0}^{\infty} \frac{1}{4^i} \phi(2^i x, 0) \quad \left( \sum_{i=1}^{\infty} 4^i \phi\left(\frac{x}{2^i}, 0\right), \text{ respectively} \right)$$

converges and

$$\lim_{n \rightarrow \infty} \frac{\phi(2^n x, 2^n y)}{4^n} = 0 \quad \left( \lim_{n \rightarrow \infty} 4^n \phi\left(\frac{x}{2^n}, \frac{y}{2^n}\right) = 0 \right)$$


for all  $x, y \in X$ . Suppose that a function  $f : X \rightarrow Y$  satisfies

$$\|f(2x + y) + f(x + 2y) - 4f(x + y) - f(x) - f(y)\| \leq \phi(x, y)$$

for all  $x, y \in X$ . Then there exists a unique quadratic function  $T : X \rightarrow Y$  which satisfies the equation (1.4) and the inequality

$$(4.11) \quad \|f(x) - T(x)\| \leq \frac{1}{4} \sum_{i=0}^{\infty} \frac{1}{4^i} \phi(2^i x, 0) + \frac{\|f(0)\|}{3}$$

$$\left( \|f(x) - T(x)\| \leq \frac{1}{4} \sum_{i=1}^{\infty} 4^i \phi\left(\frac{x}{2^i}, 0\right) \right)$$



An Inequality which Arises in  
the Absence of the Mountain  
Pass Geometry

Ick-Soon Chang and  
Hark-Mahn Kim

Title Page

Contents



Go Back

Close

Quit

Page 22 of 28

for all  $x \in X$ . The function  $T$  is given by

$$T(x) = \lim_{n \rightarrow \infty} \frac{f(2^n x)}{4^n} \quad \left( T(x) = \lim_{n \rightarrow \infty} 4^n f\left(\frac{x}{2^n}\right) \right)$$

for all  $x \in X$ .

From Remark 4.1, we obtain the following corollary concerning the stability of the equation (1.4). We note that  $p$  need not be equal to  $q$  and  $\|f(0)\| = 0$  if  $p > 0$ .

**Corollary 4.4.** *Let  $X$  and  $Y$  be a real normed space and a Banach space, respectively, and let  $\varepsilon, p, q$  be real numbers such that  $\varepsilon \geq 0$ ,  $q > 0$  and either  $p, q < 2$  or  $p, q > 2$ . Suppose that a function  $f : X \rightarrow Y$  satisfies*

$$\|f(2x + y) + f(x + 2y) - 4f(x + y) - f(x) - f(y)\| \leq \varepsilon(\|x\|^p + \|y\|^q)$$


for all  $x, y \in X$ . Then there exists a unique quadratic function  $T : X \rightarrow Y$  which satisfies the equation (1.4) and the inequality

$$\|f(x) - T(x)\| \leq \frac{\varepsilon}{|4 - 2^p|} \|x\|^p + \frac{\|f(0)\|}{3}$$

for all  $x \in X$  and for all  $x \in X - \{0\}$  if  $p < 0$ . The function  $T$  is given by

$$T(x) = \lim_{n \rightarrow \infty} \frac{f(2^n x)}{4^n} \quad \text{if } p, q < 2 \quad \left( T(x) = \lim_{n \rightarrow \infty} 4^n f\left(\frac{x}{2^n}\right) \quad \text{if } p, q > 2 \right)$$

for all  $x \in X$ . Further, if for each fixed  $x \in X$  the mapping  $t \mapsto f(tx)$  from  $\mathbb{R}$  to  $Y$  is continuous, then  $T(rx) = r^2 T(x)$  for all  $r \in \mathbb{R}$ .



An Inequality which Arises in  
the Absence of the Mountain  
Pass Geometry

Ick-Soon Chang and  
Hark-Mahn Kim

Title Page

Contents



Go Back

Close

Quit

Page 23 of 28

As a consequence of the above Theorem 4.1, we have the following.

**Corollary 4.5.** *Let  $X$  and  $Y$  be a real normed space and a Banach space, respectively, and let  $\varepsilon \geq 0$ ,  $0 < p \neq 2$  be real numbers. Suppose that a function  $f : X \rightarrow Y$  satisfies*

$$\|f(2x + y) + f(x + 2y) - 4f(x + y) - f(x) - f(y)\| \leq \varepsilon(\|x\|^p + \|y\|^p)$$

for all  $x, y \in X$ . Then there exists a unique quadratic function  $T : X \rightarrow Y$  which satisfies the equation (1.4) and the inequality

$$\|f(x) - T(x)\| \leq \frac{3\varepsilon}{|9 - 3^p|} \|x\|^p$$

for all  $x \in X$ . The function  $T$  is given by


$$T(x) = \lim_{n \rightarrow \infty} \frac{f(3^n x)}{9^n} \quad \text{if } 0 < p < 2 \quad \left( T(x) = \lim_{n \rightarrow \infty} 9^n f\left(\frac{x}{3^n}\right) \quad \text{if } p > 2 \right)$$

for all  $x \in X$ . Further, if for each fixed  $x \in X$  the mapping  $t \mapsto f(tx)$  from  $\mathbb{R}$  to  $Y$  is continuous, then  $T(rx) = r^2 T(x)$  for all  $r \in \mathbb{R}$ .

The following corollary is an immediate consequence of Theorem 4.1.

**Corollary 4.6.** *Let  $X$  and  $Y$  be a real normed space and a Banach space, respectively, and let  $\varepsilon \geq 0$  be a real number. Suppose that a function  $f : X \rightarrow Y$  satisfies*

$$(4.12) \quad \|f(2x + y) + f(x + 2y) - 4f(x + y) - f(x) - f(y)\| \leq \varepsilon$$



An Inequality which Arises in the Absence of the Mountain Pass Geometry

Ick-Soon Chang and  
Hark-Mahn Kim

Title Page

Contents



Go Back

Close

Quit

Page 24 of 28



for all  $x, y \in X$ . Then there exists a unique quadratic function  $T : X \rightarrow Y$  defined by  $T(x) = \lim_{n \rightarrow \infty} \frac{f(2^n x)}{4^n}$  which satisfies the equation (1.4) and the inequality


$$(4.13) \quad \|f(x) - T(x)\| \leq \frac{5\varepsilon}{16}$$

for all  $x \in X$ . Further, if for each fixed  $x \in X$  the mapping  $t \mapsto f(tx)$  from  $\mathbb{R}$  to  $Y$  is continuous, then  $T(rx) = r^2T(x)$  for all  $r \in \mathbb{R}$ .

**Remark 4.2.** If we put  $y = x = 0$  in the inequality of (4.12), we get  $4\|f(0)\| \leq \varepsilon$ . Applying Remark 4.1 to (4.12), we know that there exists a unique quadratic function  $T : X \rightarrow Y$  defined by  $T(x) = \lim_{n \rightarrow \infty} \frac{f(2^n x)}{4^n}$  which satisfies the equation (1.4) and the inequality

$$\|f(x) - T(x)\| \leq \frac{\varepsilon}{3} + \frac{\|f(0)\|}{3} \leq \frac{5\varepsilon}{12}$$

for all  $x \in X$ . But we have a better possible upper bound (4.13) than that of the last inequality.




---

An Inequality which Arises in the Absence of the Mountain Pass Geometry

Ick-Soon Chang and  
Hark-Mahn Kim

---

Title Page

Contents



Go Back


Close

Quit

Page 25 of 28

## References

- [1] J. ACZÉL AND J. DHOMBRES, *Functional Equations in Several Variables*, Cambridge Univ. Press, 1989.
- [2] J. BAKER, The stability of the cosine equation, *Proc. Amer. Math. Soc.*, **80** (1980), 411–416.
- [3] P.W. CHOLEWA, Remarks on the stability of functional equations, *Aequationes Math.*, **27** (1984), 76–86.
- [4] S. CZERWIK, On the stability of the quadratic mapping in normed spaces, *Abh. Math. Sem. Univ. Hamburg*, **62** (1992), 59–64.
- [5] S. CZERWIK, The stability of the quadratic functional equation, In *Stability of Mappings of Hyers-Ulam Type* (edited by Th. M. Rassias and J. Tabor), Hadronic Press, Florida, 1994, 81–91.
- [6] G.L. FORTI, Hyers-Ulam stability of functional equations in several variables, *Aequationes Math.*, **50** (1995), 143–190.
- [7] P. GÁVRUTA, A generalization of the Hyers-Ulam-Rassias Stability of approximately additive mappings, *J. Math. Anal. Appl.*, **184** (1994), 431–436.
- [8] A. GRABIEC, The generalized Hyers-Ulam stability of a class of functional equations, *Publ. Math. Debrecen*, **48** (1996), 217–235.
- [9] D.H. HYERS, On the stability of the linear functional equation, *Proc. Natl. Acad. Sci.*, **27** (1941), 222–224.



---

An Inequality which Arises in  
the Absence of the Mountain  
Pass Geometry

Ick-Soon Chang and  
Hark-Mahn Kim

---

Title Page

Contents




Go Back

Close

Quit

Page 26 of 28

- [10] D.H. HYERS, G. ISAC AND Th. M. RASSIAS, *Stability of Functional Equations in Several Variables*, Birkhäuser, Basel, 1998.
- [11] D.H. HYERS, G. ISAC AND Th. M. RASSIAS, On the asymptoticity aspect of Hyers-Ulam stability of mappings, *Proc. Amer. Math. Soc.*, **126** (1998), 425–430.
- [12] D.H. HYERS AND Th. M. RASSIAS, Approximate homomorphisms, *Aequationes Math.*, **44** (1992), 125–153.
- [13] K.W. JUN AND Y.H. LEE, On the Hyers-Ulam-Rassias stability of a pexiderized quadratic inequality, *Math. Ineq. Appl.*, **4**(1) (2001), 93–118.
- [14] S.M. JUNG, On the Hyers-Ulam stability of the functional equations that have the quadratic property, *J. Math. Anal. Appl.*, **222** (1998), 126–137.
- [15] S.M. JUNG, On the Hyers-Ulam-Rassias stability of a quadratic functional equation, *J. Math. Anal. Appl.*, **232** (1999), 384–393.
- [16] Pl. KANNAPPAN, Quadratic functional equation and inner product spaces, *Results Math.*, **27** (1995), 368–372.
- [17] Th. M. RASSIAS, On the stability of the linear mapping in Banach spaces, *Proc. Amer. Math. Soc.*, **72** (1978), 297–300.
- [18] Th. M. RASSIAS, On the stability of functional equations in Banach spaces, *J. Math. Anal. Appl.*, **251** (2000), 264–284.
- [19] F. SKOF, Proprietà locali e approssimazione di operatori, *Rend. Sem. Mat. Fis. Milano*, **53** (1983), 113–129.




---

An Inequality which Arises in  
the Absence of the Mountain  
Pass Geometry

Ick-Soon Chang and  
Hark-Mahn Kim

---

Title Page

Contents




Go Back

Close

Quit

Page 27 of 28

[20] S. M. ULAM, *Problems in Modern Mathematics*, Chap. VI, Science Ed., Wiley, New York, 1960.



---

**An Inequality which Arises in  
the Absence of the Mountain  
Pass Geometry**

Ick-Soon Chang and  
Hark-Mahn Kim

---

Title Page

Contents



Go Back

Close

Quit

Page 28 of 28