



# A New Domino Tiling Sequence

Roberto Tauraso

Dipartimento di Matematica

Università di Roma “Tor Vergata”

via della Ricerca Scientifica

00133 Roma, Italy

[tauraso@mat.uniroma2.it](mailto:tauraso@mat.uniroma2.it)

## Abstract

*In this short note, we prove that the sequence [A061646](#) from Neil Sloane’s Online Encyclopedia of Integer Sequences is connected with the number of domino tilings of a holey square.*

## 1 Introduction

J. Bao discusses in [1] the following conjecture due to Edward Early (see also [2]): the number of tilings of a *holey square*, that is a  $2n \times 2n$  square with a hole of size  $2m \times 2m$  removed from the center, is  $2^{n-m} \cdot k^2$  where  $k$  is an odd number which depends on  $n$  and  $m$ . The conjecture has been proven for  $m = 1, 2, \dots, 6$  and numerical experiments indicate that it holds for several other values of  $m$ . Here we are going to show that the conjecture is verified also for  $m = n - 2$  and, perhaps what is more surprising, that the corresponding odd numbers  $k_n$  give a sequence already contained in Sloane’s Encyclopedia [3], namely [A061646](#).

Before proving this result we supply some definitions and notations. The Fibonacci numbers [A000045](#) are defined as the sequence of integers

$$f_0 = 0, f_1 = 1, \text{ and } f_n = f_{n-1} + f_{n-2} \text{ for all } n \geq 2.$$

We will need Cassini’s formula, one of the most famous Fibonacci identities:

$$f_{n+1}f_{n-1} - f_n^2 = (-1)^n \text{ for all } n \geq 1. \tag{1}$$

The terms of the sequence [A061646](#) satisfy the recurrence relation given by

$$a_0 = 1, a_1 = 1, a_2 = 3, \text{ and } a_n = 2a_{n-1} + 2a_{n-2} - a_{n-3} \text{ for all } n \geq 3,$$

and the corresponding generating function is

$$A(x) = \frac{1 - x - x^2}{(1 + x)(1 - 3x + x^2)}.$$

The number  $a_n$  can be expressed in terms of the Fibonacci numbers, as follows:


$$a_n = f_{n+1}f_{n-1} + f_n^2 = 2f_n^2 + (-1)^n \text{ for all } n \geq 1.$$

The second formula is obtained from the first by using (1) and it says that  $a_n$  is always an odd number. The first 20 values of the sequence  $a_n$  compared with the Fibonacci numbers are as follows:

$n$	$f_n$	$a_n$
0	0	1
1	1	1
2	1	3
3	2	7
4	3	19
5	5	49
6	8	129
7	13	337
8	21	883
9	34	2311
10	55	6051
11	89	15841
12	144	41473
13	233	108577
14	377	284259
15	610	744199
16	987	1948339
17	1597	5100817
18	2584	13354113
19	4181	34961521

## 2 The Result

**Lemma 2.1.** *For all  $n, m \geq 1$  the number of domino tilings of the L-grid*







[3] N. J. A. Sloane, The On-Line Encyclopedia of Integer Sequences. Published electronically at <http://www.research.att.com/~njas/sequences/>.

---

2000 *Mathematics Subject Classification*: Primary 05B45; Secondary 11B37, 11B39.

*Keywords*: domino tilings, Fibonacci numbers.

---

(Concerned with sequences [A000045](#) and [A061646](#).)

---

Received March 26 2004; revised version received April 19 2004. Published in *Journal of Integer Sequences*, April 27 2004.

---

Return to [Journal of Integer Sequences home page](#).