

WEIGHTED MULTIDIMENSIONAL INEQUALITIES FOR
MONOTONE FUNCTIONS

SORINA BARZA, LARS-ERIK PERSSON, Luleå

(Received January 18, 1999)

Dedicated to Professor Alois Kufner on the occasion of his 65th birthday

Abstract. We discuss the characterization of the inequality

$$\left(\int_{\mathbb{R}_+^N} f^q u \right)^{1/q} \leq C \left(\int_{\mathbb{R}_+^N} f^p v \right)^{1/p}, \quad 0 < q, p < \infty,$$

for monotone functions $f \geq 0$ and nonnegative weights u and v and $N \geq 1$. We prove a new multidimensional integral modular inequality for monotone functions. This inequality generalizes and unifies some recent results in one and several dimensions.

Keywords: integral inequalities, monotone functions, several variables, weighted L^p spaces, modular functions, convex functions, weakly convex functions

MSC 2000: 26D15, 26B99

1. INTRODUCTION

Let $\mathbb{R}_+^N := \{(x_1, \dots, x_N); x_i \geq 0, i = 1, 2, \dots, N\}$ and $\mathbb{R}_+ := \mathbb{R}_+^1$. Assume that $f: \mathbb{R}_+^N \rightarrow \mathbb{R}_+$ is monotone which means that it is monotone with respect to each variable. We denote $f \downarrow$, when f is decreasing (= nonincreasing) and $f \uparrow$ when f is increasing (= nondecreasing). Throughout this paper ω, u, v are positive measurable functions defined on \mathbb{R}_+^N , $N \geq 1$.

A function P on $[0, \infty)$ is called a modular function if it is strictly increasing, with the values 0 at 0 and ∞ at ∞ . For the definition of an N-function we refer to [7]. We say that a modular function P is weakly convex if $2P(t) \leq P(Mt)$, for all $t > 0$ and some constant $M > 1$. All convex modular functions are obviously weakly convex. The function $P_1(t) = t^p$, $0 < p < 1$ and the function $P_2(t) = \exp(\sqrt{t}) - 1$ are weakly convex, but not convex. See also [6].

In order to motivate this investigation and put it into a frame we use Section 2 to present the characterization of the inequality

$$(1) \quad \left(\int_{\mathbb{R}_+^N} f^q u \right)^{1/q} \leq C \left(\int_{\mathbb{R}_+^N} f^p v \right)^{1/p}, \quad 0 < p, q < \infty,$$

for all $f \downarrow$ or $f \uparrow$.

In Section 3 we will characterize the weights ω , u and v such that

$$(2) \quad Q^{-1} \left(\int_{\mathbb{R}_+^N} Q(\omega(x)f(x)) u(x) dx \right) \leq P^{-1} \left(\int_{\mathbb{R}_+^N} P(Cf(x)) v(x) dx \right)$$

holds for modular functions P and Q , where P is weakly convex and $0 \leq f \downarrow$. Here and in the sequel $C > 0$ denotes a constant independent of f .

C O N V E N T I O N S and notation. Products and quotients of the form $0 \cdot \infty$, $\frac{\infty}{\infty}$, $\frac{0}{0}$ are taken to be 0. \mathbb{Z} stands for the set of all integers and χ_E denotes the characteristic function of a set E .

2. WEIGHTED L^p INEQUALITIES FOR MONOTONE FUNCTIONS

In the one-dimensional case the inequality (1) was characterized in [8, Proposition 1] for both alternative cases $0 < p \leq q < \infty$ and $0 < q < p < \infty$ as follows:

(a) If $N = 1$, $0 < p \leq q < \infty$, then (1) is valid for all $f \downarrow$ if and only if

$$A_0 := \sup_{t>0} \left(\int_0^t u \right)^{1/q} \left(\int_0^t v \right)^{-1/p} < \infty$$

and the constant $C = A_0$ is sharp.

(b) If $N = 1$, $0 < q < p < \infty$, $1/r = 1/q - 1/p$, then (1) is true for all $f \downarrow$ if and only if

$$B_0 := \left(\int_0^\infty \left(\int_0^t u \right)^{r/p} \left(\int_0^t v \right)^{-r/p} u(t) dt \right)^{1/r} < \infty.$$

Moreover,

$$\left(\frac{q^2}{pr} \right)^{1/p} B_0 \leq C \leq \left(\frac{r}{q} \right)^{1/r} B_0$$

and

$$B_0^r = \frac{q}{r} \frac{\left(\int_0^\infty u \right)^{r/q}}{\left(\int_0^\infty v \right)^{r/p}} + \frac{q}{p} \int_0^\infty \left(\int_0^t u \right)^{r/q} \left(\int_0^t v \right)^{-r/q} v(t) dt.$$

(c) Similar characterizations are valid when $f \uparrow$, with the only change that the integrals over $[0, t]$ are replaced by integrals over $[t, \infty]$.

Since the one-dimensional inequality (1) expresses the embedding of classical Lorentz spaces, further generalizations and references in this directions can be found in [3].

The multidimensional case was recently treated in [1, Theorem 2.2], for the case $0 < p \leq q < \infty$ and in [2, Theorem 4.1], for the case $0 < q < p < \infty$ as follows:

(a) If $0 < p \leq q < \infty$, then (1) is valid for all $f \downarrow$ if and only if

$$A_N := \sup_{D \in \mathcal{D}_d} \frac{(\int_D u)^{1/q}}{(\int_D v)^{1/p}} < \infty$$

and the constant $C = A_N$ is sharp. Here the supremum is taken over the set \mathcal{D}_d of all “decreasing” domains, i.e., for which the characteristic function is a decreasing function in each variable.

(b) If $0 < q < p < \infty$, then (1) is valid for all $f \downarrow$ if and only if

$$B_N^r := \sup_{0 \leq h \downarrow} \int_0^\infty \left(\int_{D_{h,t}} v \right)^{-r/p} d \left(- \left(\int_{D_{h,t}} u \right)^{r/q} \right) < \infty,$$

where

$$D_{h,t} = \{x \in \mathbb{R}_+^N; h(x) > t\}.$$

Moreover,

$$\frac{1}{2^{1/q}(2^{r/q} + 2^{r/p})^{1/r}} B_N \leq C \leq 4^{1/q} B_N.$$

If $N = 1$, P and Q are N-functions and $Q \circ P^{-1}$ is convex, then some weight characterizations of the inequality (2) have been obtained in [4] and [5].

For $N > 1$, P and Q N-functions and $Q \circ P^{-1}$ convex, (2) holds for all $0 \leq f \downarrow$ if and only if there exists a constant $A = A(\Phi_1, \Phi_2, u, v, \omega)$ such that, for all $\varepsilon > 0$ and $D \in \mathcal{D}_d$,

$$Q^{-1} \left(\int_D Q(\varepsilon \omega(x)) u(x) dx \right) \leq P^{-1} \left(P(A\varepsilon) \int_D v(x) dx \right).$$

This characterization can be found in [2, Theorem 2.1].

However, if Q and P are not N-functions (hence not convex) and $Q \circ P^{-1}$ is not convex, then the problem of characterizing weights for which (2) holds seems to be to a large extent open. For $N = 1$ the first characterization of this type was given in [6].

In the next section we characterize the weights for which (2) holds when P is weakly convex. This result generalizes both the corresponding one-dimensional result obtained in [6] and the multidimensional case obtained in [2]. Some particular cases of (2) will also be pointed out.

3. A MULTIDIMENSIONAL MODULAR INEQUALITY

Let $0 \leq h(x) \downarrow$ and $t > 0$. Denote

$$D_{h,t} := \{x \in \mathbb{R}_+^N; h(x) > t\},$$

and

$$\mathcal{D}_d := \bigcup_{0 \leq h \downarrow} \bigcup_{t > 0} D_{h,t}.$$

The set \mathcal{D}_d consists of all “decreasing” domains $D_{h,t}$. In particular, $\chi_{D_{h,t}}$ is decreasing in each variable. For a strictly decreasing, positive sequence $\{t_k\}$, such that $t_k \rightarrow 0$ as $k \rightarrow \infty$ we put

$$D_k = D_{h,t_k} := \{x \in \mathbb{R}_+^N; h(x) > t_k\}, k \in \mathbb{Z}.$$

Obviously, $D_{k+1} \supset D_k$ and we define

$$\Delta_k = \Delta_{h,t_k} := D_{k+1} \setminus D_k.$$

Hence, $\Delta_k \cap \Delta_n = \emptyset$, $k \neq n$ and $\mathbb{R}_+^N = \bigcup_k \Delta_k$. For simplicity we also assume in the sequel that

$$(3) \quad \int_{\mathbb{R}_+^N} v(x) dx = \infty.$$

Theorem 3.1. *Let Q and P be modular functions and P weakly convex. Then (2) holds for all $0 \leq f \downarrow$ if and only if there exists a constant $B > 0$ such that*

$$(4) \quad Q^{-1} \left(\sum_{k \in \mathbb{Z}} \int_{\Delta_k} Q \left(\frac{\varepsilon_k}{B} \omega(x) \right) u(x) dx \right) \leq P^{-1} \left(\sum_{k \in \mathbb{Z}} P(\varepsilon_k) \int_{\Delta_k} v(x) dx \right)$$

is satisfied for all positive decreasing sequences $\{\varepsilon_k\}_{k \in \mathbb{Z}}$ and all increasing sequences of decreasing sets $\{D_k\}_{k \in \mathbb{Z}}$ such that $\int_{D_k} v(x) dx = 2^k$.

Proof. The necessity follows, if we replace f in (2) by the decreasing function $f = \sum_{k \in \mathbb{Z}} \varepsilon_k \chi_{\Delta_k}$, $\{\varepsilon_k\}_k$ being a decreasing sequence.

Next we consider the sufficiency. Fix $f \downarrow$ and set $\varepsilon_k = Bt_k$, $D_k = D_{f,t_k}$ and $\Delta_k = \Delta_{f,t_k}$. Because $\mathbb{R}_+^N = \bigcup_k \Delta_k$ we obtain, using also (4) and the facts that Q, P ,

Q^{-1} , P^{-1} are increasing and f is decreasing,

$$\begin{aligned}
Q^{-1}\left(\int_{\mathbb{R}_+^N} Q(\omega(x)f(x))u(x)dx\right) &= Q^{-1}\left(\sum_{k \in \mathbb{Z}} \int_{\Delta_k} Q(\omega(x)f(x))u(x)dx\right) \\
&\leq Q^{-1}\left(\sum_{k \in \mathbb{Z}} \int_{\Delta_k} Q(\omega(x)t_k)u(x)dx\right) \\
&\leq P^{-1}\left(\sum_{k \in \mathbb{Z}} P(Bt_k) \int_{\Delta_k} v(x)dx\right) \\
&= P^{-1}\left(\sum_{k \in \mathbb{Z}} 2P(Bt_k) \int_{\Delta_{k-1}} v(x)dx\right) \\
&\leq P^{-1}\left(\sum_{k \in \mathbb{Z}} \int_{\Delta_{k-1}} 2P(Bf(x))v(x)dx\right).
\end{aligned}$$

Therefore, by using the assumption that P is weakly convex, we find that

$$\begin{aligned}
Q^{-1}\left(\int_{\mathbb{R}_+^N} Q(\omega(x)f(x))u(x)dx\right) &\leq P^{-1}\left(\sum_{k \in \mathbb{Z}} \int_{\Delta_{k-1}} P(MBf(x))v(x)dx\right) \\
&= P^{-1}\left(\int_{\mathbb{R}_+^N} P(MBf(x))v(x)dx\right),
\end{aligned}$$

i.e., (2) holds with $C = MB$. The proof is complete. \square

We will give now two important corollaries of Theorem 3.1.

Corollary 3.2. *If P and Q are as in Theorem 3.1 and $Q \circ P^{-1}$ is convex, then (2) holds if and only if, for all $\varepsilon > 0$ and decreasing sets D , there exists a $C > 0$ such that*

$$(5) \quad Q^{-1}\left(\int_D Q\left(\frac{\omega(x)}{C}P^{-1}\left(\frac{\varepsilon}{\int_D v}\right)\right)u(x)dx\right) \leq P^{-1}(\varepsilon).$$

Proof. For the necessity we just have to substitute f in (2) with the function

$$f_0(x) = \frac{P^{-1}\left(\frac{\varepsilon}{\int_D v}\right)}{C} \chi_D(x).$$

Next we prove the sufficiency, i.e., that (5) implies (2). According to Theorem 3.1 it is sufficient to prove that (5) implies (4). By applying (5) with $\varepsilon = P(C\varepsilon_k) \int_{D_{k+1}} v$ for

each decreasing set D_{k+1} and using the convexity of $Q \circ P^{-1}$ and the weak convexity of P we find that

$$\begin{aligned} \left(\sum_{k \in \mathbb{Z}} \int_{\Delta_k} Q(\varepsilon_k \omega(x)) u(x) \, dx \right) &\leq \left(\sum_{k \in \mathbb{Z}} \int_{D_{k+1}} Q(\varepsilon_k \omega(x)) u(x) \, dx \right) \\ &\leq \sum_{k \in \mathbb{Z}} Q \circ P^{-1} \left(P(C\varepsilon_k) \int_{D_{k+1}} v \right) \\ &\leq Q \circ P^{-1} \left(\sum_{k \in \mathbb{Z}} 2P(C\varepsilon_k) \int_{D_k} v \right) \\ &\leq Q \circ P^{-1} \left(\sum_{k \in \mathbb{Z}} P(MC\varepsilon_k) 2^k \right) \\ &= Q \circ P^{-1} \left(\sum_{k \in \mathbb{Z}} P(MC\varepsilon_k) \int_{\Delta_k} v \right). \end{aligned}$$

Hence (4) follows with $B = MC$ and the corollary is proved. \square

Remark. If $Q(x) = x^q$ and $P(x) = x^p$, $0 < p \leq q < \infty$, then $Q \circ P^{-1}$ is convex and the condition (5) coincides with condition (3). Hence, Corollary 3.2 generalizes Theorem 2.2(d) in [1].

Remark. For $N = 1$ the condition (5) reads

$$Q^{-1} \left(\int_0^r Q \left(\frac{\omega(x)}{B} P^{-1} \left(\frac{\varepsilon}{\int_0^r v} \right) \right) u(x) \, dx \right) \leq P^{-1}(\varepsilon), \quad \forall r > 0.$$

Thus, if $N = 1$, then Corollary 3.2 coincides with Corollary 1 in [6].

Finally we apply Theorem 3.1 with $P(x) = x^p$ and $Q(x) = x^q$, $0 < p, q < \infty$, and obtain the following result:

Corollary 3.3. *The inequality (1) holds for all $0 < f \downarrow$ if and only if there exists a constant $K = K(p, q)$ such that*

$$\left(\sum_{k \in \mathbb{Z}} \varepsilon_k^q \int_{\Delta_k} u(x) \, dx \right)^{1/q} \leq K \left(\sum_{k \in \mathbb{Z}} \varepsilon_k^p \int_{\Delta_k} v(x) \, dx \right)^{1/p}$$

for all positive decreasing sequences $\{\varepsilon_k\}_{k \in \mathbb{Z}}$ and such that $\int_{D_k} v(x) \, dx = 2^k$.

Remark. For $N = 1$ a similar characterization is given in [6]. For other multidimensional characterizations of (1) in the case $0 < p \leq q < \infty$ see [1] and in the case $0 < q < p < \infty$ see [2] (cf. Section 2).

Final remarks. (i) The results in this paper can also be formulated when we remove the technical assumption (3) (cf. [2], [8]).

(ii) Similar results to all results in this paper can be formulated also for increasing functions of several variables.

References

- [1] Barza, S.; Persson, L. E.; Soria, J.: Sharp weighted multidimensional integral inequalities for monotone functions. *Math. Nachr.* To appear.
- [2] Barza, S.; Persson, L. E.; Stepanov, V. D.: On weighted multidimensional embeddings for monotone functions. *Math. Scand.* To appear.
- [3] Carro, M. J.; Pick, L.; Soria, J.; Stepanov, V. D.: On embeddings between classical Lorentz spaces. Research Report, 1998, submitted.
- [4] Drábek, P.; Heinig, H. P.; Kufner, A.: Weighted modular inequalities for monotone functions. *J. Inequal. Appl.* 1 (1997), 183–197.
- [5] Heinig, H. P.; Kufner, A.: Hardy operators on monotone functions and sequences in Orlicz spaces. *J. Lond. Math. Soc.*, 53(2) (1996), 256–270.
- [6] Heinig, H.; Lai, Q.: Weighted modular inequalities for Hardy-type operators defined on monotone functions. Research Report, 1998, submitted.
- [7] Krasnosel'ski, M. A.; Ruticki, Ya. B.: *Convex Functions and Orlicz spaces.* Noordhoff, Groningen, 1961.
- [8] Stepanov, V. D.: The weighted Hardy's inequality for nonincreasing functions. *Trans. Amer. Math. Soc.* 338 (1993), 173–186.

Authors' addresses: Sorina Barza, Lars-Erik Persson, Department of Mathematics, Luleå University of Technology, S-97187 Luleå, Sweden, e-mails: sorina@sm.luth.se, larserik@sm.luth.se.