

SOME DISCUSSIONS RELATED TO DISJOINT BER'S SUBPLANES

Milan Janić

Abstract. A projective plane \mathcal{P}_m is a Ber's subplane of a finite projective plane \mathcal{P}_n if every point and line of $\mathcal{P}_n \setminus \mathcal{P}_m$ is incident to some line and some point, respectively, of \mathcal{P}_n . It is known that the order of the plane \mathcal{P}_n and its Ber's subplane \mathcal{P}_m satisfy the equation $n = m^2$. In this article we prove some properties of finite projective planes \mathcal{P}_n having disjoint Ber's subplanes covering it.

Let \mathcal{P}_n , $n = m^2$, $n \in \mathbf{N}$ be a projective plane. Since the number of points of \mathcal{P}_n equals $n^2 + n + 1 = (m^2 + m + 1)(m^2 - m + 1)$, then, as long as the number of points is concerned, every such a projective plane \mathcal{P}_n could be covered by $m^2 - m + 1$ disjoint Ber's subplanes. We shall give an example (Theorem 1) of such a covering of the projective plane \mathcal{P}_4 , but it is still an open problem whether every projective plane of order $n = m^2$, $m \in \mathbf{N}$ has a covering by its disjoint Ber's subplanes.

THEOREM 1. *The plane \mathcal{P}_4 has a covering by its disjoint Ber's subplanes.*

Proof. Since $4 = 2^2$ and $2^2 - 2 + 1 = 3$, Ber's subplanes of \mathcal{P}_4 are of order 2 and if disjoint Ber's subplanes cover \mathcal{P}_4 , their number is 3. Let p_i^j and P_i^j , $i = 1, 2, \dots, 7$, $j = 1, 2, 3$ be the lines and the points of \mathcal{P}_4 with the incidences given by:

$$\begin{aligned} p_1^1 &= \{P_2^1, P_3^1, P_5^1, P_1^2, P_1^3\}, & p_1^2 &= \{P_2^2, P_3^2, P_5^2, P_1^1, P_1^3\}, & p_1^3 &= \{P_2^3, P_3^3, P_5^3, P_1^1, P_1^2\}, \\ p_2^1 &= \{P_1^1, P_4^1, P_5^1, P_6^2, P_6^3\}, & p_2^2 &= \{P_1^2, P_4^2, P_5^2, P_6^1, P_6^3\}, & p_2^3 &= \{P_1^3, P_4^3, P_5^3, P_6^1, P_6^2\}, \\ p_3^1 &= \{P_1^1, P_3^1, P_6^1, P_7^2, P_7^3\}, & p_3^2 &= \{P_7^1, P_1^2, P_3^2, P_6^2, P_7^3\}, & p_3^3 &= \{P_7^1, P_7^2, P_1^3, P_3^3, P_6^3\}, \\ p_4^1 &= \{P_2^1, P_4^1, P_6^1, P_3^2, P_3^3\}, & p_4^2 &= \{P_3^1, P_2^2, P_4^2, P_6^2, P_3^3\}, & p_4^3 &= \{P_3^1, P_3^2, P_2^3, P_4^3, P_6^3\}, \\ p_5^1 &= \{P_1^1, P_2^1, P_7^1, P_4^2, P_4^3\}, & p_5^2 &= \{P_4^1, P_1^2, P_2^2, P_7^2, P_4^3\}, & p_5^3 &= \{P_4^1, P_4^2, P_1^3, P_2^3, P_7^3\}, \\ p_6^1 &= \{P_3^1, P_4^1, P_7^1, P_5^2, P_5^3\}, & p_6^2 &= \{P_5^1, P_3^2, P_4^2, P_7^2, P_5^3\}, & p_6^3 &= \{P_5^1, P_5^2, P_3^3, P_4^3, P_7^3\}, \\ p_7^1 &= \{P_5^1, P_6^1, P_7^1, P_2^2, P_2^3\}, & p_7^2 &= \{P_2^1, P_5^2, P_6^2, P_7^2, P_2^3\}, & p_7^3 &= \{P_2^1, P_2^2, P_5^3, P_6^3, P_7^3\}. \end{aligned}$$

It is easy to check that $\mathcal{P}_2^1 = \{P_1^1, \dots, P_7^1\}$, $\mathcal{P}_2^2 = \{P_1^2, \dots, P_7^2\}$ and $\mathcal{P}_2^3 = \{P_1^3, \dots, P_7^3\}$ are disjoint Ber's subplanes which cover \mathcal{P}_4 . The incidences are probably more transparent in the figure 1. ■

Figure 1. A covering of \mathcal{P}_4 by disjoint Ber's subplanes

Now we are going to prove some properties of a covering of \mathcal{P}_n by disjoint Ber's subplanes. We say that a line p is incident to some subset \mathcal{P} of \mathcal{P}_n if the set \mathcal{P} is incident to at least two points of p .

THEOREM 2. *Let \mathcal{P}_m^i , $i = 1, \dots, s$ be disjoint Ber's subplanes of a projective plane \mathcal{P}_n . We denote by \mathcal{P}^s the set $\bigcup_{i=1}^s \mathcal{P}_m^i$ and by \mathcal{Q}^s the set $\mathcal{P}_n \setminus \mathcal{P}^s$. Each point of \mathcal{P}^s is incident to exactly $m+s$ lines of the Ber's subplanes \mathcal{P}_m^i , $i = 1, \dots, s$. Each point of \mathcal{Q}^s is incident to exactly s lines of the Ber's subplanes \mathcal{P}_m^i , $i = 1, \dots, s$.*

Proof. Every point of \mathcal{P}_n is incident to at least one line of each subplane \mathcal{P}_m^i , $i \leq s$ by the definition of a Ber's subplane. If $P \in \mathcal{P}_m^i$, then P is incident to $m+1$ lines of \mathcal{P}_m^i and to a single line of each \mathcal{P}_m^j , $j \neq i$. (If a point is incident to two lines of a subplane it is incident to that subplane.) Therefore, if P is a point of \mathcal{P}^s , it is incident to $m+1+s-1 = m+s$ lines of the subplanes \mathcal{P}_m^i , $i \leq s$. On the other hand, the points of \mathcal{Q}^s are incident only to a single line of each subplane \mathcal{P}_m^i , $i \leq s$, which proves the theorem. ■

We shall call the property given by the previous theorem *the homogeneity of finite projective planes*. Let us prove one more property of disjoint Ber's subplanes.

THEOREM 3. *If a plane \mathcal{P}_n , $n = m^2$, $m \in \mathbf{N}$ contains $m^2 - m$ disjoint Ber's subplanes \mathcal{P}_m^i , $i = 1, \dots, m^2 - m$, of order m , then $\mathcal{Q} = \mathcal{P}_n \setminus \bigcup_{i=1}^{m^2-m} \mathcal{P}_m^i$ is a Ber's subplane.*

Proof. Since every line p is incident to a single point or to $m+1$ points of each \mathcal{P}_m^i , $i \leq m^2 - m$, then it has $k(m+1) + (m^2 - m - k) + l = m^2 + 1$ points, where k is the number of the subplanes \mathcal{P}_m^i , $i \leq m^2 - m$ to which it is incident and l the number of the points of $p \cap \mathcal{Q}$. Hence, $k > 1$ is impossible, $k = 1$ implies $l = 1$, and $k = 0$ implies $l = m+1$, which proves that a line intersects \mathcal{Q} in a single point if it is not incident to \mathcal{Q} , and in $m+1$ points otherwise.

Every point of \mathcal{P}_n is incident to a single line of each Ber's subplane to which it does not belong and to $m+1$ lines of the subplane to which it belongs. This implies that every point P of \mathcal{Q} is incident to $m^2 - m$ lines of the subplanes \mathcal{P}_m^i , $i \leq m^2 - m$ and $m^2 + 1 - (m^2 - m) = m+1$ lines not incident to the subplanes \mathcal{P}_m^i , $i \leq m^2 - m$. Hence, every point P of \mathcal{Q} is incident to $m+1$ lines of \mathcal{Q} . By the Theorem 2, every point of $\mathcal{P}_n \setminus \mathcal{Q}$ is incident to $m + (m^2 - m)$ lines of the subplanes \mathcal{P}_m^i , $i \leq m^2 - m$, and therefore, to a single line of \mathcal{Q} .

Since \mathcal{Q} contains $m^2 + m + 1 = (n^2 + n + 1) - (m^2 + m + 1)(m^2 - m)$ points lying on lines with $m+1$ such points (points of \mathcal{Q}), and since each point of \mathcal{Q} is incident to $m+1$ such lines, as it is well known, \mathcal{Q} is a projective subplane of order m . Finally, \mathcal{Q} is a Ber's subplane according to the properties proved above. ■

REFERENCES

- [1] D. R. Hughes and F. C. Piper, *Projective planes*, Berlin 1972
- [2] Ф. Картеси, *Введение в конечные геометрии*, Наука, Москва 1980
- [3] M. Hall Jr., *The theory of groups*, Mac Millan, New York 1960
- [4] M. Hall Jr., *Projective planes*, Trans. Math. **80**, 1955

(received 22.02.1993, in revised form 23.10.1994.)