

C. Fontanari

## MODULI OF CURVES VIA ALGEBRAIC GEOMETRY

**Abstract.** Here we discuss some open problems about moduli spaces of curves from an algebro-geometric point of view. In particular, we focus on Arbarello stratification and we show that its top dimensional stratum is affine.

The moduli space  $\overline{\mathcal{M}}_{g,n}$  of stable  $n$ -pointed genus  $g$  curves is by now a widely explored subject (see for instance the book [10] and the references therein), but many interesting problems in the field are still unsolved, both from a topological and a geometrical point of view. Even though various methods have been fruitfully applied (e. g. Teichmüller spaces, Hodge theory, G.I.T., ...), a purely algebro-geometric approach seems to be quite powerful and rather promising as well. We wish to mention at least the recent paper [3] by Enrico Arbarello and Maurizio Cornalba: as the authors point out in the introduction, what is really new there is the method of proof, which is based on standard algebro-geometric techniques.

Indeed, the only essential result borrowed from geometric topology is a vanishing theorem due to John Harer. Namely, the fact that  $H_k(\mathcal{M}_{g,n})$  vanishes for  $k > 4g - 4 + n$  if  $n > 0$  and for  $k > 4g - 5$  if  $n = 0$  was deduced in [9] from the construction of a  $(4g - 4 + n)$ -dimensional spine for  $\mathcal{M}_{g,n}$  by means of Strebel differentials. On the other hand, it is conceivable that Harer's vanishing is only the tip of an iceberg of deeper geometrical properties. For instance, a conjecture of Eduard Looijenga says that  $\mathcal{M}_g$  is a union of  $g - 1$  open subsets (see [7], Conjecture 11.3), but (as far as we know) there are no advances in this direction. Another strategy (see [8], Problem 6.5) in order to avoid the use of Strebel's differentials in the proof of Harer's theorem is to look for an orbifold stratification of  $\mathcal{M}_g$  with  $g - 1$  affine subvarieties as strata.

A natural candidate for such a stratification is provided by a flag of subvarieties introduced by Enrico Arbarello in his Ph.D. thesis. Namely, for each integer  $n$ ,  $2 \leq n \leq g$ , he defined the subvariety  $W_{n,g} \subset \mathcal{M}_g$  as the sublocus of  $\mathcal{M}_g$  described by those points of  $\mathcal{M}_g$  which correspond to curves of genus  $g$  which can be realized as  $n$ -sheeted coverings of  $\mathbb{P}^1$  with a point of total ramification (see [2] p. 1). The natural expectation (see [1] p. 326 but also [12] p. 310) was that  $W_{n,g} \setminus W_{n-1,g}$  does not contain any complete curve. About ten years later, Steven Diaz was able to prove that a slightly different flag of subvarieties enjoys such a property and he deduced from this fact his celebrated bound on the dimension of complete subvarieties in  $\mathcal{M}_g$  (see [5]). It remains instead an open question whether or not the open strata of the Arbarello flag admit complete curves (see [10] p. 291).

Perhaps an even stronger conjecture could be true: since  $W_{2,g}$  is the hyperelliptic locus, which is well-known to be affine (see for instance [11] p. 320), one may wonder whether all the open strata  $W_{n,g} \setminus W_{n-1,g}$  are affine. We were not able to prove this statement in full generality; however, we found an elementary proof that the top dimensional stratum is indeed affine.

**THEOREM 1.** *If  $g \geq 3$  then  $\mathcal{M}_g \setminus W_{g-1,g}$  is affine.*

*Proof.* Since  $\mathcal{M}_g \setminus W_{g-1,g} = \overline{\mathcal{M}}_g \setminus (\text{supp}(\overline{W}_{g-1,g}) \cup \partial\overline{\mathcal{M}}_g)$ , it is sufficient to prove that  $\text{supp}(\overline{W}_{g-1,g}) \cup \partial\overline{\mathcal{M}}_g$  is the support of an effective ample divisor on  $\overline{\mathcal{M}}_g$ . The class of  $\overline{W}_{g-1,g}$  in the Picard group of  $\overline{\mathcal{M}}_g$  was computed by Steven Diaz in his Ph.D. thesis (see [6]), so we know that

$$[\overline{W}_{g-1,g}] = a\lambda - \sum_i b_i \delta_i$$

where

$$\begin{aligned} a &:= \frac{g^2(g-1)(3g-1)}{2} \\ b_0 &:= \frac{(g-1)^2g(g+1)}{6} \\ b_i &:= \frac{i(g-i)g(g^2+g-4)}{2} \quad (i > 0). \end{aligned}$$

In particular, notice that if  $g \geq 3$  then  $a > 11$  and  $b_i > 1$  for every  $i$ . Consider now the following divisor on  $\overline{\mathcal{M}}_g$ :

$$D := \overline{W}_{g-1,g} + \sum_i (b_i - 1)\Delta_i.$$

Since  $b_i > 1$  we see that  $D$  is effective; moreover, we have  $\text{supp}(D) = \text{supp}(\overline{W}_{g-1,g}) \cup \partial\overline{\mathcal{M}}_g$ . We claim that  $D$  is ample. Indeed,

$$\begin{aligned} [D] &= [\overline{W}_{g-1,g}] + \sum_i (b_i - 1)\delta_i \\ &= a\lambda - \sum_i b_i \delta_i + \sum_i b_i \delta_i - \sum_i \delta_i \\ &= a\lambda - \delta. \end{aligned}$$

Since  $a > 11$  we may deduce that  $D$  is ample from the Cornalba-Harris criterion (see [4], Theorem 1.3), so the proof is over.  $\square$

## References

- [1] ARBARELLO E., *Weierstrass points and moduli of curves*, *Compositio Math.* **29** (1974), 325–342.
- [2] ARBARELLO E., *On subvarieties of the moduli space of curves of genus  $g$  defined in terms of Weierstrass points*, *Atti Accad. Naz. Lincei Mem. Cl. Sci. Fis. Mat. Natur. Sez. I* **8** **15** (1978), no. 1, 3–20.
- [3] ARBARELLO E. AND CORNALBA M., *Calculating cohomology groups of moduli spaces of curves via algebraic geometry*, *Inst. Hautes Études Sci. Publ. Math.* **88** (1998), 97–127.
- [4] CORNALBA M. AND HARRIS J., *Divisor classes associated to families of stable varieties, with applications to the moduli space of curves*, *Ann. Sc. Ec. Norm. Sup.* **21** (4) (1988), 455–475.
- [5] DIAZ S., *A bound on the dimensions of complete subvarieties of  $\mathcal{M}_g$* , *Duke Math. J.* **151** (1984), 405–408.

- [6] DIAZ S., *Exceptional Weierstrass points and the divisor on moduli space that they define*, Memoirs of the Am. Math. Soc. **327** (1985).
- [7] FABER C. AND LOOIJENGA E., *Remarks on moduli of curves. Moduli of curves and abelian varieties*, Aspects Math. **33** (1999), 23–45.
- [8] HAIN R. AND LOOIJENGA E., *Mapping class groups and moduli spaces of curves*, Proc. Symp. Pure Math. AMS **62** (1998), 97–142.
- [9] HARER J., *The virtual cohomological dimension of the mapping class group of an orientable surface*, Inv. Math. **84** (1986), 157–176.
- [10] HARRIS J. AND MORRISON I., *Moduli of curves*, Graduate Texts in Math. **187**, Springer 1998.
- [11] MORRISON I., *Subvarieties of moduli spaces of curves: open problems from an algebro-geometric point of view*, Contemp. Math. AMS **150** (1993), 317–343.
- [12] MUMFORD D., *Towards an enumerative geometry of the moduli space of curves*, Progress in Math. **36**, Birkhäuser, Boston 1983, 271–328.

**AMS Subject Classification:** 14H10, 14H55.

Claudio FONTANARI  
Dipartimento di Matematica  
Università degli studi di Trento  
Via Sommarive 14  
38050 Povo (Trento), ITALIA  
e-mail: fontanar@science.unitn.it

