


*Geometry & Topology*  
 Volume 4 (2000) 149{170  
 Published: 11 March 2000



## Double point self-intersection surfaces of immersions

Mohammad A Asadi-Golmankhaneh  
 Peter J Eccles

*Department of Mathematics, University of Urmia*  
*PO Box 165, Urmia, Iran*

and

*Department of Mathematics, University of Manchester*  
*Manchester, M13 9PL, UK*

Email: pj eccles@man. ac. uk

### Abstract

A self-transverse immersion of a smooth manifold  $M^{k+2}$  in  $\mathbb{R}^{2k+2}$  has a double point self-intersection set which is the image of an immersion of a smooth surface, the double point self-intersection surface. We prove that this surface may have odd Euler characteristic if and only if  $k \equiv 1 \pmod{4}$  or  $k+1$  is a power of 2. This corrects a previously published result by Andras Szűcs [22].

The method of proof is to evaluate the Stiefel-Whitney numbers of the double point self-intersection surface. By the methods of [2] these numbers can be read off from the Hurewicz image  $h(\sigma) \in H_{2k+2}(\mathbb{Z}/2) \cong MO(k)$  of the element  $\sigma \in H_{2k+2}(\mathbb{Z}/2) \cong MO(k)$  corresponding to the immersion under the Pontrjagin-Thom construction.

**AMS Classification numbers** Primary: 57R42

Secondary: 55R40, 55Q25, 57R75

**Keywords:** immersion, Hurewicz homomorphism, spherical class, Hopf invariant, Stiefel-Whitney number

Proposed: Ralph Cohen  
 Seconded: Gunnar Carlsson, Yasha Eliashberg

Received: 30 July 1999  
 Accepted: 29 February 2000

*Copyright Geometry and Topology*

## 1 Introduction

Let  $f: M^{n-k} \hookrightarrow \mathbb{R}^n$  be a self-transverse immersion of a compact closed smooth  $(n-k)$ -dimensional manifold in  $n$ -dimensional Euclidean space ( $0 < k < n$ ). A point of  $\mathbb{R}^n$  is an  $r$ -fold self-intersection point of the immersion if it is the image under  $f$  of (at least)  $r$  distinct points of the manifold. The self-transversality of  $f$  implies that the set of  $r$ -fold self-intersection points (the  $r$ -fold self-intersection set) is itself the image of an immersion

$${}_r(f): {}_r(f) \hookrightarrow \mathbb{R}^n$$

of a compact manifold  ${}_r(f)$ , the  $r$ -fold self-intersection manifold, of dimension  $n-rk$ , although in general this immersion will not be self-transverse. It is natural to ask what can be said about this  $r$ -fold intersection manifold: which manifolds can arise for each value of  $n$ ,  $k$  and  $r$ ?

The simplest case is when  $n = rk$  so that the self-intersection manifold is a finite set of points and this case was the first to be considered in detail (see [3], [10], [11], [12], [16]).

More recently cases of higher dimensional self-intersection manifolds have been considered. Andras Szűcs was one of the first to do so and in [22] considered the simplest case of interest, when the double point self-intersection manifold is a surface; this arises for  $n = 2k + 2$ . In this paper we return to this case using different methods to Szűcs. Our main result is the following.

**Theorem 1.1** *For  $k \geq 1$ , there exists an immersion  $f: M^{k+2} \hookrightarrow \mathbb{R}^{2k+2}$  with double point self-intersection manifold of odd Euler characteristic if and only if  $k \equiv 1 \pmod{4}$  or  $k+1$  is a power of 2.*

This result should be contrasted with Szűcs' result which asserted that double point self-intersection manifolds of odd Euler characteristic can occur only if  $k \equiv 1 \pmod{4}$ . Szűcs's approach used differential topology and the argument in the case  $k \equiv 3 \pmod{4}$  was particularly delicate.

Our approach uses algebraic topology and in particular the correspondence between bordism groups and homotopy groups of Thom complexes. In [2] we described a general approach to these problems which gives a method for determining the bordism class of the self-intersection manifolds of any immersion: the unoriented bordism class of a manifold is determined by its Stiefel-Whitney numbers and the Stiefel-Whitney numbers of the self-intersection manifolds of an immersion can be read off from certain homological information about the

immersion. For double point self-intersection surfaces the situation is particularly simple since there are only two bordism classes: a compact surface is a boundary if its Euler characteristic is even and is a non-boundary bordant to the projective plane if its Euler characteristic is odd.

The paper is organized as follows. In Section 2 we recall the results needed from [2], establish our basic notation and outline the proof of the theorem covering the steps which apply in all cases. In Section 3 we complete the proof in the easiest case of  $k$  even and this is followed by the proofs for  $k \equiv 1 \pmod 4$  in Section 4 and  $k \equiv 3 \pmod 4$  in Section 5. Finally, in Section 6, we comment on the relationship between our results and those in Szűcs' paper [22]. Almost everything in that paper is confirmed by our methods.

**Acknowledgements** Most of the results in this paper are contained in the first author's thesis [1] which also contains other applications of these methods. He was supported by the University of Urmia and the Ministry of Culture and Higher Education of the Islamic Republic of Iran during his time as a student at the University of Manchester.

## 2 The Stiefel Whitney characteristic numbers of the double point self-intersection manifold

Let  $\text{Imm}(n - k; k)$  denote the group of bordism classes of immersions  $M^{n-k} \looparrowright \mathbb{R}^n$  of compact closed smooth manifolds in Euclidean  $n$ -space. Details of (co)bordism in this setting have been given by R Wells in [24]. By general position every immersion is regularly homotopic (and so bordant) to a self-transverse immersion and so each element of  $\text{Imm}(n - k; k)$  can be represented by a self-transverse immersion. In the same way bordisms between self-transverse immersions can be taken to be self-transverse; it is clear that such a bordism will induce a bordism of the immersions of the double point self-intersection manifolds so that  $f \looparrowright \mathbb{R}^n \rightarrow \mathbb{R}^{2n}$  induces a double point self-intersection map

$$\mathbb{Z}_2: \text{Imm}(n - k; k) \rightarrow \text{Imm}(n - 2k; 2k):$$

Wells shows how  $\text{Imm}(n - k; k)$  may be described as a stable homotopy group. Let  $MO(k)$  denote the Thom complex of a universal  $O(k)$ -bundle  $\pi: EO(k) \rightarrow BO(k)$  (see [18] for basic material on vector bundles, Thom complexes and bordism theory). Using the Pontrjagin-Thom construction, Wells describes an isomorphism

$$\mathbb{Z}_2: \text{Imm}(n - k; k) \cong \mathbb{Z}_2 \otimes_n MO(k):$$

Write  $QX$  for the direct limit  $QX = \lim_{\leftarrow} \Sigma^{-n} X$ , where  $\Sigma^{-n}$  denotes the reduced suspension functor and  $\Omega$  denotes the loop space functor. By the adjointness of the functors  $\Sigma^{-n}$  and  $\Omega$ ,  $\Sigma^{-n} MO(k) = \Omega^n QMO(k)$ . We consider the  $\mathbb{Z}/2$ -homology Hurewicz homomorphism

$$h: \Sigma^{-n} MO(k) \rightarrow \Omega^n QMO(k) \quad H_n QMO(k) = H_n(QMO(k); \mathbb{Z}/2):$$

(Throughout this paper we use  $H$  and  $H$  to denote homology and cohomology with  $\mathbb{Z}/2$  coefficients.)

The main result of [2] describes how, for a self-transverse immersion  $f: M^{n-k} \hookrightarrow \mathbb{R}^n$  corresponding to  $\Sigma^{-n} MO(k)$ , the Hurewicz image  $h(f) \in H_n QMO(k)$  determines the normal Stiefel-Whitney numbers of the self-intersection manifold  $\sigma_r(f)$ .

To state this result in the case of double point self-intersection manifolds we need some preliminaries.

The *quadratic construction* on a pointed space  $X$  is defined to be

$$D_2 X = X \wedge X \rtimes_{\mathbb{Z}/2} S^1 = X \wedge X \wedge_{\mathbb{Z}/2} S^1 = f \circ g \wedge_{\mathbb{Z}/2} S^1;$$

where the non-trivial element of the group  $\mathbb{Z}/2$  acts on  $X \wedge X$  by permuting the co-ordinates and on the infinite sphere  $S^1$  by the antipodal action. There is a natural map

$$h^2: QX \rightarrow QD_2 X$$

known as a *stable James-Hopf map* which induces a *stable Hopf invariant*

$$h^2: \Sigma^{-n} X \rightarrow \Sigma^{-n} D_2 X$$

(see [4] and [19]).

If the self-transverse immersion  $f: M^{n-k} \hookrightarrow \mathbb{R}^n$  corresponds to the element  $\Sigma^{-n} MO(k)$ , then the immersion of the double point self-intersection manifold  $\sigma_2(f): \sigma_2(f) \hookrightarrow \mathbb{R}^n$  corresponds to the element  $h^2(f) \in \Sigma^{-n} D_2 MO(k)$  given by the stable Hopf invariant (see [15], [20], [21], [23]).

The immersion  $\sigma_2(f)$  corresponds to an element in the stable homotopy of  $D_2 MO(k)$  because the immersion of the double point self-intersection manifold automatically acquires additional structure on its normal bundle, namely that at each point  $f(x_1) = f(x_2)$  the normal  $2k$ -dimensional space is decomposed as the direct sum of the two (unordered)  $k$ -dimensional normal spaces of  $f$  at the points  $x_1$  and  $x_2$ . The universal bundle for this structure is

$$k \oplus k \wedge_{\mathbb{Z}/2} 1: EO(k) \wedge_{\mathbb{Z}/2} EO(k) \wedge_{\mathbb{Z}/2} S^1 \rightarrow BO(k) \wedge_{\mathbb{Z}/2} BO(k) \wedge_{\mathbb{Z}/2} S^1$$

which has the Thom complex  $D_2 MO(k)$ .

Forgetting this additional structure on the immersion corresponds to applying the map

$$: \mathbb{S}_n D_2 MO(k) \rightarrow \mathbb{S}_n MO(2k)$$

induced by the map of Thom complexes  $\mathbb{S}_n D_2 MO(k) \rightarrow \mathbb{S}_n MO(2k)$  which comes from the map  $BO(k) \rightarrow BO(k) \times_{\mathbb{Z}_2} S^1 \rightarrow BO(2k)$  classifying the bundle  $\xi_k \oplus \xi_k \oplus \mathbb{Z}_2$ . Thus, we have the following commutative diagram.

$$\begin{array}{ccc}
 I(n-k; k) & \xrightarrow{\quad \tau \quad} & I(n-2k; 2k) \\
 \downarrow \cong & & \downarrow \cong \\
 \mathbb{S}_n MO(k) & \xrightarrow{\quad h^2 \quad} & \mathbb{S}_n D_2 MO(k) \xrightarrow{\quad} \mathbb{S}_n MO(2k)
 \end{array} \tag{1}$$

Turning now to homology, observe that, by adjointness, the stable James-Hopf map  $h^2: QX \rightarrow QD_2X$  gives a stable map  $\mathbb{S}^1 QX \rightarrow \mathbb{S}^1 D_2X$  inducing a map in homology  $h^2: H_n QX \rightarrow H_n D_2X$ . This gives the following commutative diagram.

$$\begin{array}{ccccc}
 \mathbb{S}_n MO(k) & \xrightarrow{\quad h^2 \quad} & \mathbb{S}_n D_2 MO(k) & \xrightarrow{\quad} & \mathbb{S}_n MO(2k) \\
 \downarrow \cong & & \downarrow h^S & & \downarrow h^S \\
 \mathbb{S}_n QMO(k) & & \mathbb{S}_n QD_2 MO(k) & & \mathbb{S}_n QMO(2k) \\
 \downarrow h & & \downarrow \cong & & \downarrow \cong \\
 H_n QMO(k) & \xrightarrow{\quad h^2 \quad} & H_n QD_2 MO(k) & \xrightarrow{\quad} & H_n QMO(2k)
 \end{array} \tag{2}$$

In this diagram the second and third vertical maps are stable Hurewicz homomorphisms defined using the fact the Hurewicz homomorphisms commute with suspension. The first square commutes by the definition of the stable Hurewicz map and by naturality, and the second square commutes by naturality.

Notice that the normal Stiefel-Whitney numbers (and so the bordism class) of the double point self-intersection manifold  $\Sigma_2(f)$  of an immersion  $f: M^{n-k} \hookrightarrow \mathbb{R}^n$  corresponding to  $\tau \in \mathbb{S}_n MO(k)$  are determined by (and determine) the Hurewicz image  $h^S(\tau)$  of the element  $\tau = h^2(\tau) \in \mathbb{S}_n MO(2k)$  corresponding to the immersion  $\Sigma_2(f)$ . To be more explicit in the case under consideration we recall the structure of  $H^* MO(k)$ .

Let  $e_i \in H_i BO(1) = \mathbb{Z} = 2$  be the non-zero element (for  $i \equiv 0$ ). For each sequence  $I = (i_1; i_2; \dots; i_k)$  of non-negative integers we define

$$e_I = e_{i_1} e_{i_2} \dots e_{i_k} = \binom{k}{I} (e_{i_1} \ e_{i_2} \ \dots \ e_{i_k}) \in H BO(k)$$

where  $\binom{k}{I}: BO(1)^k \rightarrow BO(k)$  is the map which classifies the product of the universal line bundles. The dimension of  $e_I$  is  $|I| = i_1 + i_2 + \dots + i_k$ .

From the definition of  $\binom{k}{I}$ ,  $e_{i_1} e_{i_2} \dots e_{i_k} = e_{i_{\sigma(1)}} e_{i_{\sigma(2)}} \dots e_{i_{\sigma(k)}}$  for each  $\sigma \in S_k$ , where  $S_k$  is the permutation group on  $k$  elements. Thus each such element can be written as  $e_{i_1} e_{i_2} \dots e_{i_k}$  where  $i_1 \leq i_2 \leq \dots \leq i_k$  and it follows by a counting argument that

$$f e_{i_1} e_{i_2} \dots e_{i_k} j_0 \ i_1 \ i_2 \ \dots \ i_k g$$

is a basis for  $H BO(k)$  (see [14] Proposition 2.4.3).

The sphere bundle of the universal  $O(k)$  bundle is given up to homotopy by the inclusion  $BO(k-1) \rightarrow BO(k)$  and so the Thom complex  $MO(k)$  is homotopy equivalent to the quotient space  $BO(k) = BO(k-1)$ . It follows that

$$f e_{i_1} e_{i_2} \dots e_{i_k} j_1 \ i_1 \ i_2 \ \dots \ i_k g$$

is a basis for  $H MO(k)$ .

By Diagrams (1) and (2), the double point self-intersection surface of an immersion  $M^{k+2} \hookrightarrow \mathbb{R}^{2k+2}$  may be identified up to bordism by using the stable Hurewicz homomorphism

$$h^S: \pi_{2k+2}^S MO(2k) \rightarrow H_{2k+2} MO(2k):$$

From the above,  $H_{2k+2} MO(2k)$  has a basis  $f e_1^{2k-1} e_3; e_1^{2k-2} e_2^2 g$ . The element  $e_1^{2k-1} e_3 \in \pi_{2k+2}^S MO(2k)$  corresponds to an immersion of a non-boundary (ie, a surface of odd Euler characteristic, bordant to the real projective plane) if and only if  $h^S(e_1^{2k-1} e_3) = e_1^{2k-1} e_3$ , the only non-zero stably spherical element. This corresponds to the fact that a surface  $L$  is a non-boundary if and only if the normal Stiefel-Whitney number  $\overline{w}_1^2[L] = 1$  (see [2] Proposition 3.4).

For  $k > 1$  we are in the stable range,  $\pi_{2k+2}^S MO(2k) = \mathbb{Z} = 2$ , and  $h^S$  is a monomorphism: any two immersions of bordant manifolds are bordant. On the other hand, for  $k = 1$  the group  $\text{Imm}(2; 4) = \pi_4^S MO(2)$  is infinite and  $h^S$  is not a monomorphism. The bordism class of an immersion  $L^2 \hookrightarrow \mathbb{R}^4$  is not determined by  $L$ .

This discussion can be summed up in the following theorem which follows essentially from Diagrams (1) and (2).

**Theorem 2.1** Suppose that  $f: M^{k+2} \looparrowright \mathbb{R}^{2k+2}$  is a self-transverse immersion corresponding to  $\in \mathbb{Z} \oplus_{2k+2} MO(k)$ . Then the double point self-intersection surface  $\Sigma_2(f)$  has odd Euler characteristic and so is a non-boundary if and only if

$$h^2 h(\Sigma_2(f)) = e_1^{2k-1} e_3 \in \mathbb{Z} \oplus_{2k+2} MO(2k): \quad \square$$

The map  $h^2: H_n QMO(k) \rightarrow H_n D_2 MO(k)$  in Diagram (2) is very easy to describe in terms of the description of  $H QX$  as a Pontrjagin ring provided by Dyer and Lashof (see [9] or [17]). They make use of the Kudo-Araki operations  $Q^i: H_m QX \rightarrow H_{m+i} QX$ . These are trivial for  $i < m$  and equal to the Pontrjagin square for  $i = m$ . If  $I$  denotes the sequence  $(i_1; i_2; \dots; i_r)$  then we write  $Q^I x = Q^{i_1} Q^{i_2} \dots Q^{i_r} x$ . The sequence  $I$  is *admissible* if  $i_j \leq i_{j+1}$  for  $1 \leq j < r$  and its *excess* is given by  $e(I) = i_1 - i_2 - \dots - i_r$ . With this notation we can give the description of  $H QX$  as a polynomial algebra: if  $f: X \rightarrow \mathbb{Z} \oplus_{2k+2} MO(k)$  is a homogeneous basis for  $H X \rightarrow H QX$  where  $X$  is a path-connected space then

$$H QX = \mathbb{Z}\langle Q^I x_j \mid I \text{ admissible of excess } e(I) > \dim x_j \rangle$$

Thus a basis for  $H QX$  is provided by the monomials in the polynomial generators.

We may define a height function  $ht$  on the monomial generators of  $H QX$  by  $ht(x_j) = 1$ ,  $ht(Q^i u) = 2ht(u)$  and  $ht(u \smile v) = ht(u) + ht(v)$  (where  $u \smile v$  represents the Pontrjagin product). The following is a special case of Lemma 2.3 in [2].

**Lemma 2.2** The homomorphism  $h^2: H QX \rightarrow H D_2 X$  is given by projection onto the monomial generators of height 2. The kernel is spanned by the set of monomials of height other than 2. □

**Corollary 2.3** A basis for  $H_{2k+2} D_2 MO(k)$  is given by the following set:

$$\{e_1^k, e_1^{k-1} e_3, e_1^k e_1^{k-2} e_2^2, e_1^{k-1} e_2, e_1^{k-1} e_2, Q^{k+2} e_1^k\} \quad \square$$

The map  $h^2: H_n D_2 MO(k) \rightarrow H_n MO(2k)$  of Diagram (2) is also determined in [2]. In the case of  $n = 2k + 2$ , Theorem 3.1 of [2] gives the following.

**Lemma 2.4** The homomorphism  $h^2: H_{2k+2} D_2 MO(k) \rightarrow H_{2k+2} MO(2k)$  is determined by the following values:

$$\begin{aligned} (e_1^k, e_1^{k-1} e_3) &= e_1^{2k-1} e_3; \\ (e_1^k, e_1^{k-2} e_2^2) &= (e_1^{k-1} e_2, e_1^{k-1} e_2) = e_1^{2k-2} e_2^2; \end{aligned}$$

$$(Q^{k+2}e_1^k) = \begin{cases} 0 & \text{for } k \equiv 0 \pmod{4}; \\ e_1^{2k-1}e_3 & \text{for } k \equiv 1 \pmod{4}; \\ e_1^{2k-2}e_2^2 & \text{for } k \equiv 2 \pmod{4}; \\ e_1^{2k-1}e_3 + e_1^{2k-2}e_2^2 & \text{for } k \equiv 3 \pmod{4}; \end{cases}$$

**Proof** The first three results are immediate from [2] Theorem 3.1.

For the fourth we apply the formula given in the theorem:

$$\begin{aligned} (Q^{k+2}e_1^k) &= \sum_{m_1+m_2+\dots+m_k=2} \prod_{j=1}^k \binom{m_j-1}{0} e_1 e_{m_j+1} \quad (m_j \geq 0) \\ &= \sum_{m_1+m_2+\dots+m_k=2} e_1 e_{m_1+1} e_1 e_{m_2+1} \dots e_1 e_{m_k+1} \quad (m_j \geq 0) \\ &= \binom{k}{1} e_1^{2k-1}e_3 + \binom{k}{2} e_1^{2k-2}e_2^2 \end{aligned}$$

which gives the required result. □

To prove Theorem 1.1 using Theorem 2.1 we determine the image of the spherical classes in  $H_{2k+2}QMO(k)$ , ie, the classes in the image of  $h: H_{2k+2}QMO(k) \rightarrow H_{2k+2}QMO(k)$ , under  $h^2$  (see Diagram (2)). A complete description of these spherical classes is not necessary for it is sufficient to observe the following well-known properties of spherical classes (which are immediate from  $H^*S^n$  by naturality).

**Lemma 2.5** (a) *If an homology class  $u \in H_n X$  is spherical then it is primitive with respect to the cup coproduct, ie,*

$$(u) = u \otimes 1 + 1 \otimes u;$$

where  $\Delta: H_n(X \times X) \rightarrow \bigoplus_i H_i X \otimes H_{n-i} X$  is the map induced by the diagonal map.

(b) *If an homology class  $u \in H_n X$  is spherical (or stably spherical, ie, in the image of  $h^S: H_n^S X \rightarrow H_n X$ ) then it is annihilated by the reduced Steenrod algebra, ie,*

$$Sq^i(u) = 0$$

for all  $i > 0$ , where  $Sq^i: H_n X \rightarrow H_{n-i} X$  is the vector space dual of the usual Steenrod square cohomology operation  $Sq^i: H^{n-i} X \rightarrow H^n X$ . □

To apply the first of these observations we determine the image of the coproduct primitive submodule of  $H_{2k+2}QMO(k)$  in  $H_{2k+2}D_2MO(k)$ .



**Lemma 2.6** Suppose that  $k > 2$ . Then a basis for the coproduct primitive submodule  $PH_{2k+2}QMO(k)$  is given by the following set of elements:

$$f e_1 e_{i_2} \cdots e_{i_k} j \quad i_2 \quad \cdots \quad i_k g [ f e_2^{k-2} e_3^2 + e_1^k e_1^{k-2} e_2^2; e_1^{k-1} e_2 e_1^{k-1} e_2; Q^{k+2} e_1^k g;$$

For  $k = 1$ , a basis for  $PH_4QMO(1)$  is given by

$$f Q^3 e_1; e_1 e_1 e_1 e_1 g;$$

For  $k = 2$ , a basis for  $PH_6QMO(2)$  is given by

$$f e_1 e_5; e_3^2 + e_1^2 e_2^2 + e_1^2 e_1^2 e_1^2; e_1 e_2 e_1 e_2; Q^4 e_1^2 g;$$

**Proof** Recall that  $(e_i) = \prod_j e_j e_{i-j}$ . This determines  $(e_i)$  by naturality.

First of all observe that a basis element  $e_{i_1} e_{i_2} \cdots e_{i_k}$  ( $i_1 \quad i_2 \quad \cdots \quad i_k$ ) of  $HMO(k)$  is primitive if and only if  $i_1 = 1$ . To complete the proof we evaluate the coproduct on the non-primitive height 1 basis elements and on all the basis elements of greater height in this dimension.

For  $k > 2$  there are no basis elements of height greater than 2. For simplicity we use the reduced coproduct  $\tilde{\sim}(u) = (u) - u \quad 1 - 1 \quad u$  so that  $u$  is primitive when  $\tilde{\sim}(u) = 0$ . Straightforward calculations give the following:

$$\begin{aligned} \tilde{\sim}(e_2^{k-2} e_3^2) &= e_1^k e_1^{k-2} e_2^2 + e_1^{k-2} e_2^2 e_1^k; \\ \tilde{\sim}(e_2^{k-1} e_4) &= e_1^k e_1^{k-1} e_3 + e_1^{k-1} e_2 e_1^{k-1} e_2 + e_1^{k-1} e_3 e_1^k; \\ \tilde{\sim}(e_1^k e_1^{k-1} e_3) &= e_1^k e_1^{k-1} e_3 + e_1^{k-1} e_3 e_1^k; \\ \tilde{\sim}(e_1^k e_1^{k-2} e_2^2) &= e_1^k e_1^{k-2} e_2^2 + e_1^{k-2} e_2^2 e_1^k; \\ \tilde{\sim}(e_1^{k-1} e_2 e_1^{k-1} e_2) &= 0; \\ \tilde{\sim}(Q^{k+2} e_1^k) &= 0. \end{aligned}$$

The lemma follows immediately from these results.

Similar calculations give the results for  $k = 1$  and  $k = 2$ . □

This lemma has the following immediate corollary.

**Corollary 2.7** For  $k > 1$ , a basis for the projection of the coproduct primitive submodule  $h^2 PH_{2k+2}QMO(k) \quad H_{2k+2}D_2MO(k)$  is given by the following set of elements:

$$f e_1^k e_1^{k-2} e_2^2; e_1^{k-1} e_2 e_1^{k-1} e_2; Q^{k+2} e_1^k g;$$

For  $k = 1$  a basis for  $h^2 PH_4QMO(1)$  is given by  $f Q^3 e_1 g$ . □

To complete the proof of the main theorem it is convenient to consider various cases depending on the value of  $k$  modulo 4.

### 3 Case 1: $k$ even

**Theorem 3.1** For even  $k$ , the double point self-intersection surface of each self-transverse immersion  $f: M^{k+2} \looparrowright \mathbb{R}^{2k+2}$  has even Euler characteristic and so is a boundary.

To prove this result we evaluate dual Steenrod operations on the elements given by Corollary 2.7.

**Lemma 3.2** For even  $k$ , we have the following results in  $H D_2 MO(k)$ :

$$\begin{aligned} Sq^1(e_1^k e_1^{k-2} e_2^2) &= Sq^1(e_1^{k-1} e_2 e_1^{k-1} e_2) = 0; \\ Sq^1(Q^{k+2} e_1^k) &= Q^{k+1} e_1^k; \\ Sq^2(e_1^k e_1^{k-2} e_2^2) &= Sq^2(e_1^{k-1} e_2 e_1^{k-1} e_2) = e_1^k e_1^k. \end{aligned}$$

**Proof** These are immediate from the Steenrod squares in  $BO(1) = \mathbb{R}P^1$  ( $Sq^i e_j = \binom{j-i}{i} e_{j-i}$ ) and the Nishida relations (see [17]).  $\square$

**Corollary 3.3** For even  $k$ , given  $2 \in \mathcal{S}_{2k+2} MO(k)$ ,

$$h^2 h(\ ) = (e_1^k e_1^{k-2} e_2^2 + e_1^{k-1} e_2 e_1^{k-1} e_2)$$

for  $2 \in \mathbb{Z}=2$ .

**Proof** By Lemma 2.5(a)  $h^2 h(\ )$  is a linear combination of the elements given by Corollary 2.7. But since, by Lemma 2.5(b), it is annihilated by  $Sq^1$  and  $Sq^2$  only elements of the given form can arise.  $\square$

**Proof of Theorem 3.1** Suppose that  $f$  corresponds to  $2 \in \mathcal{S}_{2k+2} QMO(k)$ . Then

$$h^2 h(\ ) = (e_1^k e_1^{k-2} e_2^2 + e_1^{k-1} e_2 e_1^{k-1} e_2) = (e_1^{2k-2} e_2^2 + e_1^{2k-2} e_2^2) = 0$$

by Lemma 2.4.

The result follows by Theorem 2.1.  $\square$

We can still ask whether there exists an immersion  $f: M^{k+2} \looparrowright \mathbb{R}^{2k+2}$  corresponding to  $2 \in \mathcal{S}_{2k+2} MO(k)$  with  $h^2 h(\ ) = e_1^k e_1^{k-2} e_2^2 + e_1^{k-1} e_2 e_1^{k-1} e_2$ . To answer this we first observe the following result.

**Proposition 3.4** For even  $k$ , given an immersion  $f: M^{k+2} \looparrowright \mathbb{R}^{2k+2}$  corresponding to  $\alpha \in \mathcal{S}_{2k+2} MO(k)$ , the Hurewicz image  $h(\alpha) \in H_{2k+2} QMO(k)$  is determined by the bordism class of  $M$ .

**Proof** Suppose that  $k > 2$ . Notice that the height 1 part of  $h(\alpha)$  is given by  $h^S(\alpha) \in H_{2k+2} MO(k) \subset H_{2k+2} QMO(k)$  (see Diagram (1) in [2]) and so

$$h(\alpha) = h^S(\alpha) + h^2 h(\alpha)$$

using the obvious inclusion  $H \subset D_2 MO(k) \subset H \subset QMO(k)$  as the elements of height 2. Furthermore,  $h^S(\alpha)$  is determined by the normal Stiefel-Whitney numbers of  $M$  and so by the bordism class of  $M$  (see [2] Lemma 2.2).

Suppose that  $f_1: M_1^{k+2} \looparrowright \mathbb{R}^{2k+2}$  and  $f_2: M_2^{k+2} \looparrowright \mathbb{R}^{2k+2}$  are two immersions of bordant manifolds  $M_1$  and  $M_2$  corresponding to  $\alpha_1, \alpha_2 \in \mathcal{S}_{2k+2} MO(k)$  respectively. Since  $M_1$  and  $M_2$  are bordant manifolds  $h^S(\alpha_1) = h^S(\alpha_2)$ . It follows that

$$\begin{aligned} h(\alpha_1) - h(\alpha_2) &= h^2 h(\alpha_1) - h^2 h(\alpha_2) \\ &= h^2 h(\alpha_1 - \alpha_2) \\ &= (e_1^k - e_1^{k-2} e_2^2 + e_1^{k-1} e_2 - e_1^{k-1} e_2) \end{aligned}$$

by Corollary 3.3. However  $e_1^k - e_1^{k-2} e_2^2 + e_1^{k-1} e_2 - e_1^{k-1} e_2 \in H_{2k+2} QMO(k)$  is not primitive (by Lemma 2.6) and so not spherical.

Hence  $h(\alpha_1) - h(\alpha_2) = 0$  and so  $h(\alpha_1) = h(\alpha_2)$  as required.

For  $k = 2$  the above proof has to be modified to take account of height 3 elements. This leads in this case to

$$h(\alpha_1) - h(\alpha_2) = (e_1^2 - e_2^2 + e_1 e_2 - e_1 e_2 + e_1^2 - e_1^2 - e_1^2)$$

which gives the same result. □

From now on we write  $\nu_1(m)$  for the number of digits 1 in the dyadic expression for the positive integer  $m$ .

**Proposition 3.5** For even  $k > 2$ , given an immersion  $f: M^{k+2} \looparrowright \mathbb{R}^{2k+2}$  corresponding to  $\alpha \in \mathcal{S}_{2k+2} MO(k)$ , the Hurewicz image  $h(\alpha) \in H_{2k+2} QMO(k)$  is given as follows:

$$h(\alpha) = \begin{cases} h^S(\alpha); & \text{if the normal Stiefel-Whitney number } \overline{w}_2 \overline{w}_k[M] = 0; \\ h^S(\alpha) + e_1^k - e_1^{k-2} e_2^2 + e_1^{k-1} e_2 - e_1^{k-1} e_2; & \text{if } \overline{w}_2 \overline{w}_k[M] = 1. \end{cases}$$

The second case arises if and only if  $\nu_1(k+2) \geq 2$ . In this case, writing  $k+2 = 2^r + 2^s$  where  $r \geq s \geq 1$ , we can take  $M$  to be  $\mathbb{R}P^{2^r} \times \mathbb{R}P^{2^s}$ .

For  $k = 2$ , given  $f: M^4 \looparrowright \mathbb{R}^6$  corresponding to  $\in \mathbb{S}^2 MO(2)$ , the Hurewicz image  $h(\ ) \in H_6 QMO(2)$  is given as follows:

$$h(\ ) = \begin{cases} 0; & \text{if } \mathbb{W}_2^2[M] = 0, \\ e_3^2 + e_1^2 e_2^2 + e_1 e_2 e_1 e_2 + e_1^2 e_1^2 e_1^2; & \text{if } \mathbb{W}_2^2[M] = 1. \end{cases}$$

**Proof** Suppose that  $k > 2$  and that  $f: M^{k+2} \looparrowright \mathbb{R}^{2k+2}$  is an immersion corresponding to  $\in \mathbb{S}^2_{2k+2} MO(k)$ .

If  $\in = 0$  in Corollary 3.3 then  $h(\ ) = h^S(\ )$ . By Lemma 2.6,  $\in = 1$  if and only if  $e_2^{k-2} e_3^2$  has coefficient 1 when  $h^S(\ )$  is written in terms of the basis  $f e_i g$ . This occurs if and only if the Stiefel-Whitney number  $\mathbb{W}_2 \mathbb{W}_k[M] = 1$ . For, by [2] Lemma 2.2,  $\mathbb{W}_2 \mathbb{W}_k[M]$  is given by the Kronecker product  $h \mathbb{W}_2 \mathbb{W}_k^2; h^S(\ ) i$  in  $MO(k)$ . By naturality, this product can be evaluated in  $BO(1)^k$  using  $\binom{k}{i}$  and the fact that  $\binom{k}{i} \mathbb{W}_i = \sum_{j=0}^i \binom{k-1}{j} \mathbb{W}_j \mathbb{W}_{i-j} \in \mathbb{Z}[x_1, x_2, \dots, x_k] \cong \mathbb{Z}[BO(1)^k]$ , the  $i$ th elementary symmetric polynomial. In this case  $\binom{k}{i} (\mathbb{W}_2 \mathbb{W}_k^2) = x_1^3 x_2^3 x_3^2 \dots x_k^2$  and so the Kronecker product  $h \mathbb{W}_2 \mathbb{W}_k^2; h^S(\ ) i$  is given by the coefficient of  $e_2^{k-2} e_3^2$ . Hence  $\in = 1$  if and only if  $\mathbb{W}_2 \mathbb{W}_k[M] = 1$ .

In the case  $(k + 2) > 2$ , by a theorem of R L W Brown ([6] Theorem 5.1), there exists an embedding  $f_1: M_1 \hookrightarrow \mathbb{R}^{2k+2}$  of a manifold  $M_1$  bordant to  $M$ . Suppose that  $f_1$  corresponds to  $\in_1 \in \mathbb{S}^2_{2k+2} MO(k)$ . Then  $h^2 h(\in_1) = 0$  since  $f_1$  is an embedding with no double points. Hence, by Proposition 3.4,  $h^2 h(\in) = 0$  and so  $h(\ ) = h^S(\ )$ .

For  $(k + 2) \geq 2$  put  $k + 2 = 2^r + 2^s$  where  $r \leq s - 1$ . Then by Whitney's immersion theorem there are immersions  $\mathbb{R}P^{2^r} \looparrowright \mathbb{R}^{2^{r+1}-1}$  and  $\mathbb{R}P^{2^s} \looparrowright \mathbb{R}^{2^{s+1}-1}$  and the product of these gives an immersion  $f: \mathbb{R}P^{2^r} \times \mathbb{R}P^{2^s} \looparrowright \mathbb{R}^{2^{r+1}+2^{s+1}-2} = \mathbb{R}^{2k+2}$ . Finally a standard verification shows that the normal Stiefel-Whitney number  $\mathbb{W}_2 \mathbb{W}_{2^r+2^s-2}[\mathbb{R}P^{2^r} \times \mathbb{R}P^{2^s}] = 1$ .

The argument for  $k = 2$  is almost identical. The manifold  $M^4$  either is a boundary or is bordant to  $\mathbb{R}P^2 \times \mathbb{R}P^2$  depending on the value of the normal Stiefel-Whitney number  $\mathbb{W}_2^2[M]$ . The presence of the height 3 term  $e_1^2 e_1^2 e_1^2$  in the Hurewicz image shows that in this case any immersion of a non-boundary has an odd number of triple points.  $\square$

### 4 Case 2: $k \equiv 1 \pmod 4$

**Theorem 4.1** For  $k \equiv 1 \pmod 4$ , given any manifold  $M^{k+2}$  there exists an immersion  $M^{k+2} \looparrowright \mathbb{R}^{2k+2}$  with double point self-intersection surface of odd

Euler characteristic (and so a non-boundary) and there exists another immersion with double point self-intersection surface of even Euler characteristic (and so a boundary).

**Proof** For  $k \equiv 1 \pmod{4}$ ,  $(k + 2) \equiv 2$  and so there exists an immersion  $f: M^{k+2} \looparrowright \mathbb{R}^{2k+2}$  by R Cohen's immersion theorem ([7]). Furthermore, for such  $k$  there is an immersion  $S^{k+2} \looparrowright \mathbb{R}^{2k+2}$  with double point manifold of odd Euler characteristic ([13] Theorem 1.2). Taking the connected sum of this immersion and  $f$  gives an immersion of  $M$  with a double point self-intersection surface with Euler characteristic of opposite parity to that of the double point self-intersection surface of  $f$ . Hence both parities can arise.  $\square$

Again it is natural to ask about the Hurewicz image  $h(\ ) \in H_{2k+2}MO(k)$  of the elements  $\in \Sigma_{2k+2}MO(k)$  corresponding to these immersions. The situation is the following.

**Proposition 4.2** For  $k \equiv 1 \pmod{4}$  such that  $k > 1$ , given an immersion  $f: M^{k+2} \looparrowright \mathbb{R}^{2k+2}$  corresponding to  $\in \Sigma_{2k+2}MO(k)$ , then the Hurewicz image  $h(\ ) \in H_{2k+2}QMO(k)$  is given as follows:

$$h(\ ) = \begin{cases} h^S(\ ); & \text{if } \chi_2(f) \text{ has even Euler characteristic,} \\ h^S(\ ) + Q^{k+2}e_1^k; & \text{if } \chi_2(f) \text{ has odd Euler characteristic.} \end{cases}$$

For  $k = 1$ , given an immersion  $f: M^3 \looparrowright \mathbb{R}^4$  corresponding to  $\in \Sigma_4MO(1)$ , then the Hurewicz image  $h(\ ) \in H_4QMO(1)$  is given as follows:

$$h(\ ) = \begin{cases} 0; & \text{if } \chi_2(f) \text{ has even Euler characteristic,} \\ Q^3e_1 + e_1e_1e_1e_1; & \text{if } \chi_2(f) \text{ has odd Euler characteristic.} \end{cases}$$

**Proof** Evaluating the dual Steenrod operations on the elements of Corollary 2.7 we obtain

$$\begin{aligned} Sq^1(e_1^k e_1^{k-2} e_2^2) &= Sq^1(e_1^{k-1} e_2 e_1^{k-1} e_2) = Sq^1(Q^{k+2} e_1^k) = 0; \\ Sq^2(e_1^k e_1^{k-2} e_2^2) &= Sq^2(e_1^{k-1} e_2 e_1^{k-1} e_2) = e_1^k e_1^k; \\ Sq^2(Q^{k+2} e_1^k) &= 0. \end{aligned}$$

Hence  $h^2 h(\ )$  is in the submodule spanned by the set

$$f e_1^k e_1^{k-2} e_2^2 + e_1^{k-1} e_2 e_1^{k-1} e_2; Q^{k+2} e_1^k g;$$

For  $k \equiv 1 \pmod{4}$  and  $k > 1$  it follows that  $(k + 2) > 2$ ; so, by RLW Brown's embedding theorem ([6] Theorem 5.1),  $M$  is bordant to a manifold  $M_1$  which

has an embedding  $f_1: M_1 \hookrightarrow \mathbb{R}^{2k+2}$ . Let  $f_1$  correspond to  $\alpha \in \mathbb{Z}/2^{k+2}MO(k)$ . Then  $h^2 h(\alpha) = 0$  since  $f_1$  is an embedding. Furthermore,  $h^S(\alpha) = h^S(\alpha)$  since these are determined by the bordism class. It follows, by Lemma 2.6, that  $h^2 h(\alpha) = h(\alpha - \beta) = Q^{k+2}e_1^k$  since this is necessarily a primitive class in  $H_{2k+2}QMO(k)$ . Hence  $h(\alpha) = h^S(\alpha) + Q^{k+2}e_1^k$  where, by Theorem 2.1 and Lemma 2.4,  $\alpha$  gives the parity of the Euler characteristic of the double point self-intersection surface of the immersion.

For  $k = 1$ , given any immersion  $f: M^3 \looparrowright \mathbb{R}^4$  the manifold  $M$  is necessarily a boundary since this is true of all 3-manifolds and so, if  $\alpha \in \mathbb{Z}/4MO(1) = \mathbb{Z}/2$  is the corresponding element,  $h^S(\alpha) = 0$ . The formula for  $h(\alpha)$  then follows from the value for the Hurewicz image coming from [10] Proposition 3.4. The presence of the height 4 term  $e_1 e_1 e_1 e_1$  when  $\alpha_2(f)$  has odd Euler characteristic indicates that such immersions have an odd number of quadruple points. □

### 5 Case 3: $k \equiv 3 \pmod 4$

**Theorem 5.1** *For  $k \equiv 3 \pmod 4$ , there exists a manifold  $M^{k+2}$  with a self-transverse immersion  $M^{k+2} \looparrowright \mathbb{R}^{2k+2}$  for which the Euler characteristic of the double point self-intersection surface is odd (so that it is a non-boundary) if and only if  $k + 1$  is a power of 2. In this case, the parity of the Euler characteristic of the double point self-intersection surface of a self-transverse immersion  $M^{k+2} \looparrowright \mathbb{R}^{2k+2}$  is given by the normal Stiefel-Whitney number  $\mathbb{W}_2 \mathbb{W}_k[M]$ .*

To prove this we begin as in Case 1.

**Lemma 5.2** *For  $k \equiv 3 \pmod 4$ , we have the following results in  $H D_2 MO(k)$ :*

$$\begin{aligned} Sq^1(e_1^k e_1^{k-2} e_2^2) &= Sq^1(e_1^{k-1} e_2 e_1^{k-1} e_2) = Sq^1(Q^{k+2} e_1^k) = 0; \\ Sq^2(e_1^k e_1^{k-2} e_2^2) &= Sq^2(e_1^{k-1} e_2 e_1^{k-1} e_2) = Sq^2(Q^{k+2} e_1^k) = e_1^k e_1^k. \quad \square \end{aligned}$$

**Corollary 5.3** *For  $k \equiv 3 \pmod 4$ , given  $\alpha \in \mathbb{Z}/2^{k+2}MO(k)$ ,  $h^2 h(\alpha)$  lies in the submodule of  $H_{2k+2} D_2 MO(k)$  spanned by the following set:*

$$f e_1^k e_1^{k-2} e_2^2 + e_1^{k-1} e_2 e_1^{k-1} e_2; e_1^k e_1^{k-2} e_2^2 + Q^{k+2} e_1^k; g \quad \square$$

To prove Theorem 5.1 we determine which of the elements in the submodule of Corollary 5.3 can arise as a value of  $h^2 h$ . First of all we observe that, for



and so lies in the submodule of  $H_{2k+2}QMO(k)$  spanned by the set given by Corollary 5.3.

We consider the non-zero elements of this submodule in turn. First of all, by Lemma 2.6, neither  $e_1^k - e_1^{k-2}e_2^2 + e_1^{k-1}e_2 - e_1^{k-1}e_2$  nor  $e_1^k - e_1^{k-2}e_2^2 + Q^{k+2}e_1^k$  is primitive and so neither of these elements can be spherical. This leaves the element  $e_1^{k-1}e_2 - e_1^{k-1}e_2 + Q^{k+2}e_1^k$ , which is primitive.

**Lemma 5.7** *The element  $e_1^{k-1}e_2 - e_1^{k-1}e_2 + Q^{k+2}e_1^k \in H_{2k+2}QMO(k)$  is not spherical.*

Accepting this for the moment, it follows that  $h(\alpha) - h(\beta) = h(\alpha - \beta) = 0$  and so  $h(\alpha) = h(\beta)$  as required.  $\square$

The proof of Lemma 5.7 needs an additional idea. Certain elements  $\in H_{2n}QX = \mathbb{S}_n X$  are detected by their Hurewicz image in  $H_n QX$ . Elements may also be detected by a cohomology operation in  $H^*C$  where the stable space  $C$  is the mapping cone of the stable map  $\rho : S^n \rightarrow X$ . It is interesting to consider the relationship between these two approaches. One of the most basic facts is the following.

**Proposition 5.8** *Given  $\alpha \in H_{2m}QX = \mathbb{S}_{2m} X$ , if  $h(\alpha) = u^2 \in H_{2m}QX$  where  $u \in H_m X$ , then  $Sq^{m+1}\bar{u} \neq 0 \in H^{2m+1}C$ , where  $\bar{u} \in H^m C$  is any class such that the Kronecker product  $hi \bar{u}; ui = 1$  writing  $i : X \rightarrow C$  for the natural inclusion map.*

Notice that  $i Sq^{m+1}\bar{u} = Sq^{m+1}i \bar{u} = 0 \in H^{2m+1}X$  for dimensional reasons and so  $Sq^{m+1}\bar{u}$  lies in  $\rho^* H^{2m+1}S^{2m+1} = \mathbb{Z}/2$  where  $\rho : C \rightarrow S^{2m+1}$  is the natural projection map. When  $Sq^{m+1}\bar{u} \neq 0$  the stable space  $C$  requires at least one suspension before it can exist as an unstable space.

**Proof of Proposition 5.8** This proposition can be thought of as describing how detection by a certain James-Hopf invariant corresponds to detection by a Steenrod-Hopf invariant. The proposition follows from work of Boardman and Steer ([5] Corollary 5.15). The special case of  $X = \mathbb{R}P^1$  appears as [11] Lemma 4.2 and the justification given there extends to the general case.  $\square$

**Proof of Lemma 5.7** Suppose, for contradiction, that there exists an element  $\alpha \in H_{2k+2}QMO(k)$  such that

$$h(\alpha) = e_1^{k-1}e_2 - e_1^{k-1}e_2 + Q^{k+2}e_1^k \in H_{2k+2}QMO(k):$$



Adjointness gives a natural isomorphism  $\tau : {}_n QX = {}_{n+1} Q X$ . Since the corresponding homology suspension  $\tau : H_n QX \rightarrow H_{n+1} Q X$  kills Pontrjagin products and commutes with the Kudo{Araki operations,

$$h(\tau) = Q^{k+2} \tau e_1^k = \tau e_1^k \tau e_1^k \in H_{2k+4} Q \tau MO(k):$$

Hence, by Proposition 5.8,

$$Sq^{k+3} \tau \neq 0 \in H^{2k+5} C_2 \tag{3}$$

where  $\tau \in H^{k+2} C_2$  is an element such that  $hi \tau; \tau e_1^k i = 1$ .

We now show that this is impossible.

First of all notice that, since  $h^S(\tau) = 0 \in H_{2k+4} \tau MO(k)$ ,  $i : H^i C_2 = H^i \tau MO(k)$  for  $i \notin 2k + 5$ . This means that away from dimension  $2k + 5$  we can do calculations in  $H C_2$  by calculating in  $H \tau MO(2)$ . In particular,  $H^{k+2} C_2 = H^{k+2} \tau MO(k) = \mathbb{Z} = 2$  is generated by  $\tau w_k$  and so  $\tau = \tau w_k$ . Here  $w_k \in H^k MO(k) = H^k BO(k)$  is the universal Stiefel{Whitney class.

Secondly, we can evaluate the action of the Steenrod squaring operations on  $H \tau MO(k)$  using the fact that the operations commute with suspension and using their action on  $H MO(k)$  which is determined by the Wu formula (see [18] Problem 8A). In evaluating the operations we use the fact that  $k \equiv 3 \pmod 4$  which means that we can write  $k = 4r - 1$  for some positive integer  $r$ .

Then,

$$\begin{aligned} Sq^{k+3} \tau &= Sq^{4r+2} \tau w_{4r-1} \\ &= (Sq^2 Sq^{4r} + Sq^1 Sq^{4r} Sq^1) \tau w_{4r-1} \text{ by the Adem relations} \\ &= Sq^2 \tau Sq^{4r} w_{4r-1} + Sq^1 \tau Sq^{4r} Sq^1 w_{4r-1} \\ &\quad \text{since the operations commute with suspension} \\ &= 0 + Sq^1 \tau w_1^2 w_{4r-1}^2 \text{ by the Wu formula and dimension} \\ &= Sq^1 \tau Sq^1 w_1 w_{4r-1}^2 \text{ by a further application of the Wu formula} \\ &= Sq^1 Sq^1 \tau w_1 w_{4r-1}^2 \text{ since } Sq^1 \text{ commutes with suspension} \\ &= 0 \text{ by the Adem relations.} \end{aligned}$$

This contradicts equation (3) and so proves the lemma, completing the proof of Proposition 5.4 and so the proof of Theorem 5.1 in the case of  $(k + 2) > 2$ .  $\square$

Now suppose that  $k \equiv 3 \pmod 4$  and  $(k + 2) = 2$ . In this case  $k + 1 = 2^r$  for some integer  $r$  such that  $r \geq 2$ .

To complete the proof of Theorem 5.1 we consider the structure of the bordism group  $MO_{k+2}$  of  $(k + 2)$ -dimensional manifolds. Recall that the unoriented

bordism ring  $MO$  is a polynomial ring with one generator in each dimension not of the form  $2^s - 1$ . We describe a manifold  $M^{k+2}$  as decomposable (up to bordism) if it represents a decomposable element in this ring and so is bordant to a union of product manifolds  $N_1^{m_1} \times N_2^{m_2}$  where  $m_1, m_2 > 0$ . We can assume that  $m_1, m_2 \geq 2$  since there is no non-boundary in dimension 1. Any indecomposable manifold of dimension  $k + 2$  can be taken as a representative of a polynomial generator in this dimension and any two such manifolds are bordant modulo decomposable manifolds. Notice that, by R Cohen's immersion theorem ([7]), every manifold in this dimension immerses in  $\mathbb{R}^{2k+2}$ .

We first of all deal with the decomposable manifolds.

**Proposition 5.9** *For  $k \equiv 3 \pmod{4}$  such that  $(k + 2) = 2$ , every manifold  $M^{k+2}$  which is decomposable in the bordism ring is bordant to a manifold which embeds in  $\mathbb{R}^{2k+2}$ . Hence the double point self-intersection surface of any immersion  $M^{k+2} \hookrightarrow \mathbb{R}^{2k+2}$  has even Euler characteristic and so is a boundary.*

**Proof** To prove that decomposable manifolds of dimension  $k + 2$  embed up to bordism it is sufficient to prove that each product  $N_1^{m_1} \times N_2^{m_2}$ , with  $m_1, m_2 \geq 2$  and  $m_1 + m_2 = k + 2$ , embeds up to bordism in  $\mathbb{R}^{2k+2}$ . To do this we make use of the following result which follows easily from the Whitney embedding theorem.

**Lemma 5.10** ([6] Lemma 2.1) *If the manifold  $N_1^{m_1}$  immerses in  $\mathbb{R}^s$ , the manifold  $N_2^{m_2}$  embeds in  $\mathbb{R}^t$  and  $s + t \geq 2m_1 + 1$  then  $N_1 \times N_2$  embeds in  $\mathbb{R}^{s+t}$ .  $\square$*

Since  $m_1 + m_2 = k + 2 = 2^r + 1$  and  $m_1, m_2 \geq 2$  it follows that  $(m_1) \geq 2$  or  $(m_2) \geq 2$ . Suppose without loss of generality that  $(m_2) \geq 2$ . Then, by R.L.W. Brown's embedding theorem ([6]),  $N_2$  is bordant to a manifold which embeds in  $\mathbb{R}^{2m_2-1}$ . In addition, by the Whitney immersion theorem,  $N_1$  immerses in  $\mathbb{R}^{2m_1-1}$ . Hence, by the lemma,  $N_1 \times N_2$  is bordant to a manifold which embeds in  $\mathbb{R}^{2m_1+2m_2-2} = \mathbb{R}^{2k+2}$ . The numerical condition of the lemma is automatically satisfied:  $s + t = 2k + 2 \geq 2m_1 + 1$  since  $m_1 \leq k$ .

The final part of the proposition is immediate from Corollary 5.5 as in the proof of Corollary 5.6.  $\square$

We turn now to the indecomposable manifolds.



manifold the same formula holds for an immersion of  $M$  by Proposition 5.4 and Proposition 5.9.

The Euler characteristic of the double point self-intersection surface is now given by Theorem 2.1 since, by Lemma 2.4,  $(e_1^k - e_1^{k-2}e_2^2 + Q^{k+2}e_1^k) = e_1^{2k-1}e_3$ .  $\square$

**Proof of Theorem 5.1** Almost everything is given by Corollary 5.6, Proposition 5.9 and Proposition 5.11. The final observation about the Stiefel-Whitney number is clear since, by Lemma 2.6,  $e_1^k - e_1^{k-2}e_2^2$  occurs in  $h(\ )$  if and only if  $e_2^{k-2}e_3^2$  does and by the argument in the proof of Proposition 3.5 this is equivalent to  $\mathbb{W}_2\mathbb{W}_k[M] = 1$ .  $\square$

## 6 Comparison with Szűcs's results

To conclude, we compare the results and methods of this paper with those of Szűcs in [22]. His methods are more geometric than ours and he gives explicit constructions to show that any surface of even Euler characteristic can occur as the double point self-intersection manifold of a self-transverse immersion  $M^{k+2} \looparrowright \mathbb{R}^{2k+2}$  if  $k \equiv 1 \pmod{4}$ , and any surface can arise if  $k \equiv 1 \pmod{4}$ .

There are two key steps in Szűcs's non-existence proof.

**Proposition 6.1** ([22] Claim 1) *For  $k \not\equiv 1 \pmod{4}$ , the parity of the Euler characteristic of the double point self-intersection surface of a self-transverse immersion  $M^{k+2} \looparrowright \mathbb{R}^{2k+2}$  depends only on the bordism class of  $M$ .*

**Proposition 6.2** ([22] Claim 3) *For  $k \not\equiv 1 \pmod{4}$ , the Euler characteristic of the double point self-intersection surface of a self-transverse immersion  $M^{k+2} \looparrowright \mathbb{R}^{2k+2}$  is even, if  $M$  is decomposable in the bordism ring.*

These are both included in our results. In fact we have proved a stronger result than Proposition 6.1 as follows.

**Proposition 6.3** *For  $k \not\equiv 1 \pmod{4}$ , given an element  $\alpha \in \mathbb{S}_{2k+2}MO(k)$  corresponding to a self-transverse immersion  $M^{k+2} \looparrowright \mathbb{R}^{2k+2}$ , the Hurewicz image  $h(\alpha) \in H_{2k+2}QMO(k)$  depends only on the bordism class of  $M$ .*

This is our Proposition 3.4 and Proposition 5.4. It implies Proposition 6.1 by Theorem 2.1.

Szucs's technique for proving Proposition 6.1 is to extend the discussion to include maps which he calls *prim* maps; these are smooth maps  $M^{k+2} \rightarrow \mathbb{R}^{2k+2}$  which arise as the projection of an immersion  $M^{k+2} \hookrightarrow \mathbb{R}^{2k+3} \rightarrow \mathbb{R}^{2k+2}$ . Szucs claims that Proposition 6.1 remains true for these maps. However, this must be false since he describes ([22] Lemma 4) how to construct a prim map  $f: V^{2r+1} \rightarrow \mathbb{R}^{2r+1}$  of an indecomposable manifold with a double point self-intersection surface of even Euler characteristic, whereas by our Proposition 5.11 the double point self-intersection surface of any immersion of  $V$  will have odd Euler characteristic.

Turning to Proposition 6.2, we have proved this explicitly for  $k \equiv 3 \pmod{4}$  as Proposition 5.9. It is immediate for even  $k$  from Theorem 3.1. It is intriguing that, whereas by our methods the case of even  $k$  is the simplest, by Szucs's geometric approach the argument in this case is quite delicate. He readily reduces the question to the case of  $\mathbb{R}P^{2r} \rightarrow \mathbb{R}P^{2s}$  (using Proposition 6.1 and Brown's embedding theorem as we do) but then dealing with these manifolds calls for some ingenuity. It is not surprising that these manifolds are the difficult ones since these are the even dimensional manifolds for which  $h^2 h(\ )$  is non-zero (by Proposition 3.5).

## References

- [1] **M A Asadi-Golmankhaneh**, *Self-intersection manifolds of immersions*, PhD Thesis, University of Manchester (1998)
- [2] **M A Asadi-Golmankhaneh**, **P J Eccles**, *Determining the characteristic numbers of self-intersection manifolds*, J. London Math. Soc. (to appear)
- [3] **T F Bancho**, *Triple points and surgery of immersed surfaces*, Proc. Amer. Math. Soc. 46 (1974) 407{413
- [4] **M G Barratt**, **P J Eccles**, *On the stable structure of  $\mathbb{R}P^{2r}$* , Topology, 13 (1974) 199{207
- [5] **J M Boardman**, **B Steer**, *On Hopf invariants*, Comment. Math. Helv. 42 (1967) 180{221
- [6] **R L W Brown**, *Immersions and embeddings up to cobordism*, Can. J. Math. 23 (1971) 1102{1115
- [7] **R L Cohen**, *The immersion conjecture for differentiable manifolds*, Annals of Math. 122 (1985) 237{328
- [8] **A Dold**, *Erzeugende der Thomschen Algebra  $\mathfrak{R}$* , Math Z. 65 (1956) 25{35
- [9] **E Dyer**, **R K Lashof**, *Homology of iterated loop spaces*, Amer. J. Math. 84 (1962) 35{88

- [10] **P J Eccles**, *Multiple points of codimension one immersions of oriented manifolds*, Math. Proc. Cambridge Philos. Soc. 87 (1980) 213{220
- [11] **P J Eccles**, *Codimension one immersions and the Kervaire invariant one problem*, Math. Proc. Cambridge Philos. Soc. 90 (1981) 483{493
- [12] **P J Eccles**, *Characteristic numbers of immersions and self-intersection manifolds*, from: "\Proceedings of the Colloquium in Topology, Szekszard, Hungary, August 1993", Bolyai Society Mathematical Studies 4, James Bolyai Mathematical Society, Budapest (1995) 197{216
- [13] **P J Eccles**, *Double point manifolds and immersions of spheres in Euclidean space*, from "\Proceedings of the William Browder Sixtieth Birthday Conference, March 1994", Annals of Math. Studies 138, Princeton University Press (1997) 125{137
- [14] **S O Kochman**, *Bordism, stable homotopy and Adams spectral sequences*, American Mathematical Society (1996)
- [15] **U Koschorke, B Sanderson**, *Self intersections and higher Hopf invariants*, Topology, 17 (1978) 283{290
- [16] **J Lannes**, *Sur les immersions de Boy*, Lecture Notes in Mathematics 1051, Springer (1984) 263{270
- [17] **J P May**, *The homology of  $E_1$  spaces*, Lecture Notes in Mathematics 533, Springer (1976) 1{68
- [18] **J W Milnor, J D Stashe** , *Characteristic classes*, Ann. of Math. Studies 76, Princeton University Press (1974)
- [19] **V P Snaith**, *A stable decomposition of  ${}^n S^n X$* , J. London Math. Soc. 7 (1974) 577{583
- [20] **A Szűcs**, *Gruppy kobordizmov I {pogruzeni I}*, Acta Math. Acad. Sci. Hungar. 27 (1976) 343{358
- [21] **A Szűcs**, *Gruppy kobordizmov I {pogruzeni II}*, Acta Math. Acad. Sci. Hungar. 28 (1976) 93{102
- [22] **A Szűcs**, *Double point surfaces of smooth immersions  $M^n \rightarrow \mathbb{R}^{2n-2}$* , Math. Proc. Cambridge Philos. Soc. 113 (1993) 601{613
- [23] **P Vogel**, *Cobordisme d'immersions*, Ann. Sci. Ecole Norm. Sup. 7 (1974) 317{358
- [24] **R Wells**, *Cobordism groups of immersions*, Topology, 5 (1966) 281{294