



NONSTANDARD PROOFS OF EGGLESTON LIKE THEOREMS

SZYMON ŻEBERSKI

ABSTRACT. We prove theorems of the following form: if $A \subseteq \mathbb{R}^2$ is a “big set”, then there exists a “big set” $P \subseteq \mathbb{R}$ and a perfect set $Q \subseteq \mathbb{R}$ such that $P \times Q \subseteq A$. We discuss cases where “big set” means: set of positive Lebesgue measure, set of full Lebesgue measure, Baire measurable set of second Baire category and comeagre set. In the first case (set of positive measure) we obtain the theorem due to Eggleston. In fact we give a simplified version of the proof given by J. Cichoń in [1]. To prove these theorems we use Shoenfield’s theorem about absoluteness for Σ_2^1 -sentences.

1. INTRODUCTION

We use standard set theoretical notation. By ω we denote the set of natural numbers. The cardinality of a set X we denote by $|X|$.

By $[0, 1]$ we denote the unit interval of the real line. By $\text{Perf}([0, 1])$ we denote the Polish space of all non-empty perfect subsets of the interval $[0, 1]$ with the Hausdorff metric. Similarly, by $\text{Perf}(\mathbb{R})$ we denote the Polish space of all non-empty perfect subsets of the real line. By \mathbf{F}_σ we denote the family of all F_σ subsets of the unit interval. By \mathbf{G}_δ we denote the family of all G_δ subsets of the unit interval.

By λ we denote Lebesgue measure on the real line, by λ^2 — Lebesgue measure on the plane. \mathbb{L} denotes the σ -ideal of null sets ($\mathbb{L} = \{A : \lambda(A) = 0\}$). By \mathbb{L}^+ we denote the family of positive Lebesgue measure sets. \mathbb{K} denotes the σ -ideal of first Baire category sets. \mathbb{K}^+ denotes the family of Baire measurable sets of second Baire category. Let us recall that if \mathcal{J} is a proper σ -ideal then $\mathcal{B} \subseteq \mathcal{J}$ is a base of \mathcal{J} iff $(\forall J \in \mathcal{J})(\exists B \in \mathcal{B}) J \subseteq B$. Furthermore, we consider a cardinal number $\text{cof}(\mathcal{J})$ defined as follows:

$$\text{cof}(\mathcal{J}) = \min\{|\mathcal{B}| : \mathcal{B} \text{ is a base of } \mathcal{J}\}.$$

It is well known that we can extend every countable standard transitive model of ZFC via forcing extension to a model of $\text{ZFC} + (\text{cof}(\mathbb{L}) = \omega_1) + (2^\omega =$

2000 *Mathematics Subject Classification.* 03E15, 28A05.

Key words and phrases. Lebesgue measure, perfect set.

ω_2). We can also extend every countable standard transitive model of ZFC via forcing extension to a model of $\text{ZFC} + (\text{cof}(\mathbb{K}) = \omega_1) + (2^\omega = \omega_2)$.

We can code Borel subsets of the real line by functions from ω^ω in an absolute way (see [5]). Functions which code Borel sets are called Borel codes. We use the following notation: if $f \in \omega^\omega$ is a Borel code, then $\#f \subseteq \mathbb{R}$ is a set coded by f .

By a canonical Polish space, we understand a countable product of spaces $\{0, 1\}^\omega$, ω^ω , $[0, 1]$, $\text{Perf}([0, 1])$ and so on. A sentence φ is a Σ_2^1 -sentence if for some canonical Polish spaces X, Y and some Borel subset $B \subseteq X \times Y$ we have $\varphi = (\exists x \in X)(\forall y \in Y) (x, y) \in B$. Spaces X, Y and the set B are called “parameters” of the sentence φ . We shall use the following classical theorem about absoluteness of Σ_2^1 -sentences (see [6]):

Theorem 1 (Shoenfield). *Suppose that $M \subseteq N$ are standard transitive models of the theory ZF such that $\omega_1^N \subseteq M$. Let φ be a Σ_2^1 -sentence with parameters from the model M . Then*

$$M \models \varphi \iff N \models \varphi.$$

Notice that if the model N is a generic extension of the standard transitive model M then both models M and N have the same ordinal numbers, so the inclusion $\omega_1^N \subseteq M$ trivially holds.

Let us note that if A is a coanalytic subset of a Polish space and $|A| > \omega_1$ then there exists a nonempty perfect subset of A . Indeed, there exists a family $\{B_\alpha\}_{\alpha < \omega_1}$ of Borel sets such that $A = \bigcup_{\alpha < \omega_1} B_\alpha$ (see [3]). Hence, if $|A| > \omega_1$ then there exists $\alpha < \omega_1$ such that $|B_\alpha| > \omega_1$, so B_α has a nonempty perfect subset.

2. MAIN RESULT

We give a new proof of Eggleston’s theorem (see [4]) and then by a simple modification of this proof we get new results about inscribing special rectangles into “big sets” in the sense of measure or category.

2.1. Measure case. In the paper [2] we can find the following result:

Theorem 2 (Cichoń, Kamburelis, Pawlikowski). *There exists a dense subset of the measure algebra $\text{Borel}(\mathbb{R})/\mathbb{L}$ of cardinality $\text{cof}(\mathbb{L})$.*

Using this theorem we can deduce the following fact:

Fact 1. *There exists a family $\mathcal{B} \subseteq \text{Perf}(\mathbb{R}) \cap \mathbb{L}^+$ such that $|\mathcal{B}| = \text{cof}(\mathbb{L})$ and*

$$(\forall A \subseteq \mathbb{R}) (\lambda(A) > 0 \longrightarrow (\exists P \in \mathcal{B}) P \subseteq A).$$

Using this fact, we can obtain a simple proof of the following theorem from [4]:

Theorem 3 (Eggleston). *Let A be a subset of the plane of positive Lebesgue measure. Then we can find perfect subsets of the real line P, Q such that $\lambda(P) > 0$ and $P \times Q \subseteq A$.*

Proof. Without loss of generality we can assume that A is a Borel set. Let V' be a generic extension of the universe V such that

$$V' \models \text{ZFC} + (\text{cof}(\mathbb{L}) = \omega_1) + (2^\omega = \omega_2).$$

For a while we will work in the universe V' . For $y \in \mathbb{R}$ let $A^y = \{x \in \mathbb{R} : (x, y) \in A\}$ and we put

$$Y = \{y \in \mathbb{R} : \lambda(A^y) > 0\}.$$

It follows from Fubini's theorem that $\lambda(Y) > 0$. In particular $|Y| = \omega_2$. Using Fact 1 we get a family $\mathcal{B} \subseteq \text{Perf}(\mathbb{R}) \cap \mathbb{L}^+$ such that

$$(\forall A \subseteq \mathbb{R}) (\lambda(A) > 0 \longrightarrow (\exists P \in \mathcal{B}) P \subseteq A)$$

and $|\mathcal{B}| = \text{cof}(\mathbb{L}) = \omega_1$. For every $y \in Y$ we can find $P \in \mathcal{B}$ such that $P \subseteq A^y$. So there exists $P \in \text{Perf}(\mathbb{R}) \cap \mathbb{L}^+$ such that $|\{y \in Y : P \subseteq A^y\}| = \omega_2$. But $\{y \in Y : P \subseteq A^y\}$ is a coanalytic set of cardinality ω_2 so we can find a perfect set Q such that $Q \subseteq \{y \in Y : P \subseteq A^y\}$. It means that in the universe V' the following sentence holds:

$$(\exists P, Q \in \text{Perf}(\mathbb{R})) (\forall r \in \mathbb{R}^2) \lambda(P) > 0 \wedge (r \in P \times Q \rightarrow r \in A).$$

But this is a Σ_2^1 -sentence, hence it is also true in the original universe V . \square

Now, we show the theorem about inscribing special rectangles into sets of full Lebesgue measure. At first, we have to formulate the following fact:

Fact 2. *There exists a family $\mathcal{F} \subseteq \mathbf{F}_\sigma$ of measure one subsets of the unit interval such that $|\mathcal{F}| = \text{cof}(\mathbb{L})$ and*

$$(\forall A \subseteq [0, 1]) (\lambda(A) = 1 \longrightarrow (\exists F \in \mathcal{F}) F \subseteq A).$$

Using this fact, we prove the following theorem:

Theorem 4. *Let A be a subset of $[0, 1]^2$ such that $\lambda^2(A) = 1$. Then we can find two sets $F, Q \subseteq [0, 1]$ such that F is a F_σ set, $\lambda(F) = 1$, Q is a perfect set and $F \times Q \subseteq A$.*

Proof. Without loss of generality A is a Borel set. Let V' be a generic extension of the universe V such that

$$V' \models \text{ZFC} + (\text{cof}(\mathbb{L}) = \omega_1) + (2^\omega = \omega_2).$$

Similarly as in the proof of theorem 3, using Fact 2, we deduce that in the universe V' the following sentence holds

$$(\exists F \in \mathbf{F}_\sigma)(\exists Q \in \text{Perf}([0, 1])) \lambda(F) = 1 \wedge F \times Q \subseteq A.$$

But we can formulate this sentence in the following way

$$(\exists f \in \omega^\omega)(\exists Q \in \text{Perf}([0, 1])) f \text{ codes } F_\sigma \text{ set} \wedge \lambda(\#f) = 1 \wedge \#f \times Q \subseteq A.$$

This is also a Σ_2^1 -sentence, hence it is true in the original universe V . \square

2.2. Category case. Now, we will formulate and prove theorems about inscribing special rectangles into “big sets” in the sense of category. We start with two observations (similar to Fact 1 and 2). Namely, we have the following facts:

Fact 3. *There exists a family $\mathcal{G} \subseteq \mathbf{G}_\delta \cap \mathbb{K}^+$ such that $|\mathcal{G}| = \text{cof}(\mathbb{K})$ and*

$$(\forall A \in \mathbb{K}^+)(\exists G \in \mathcal{G}) G \subseteq A.$$

Fact 4. *There exists a family $\mathcal{C} \subseteq \mathbf{G}_\delta$ of comeagre subsets of the unit interval such that $|\mathcal{C}| = \text{cof}(\mathbb{K})$ and*

$$(\forall A \subseteq [0, 1]) (A \text{ is comeagre} \longrightarrow (\exists C \in \mathcal{C}) C \subseteq A).$$

Using these facts, we prove the following theorems:

Theorem 5. *Let $A \subseteq [0, 1]^2$ be a Baire measurable set of second Baire category. Then we can find two sets $G, Q \subseteq [0, 1]$ such that G is a G_δ set of second Baire category, Q is a perfect set and $G \times Q \subseteq A$.*

Theorem 6. *Let $A \subseteq [0, 1]^2$ be a comeagre set. Then we can find two sets $C, Q \subseteq [0, 1]$ such that C is a comeagre G_δ set, Q is a perfect set and $C \times Q \subseteq A$.*

Proof of Theorem 5 and Theorem 6. We start with extending the universe V to V' (by generic extension) such that

$$V' \models \text{ZFC} + (\text{cof}(\mathbb{K}) = \omega_1) + (2^\omega = \omega_2).$$

Similarly as in the proof of Theorem 3 (naturally using Facts 3, 4 respectively for Theorem 5 and 6), we deduce that if A is a Baire measurable second Baire category set (Theorem 5) then in the model V' the following sentence holds:

$$(\exists G \in \mathbf{G}_\delta)(\exists Q \in \text{Perf}([0, 1])) G \in \mathbb{K}^+ \wedge G \times Q \subseteq A.$$

When A is comeagre (Theorem 6) then in the universe V' the following sentence holds:

$$(\exists C \in \mathbf{G}_\delta)(\exists Q \in \text{Perf}([0, 1])) C \text{ is comeagre} \wedge C \times Q \subseteq A.$$

Similarly as in the proof of Theorem 4, we can use Borel coding for replacing quantifiers of the type $(\exists A \in \mathbf{G}_\delta)$ by

$$(\exists a \in \omega^\omega) a \text{ codes } G_\delta \text{ set}$$

and replace A by $\#a$. After this modification we obtain equivalent sentences, which are Σ_2^1 -sentences, hence they are true in the original universe V . \square

REFERENCES

1. J. Cichoń, *Applications of Shoenfield absoluteness Theorem*, unpublished.
2. J. Cichoń, A. Kamburelis, and J. Pawlikowski, *On dense subsets of the measure algebra*, Proc. Amer. Math. Soc. **94** (1985), no. 1, 142–146. MR **86j**:04001
3. J. Cichoń, A. Kharazishvili, and B. Węglorz, *Subsets of the Real Line*, Wydawnictwo Uniwersytetu Łódzkiego, Łódź (1995), 63–70.

4. H. G. Eggleston, *Two measure properties of Cartesian product sets*, Quart. J. Math., Oxford Ser. (2) **5** (1954), 108–115. MR 16,344e
5. Thomas Jech, *Set theory*, second ed., Springer-Verlag, Berlin, 1997. MR **99b**:03061
6. J. R. Shoenfield, *The problem of predicativity*, Essays on the foundations of mathematics, Magnes Press, Hebrew Univ., Jerusalem, 1961, pp. 132–139. MR 29 #2177

INSTITUTE OF MATHEMATICS, UNIVERSITY OF WROCLAW, PL. GRUNWALDZKI 2/4,
50-384 WROCLAW, POLAND

E-mail address: `szebe@math.uni.wroc.pl`