

S.S. DRAGOMIR

School of Computer Science and Mathematics
Victoria University
PO Box 14428, MCMC 8001
VIC, Australia.

EMail: sever@csm.vu.edu.au

URL: <http://rgmia.vu.edu.au/dragomir/>



volume 6, issue 5, article 129,
2005.

Received 18 April, 2005;
accepted 05 July, 2005.

Communicated by: B. Mond

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract

Recent reverses for the discrete generalised triangle inequality and its continuous version for vector-valued integrals in Banach spaces are surveyed. New results are also obtained. Particular instances of interest in Hilbert spaces and for complex numbers and functions are pointed out as well.


2000 Mathematics Subject Classification: Primary 46B05, 46C05; Secondary 26D15, 26D10

Key words: Reverse triangle inequality, Hilbert spaces, Banach spaces, Bochner integral.

This paper is based on the talk given by the author within the "International Conference of Mathematical Inequalities and their Applications, I", December 06-08, 2004, Victoria University, Melbourne, Australia [<http://rgmia.vu.edu.au/conference>]

Contents

1	Introduction	4
2	Diaz-Metcalf Type Inequalities	9
3	Inequalities of Diaz-Metcalf Type for m Functionals	12
	3.1 The Case of Normed Spaces	12
	3.2 The Case of Inner Product Spaces	14
4	Diaz-Metcalf Inequality for Semi-Inner Products	24
5	Other Multiplicative Reverses for m Functionals	28
6	An Additive Reverse for the Triangle Inequality	34



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents




Go Back

Close

Quit

Page 2 of 99

6.1	The Case of One Functional	34
6.2	The Case of m Functionals	37
6.3	The Case of Inner Product Spaces	39
7	Other Additive Reverses for m Functionals	46
8	Applications for Complex Numbers	50
9	Karamata Type Inequalities in Hilbert Spaces	58
10	Multiplicative Reverses of the Continuous Triangle Inequality	61
10.1	The Case of One Functional	61
10.2	The Case of m Functionals	65
11	Additive Reverses of the Continuous Triangle Inequality	77
11.1	The Case of One Functional	77
11.2	The Case of m Functionals	81
12	Applications for Complex-Valued Functions	90
References		



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 3 of 99

1. Introduction

The *generalised triangle inequality*, namely

$$\left\| \sum_{i=1}^n x_i \right\| \leq \sum_{i=1}^n \|x_i\|,$$

provided $(X, \|\cdot\|)$ is a normed linear space over the real or complex field $\mathbb{K} = \mathbb{R}, \mathbb{C}$ and $x_i, i \in \{1, \dots, n\}$ are vectors in X plays a fundamental role in establishing various analytic and geometric properties of such spaces.


With no less importance, the *continuous* version of it, i.e.,

$$(1.1) \quad \left\| \int_a^b f(t) dt \right\| \leq \int_a^b \|f(t)\| dt,$$

where $f : [a, b] \subset \mathbb{R} \rightarrow X$ is a strongly measurable function on the compact interval $[a, b]$ with values in the Banach space X and $\|f(\cdot)\|$ is Lebesgue integrable on $[a, b]$, is crucial in the Analysis of vector-valued functions with countless applications in Functional Analysis, Operator Theory, Differential Equations, Semigroups Theory and related fields.

Surprisingly enough, the reverses of these, i.e., inequalities of the following type

$$\sum_{i=1}^n \|x_i\| \leq C \left\| \sum_{i=1}^n x_i \right\|, \quad \int_a^b \|f(t)\| dt \leq C \left\| \int_a^b f(t) dt \right\|,$$



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 4 of 99

with $C \geq 1$, which we call *multiplicative reverses*, or

$$\sum_{i=1}^n \|x_i\| \leq \left\| \sum_{i=1}^n x_i \right\| + M, \quad \int_a^b \|f(t)\| dt \leq \left\| \int_a^b f(t) dt \right\| + M,$$

with $M \geq 0$, which we call *additive reverses*, under suitable assumptions for the involved vectors or functions, are far less known in the literature.

It is worth mentioning though, the following reverse of the generalised triangle inequality for complex numbers


$$\cos \theta \sum_{k=1}^n |z_k| \leq \left| \sum_{k=1}^n z_k \right|,$$

provided the complex numbers $z_k, k \in \{1, \dots, n\}$ satisfy the assumption

$$a - \theta \leq \arg(z_k) \leq a + \theta, \quad \text{for any } k \in \{1, \dots, n\},$$

where $a \in \mathbb{R}$ and $\theta \in (0, \frac{\pi}{2})$ was first discovered by M. Petrovich in 1917, [22] (see [20, p. 492]) and subsequently was rediscovered by other authors, including J. Karamata [14, p. 300 – 301], H.S. Wilf [23], and in an equivalent form by M. Marden [18]. Marden and Wilf have outlined in their work the important fact that reverses of the generalised triangle inequality may be successfully applied to the location problem for the roots of complex polynomials.

In 1966, J.B. Diaz and F.T. Metcalf [2] proved the following reverse of the triangle inequality in the more general case of inner product spaces:



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 5 of 99

Theorem 1.1 (Diaz-Metcalf, 1966). Let a be a unit vector in the inner product space $(H; \langle \cdot, \cdot \rangle)$ over the real or complex number field \mathbb{K} . Suppose that the vectors $x_i \in H \setminus \{0\}$, $i \in \{1, \dots, n\}$ satisfy

$$0 \leq r \leq \frac{\operatorname{Re} \langle x_i, a \rangle}{\|x_i\|}, \quad i \in \{1, \dots, n\}.$$

Then

$$r \sum_{i=1}^n \|x_i\| \leq \left\| \sum_{i=1}^n x_i \right\|,$$

where equality holds if and only if

$$\sum_{i=1}^n x_i = r \left(\sum_{i=1}^n \|x_i\| \right) a.$$


A generalisation of this result for orthonormal families is incorporated in the following result [2].

Theorem 1.2 (Diaz-Metcalf, 1966). Let a_1, \dots, a_n be orthonormal vectors in H . Suppose the vectors $x_1, \dots, x_n \in H \setminus \{0\}$ satisfy

$$0 \leq r_k \leq \frac{\operatorname{Re} \langle x_i, a_k \rangle}{\|x_i\|}, \quad i \in \{1, \dots, n\}, \quad k \in \{1, \dots, m\}.$$

Then

$$\left(\sum_{k=1}^m r_k^2 \right)^{\frac{1}{2}} \sum_{i=1}^n \|x_i\| \leq \left\| \sum_{i=1}^n x_i \right\|,$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 6 of 99

where equality holds if and only if

$$\sum_{i=1}^n x_i = \left(\sum_{i=1}^n \|x_i\| \right) \sum_{k=1}^m r_k a_k.$$

Similar results valid for semi-inner products may be found in [15], [16] and [19].

Now, for the scalar continuous case.

It appears, see [20, p. 492], that the first reverse inequality for (1.1) in the case of complex valued functions was obtained by J. Karamata in his book from 1949, [14]. It can be stated as

$$\cos \theta \int_a^b |f(x)| dx \leq \left| \int_a^b f(x) dx \right|$$


provided

$$-\theta \leq \arg f(x) \leq \theta, \quad x \in [a, b]$$

for given $\theta \in (0, \frac{\pi}{2})$.

This result has recently been extended by the author for the case of Bochner integrable functions with values in a Hilbert space H . If by $L([a, b]; H)$, we denote the space of Bochner integrable functions with values in a Hilbert space H , i.e., we recall that $f \in L([a, b]; H)$ if and only if $f : [a, b] \rightarrow H$ is strongly measurable on $[a, b]$ and the Lebesgue integral $\int_a^b \|f(t)\| dt$ is finite, then

$$(1.2) \quad \int_a^b \|f(t)\| dt \leq K \left\| \int_a^b f(t) dt \right\|,$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 7 of 99


provided that f satisfies the condition

$$\|f(t)\| \leq K \operatorname{Re} \langle f(t), e \rangle \quad \text{for a.e. } t \in [a, b],$$

where $e \in H$, $\|e\| = 1$ and $K \geq 1$ are given. The case of equality holds in (1.2) if and only if

$$\int_a^b f(t) dt = \frac{1}{K} \left(\int_a^b \|f(t)\| dt \right) e.$$

The aim of the present paper is to survey some of the recent results concerning multiplicative and additive reverses for both the discrete and continuous version of the triangle inequalities in Banach spaces. New results and applications for the important case of Hilbert spaces and for complex numbers and complex functions have been provided as well.



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 8 of 99

2. Diaz-Metcalf Type Inequalities

In [2], Diaz and Metcalf established the following reverse of the generalised triangle inequality in real or complex normed linear spaces.

Theorem 2.1 (Diaz-Metcalf, 1966). *If $F : X \rightarrow \mathbb{K}$, $\mathbb{K} = \mathbb{R}, \mathbb{C}$ is a linear functional of a unit norm defined on the normed linear space X endowed with the norm $\|\cdot\|$ and the vectors x_1, \dots, x_n satisfy the condition*

$$(2.1) \quad 0 \leq r \leq \operatorname{Re} F(x_i), \quad i \in \{1, \dots, n\};$$

then

$$(2.2) \quad r \sum_{i=1}^n \|x_i\| \leq \left\| \sum_{i=1}^n x_i \right\|,$$

where equality holds if and only if both


$$(2.3) \quad F\left(\sum_{i=1}^n x_i\right) = r \sum_{i=1}^n \|x_i\|$$

and

$$(2.4) \quad F\left(\sum_{i=1}^n x_i\right) = \left\| \sum_{i=1}^n x_i \right\|.$$

If $X = H$, $(H; \langle \cdot, \cdot \rangle)$ is an inner product space and $F(x) = \langle x, e \rangle$, $\|e\| = 1$, then the condition (2.1) may be replaced with the simpler assumption

$$(2.5) \quad 0 \leq r \|x_i\| \leq \operatorname{Re} \langle x_i, e \rangle, \quad i = 1, \dots, n,$$



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 9 of 99

which implies the reverse of the generalised triangle inequality (2.2). In this case the equality holds in (2.2) if and only if [2]

$$(2.6) \quad \sum_{i=1}^n x_i = r \left(\sum_{i=1}^n \|x_i\| \right) e.$$

Theorem 2.2 (Diaz-Metcalf, 1966). *Let F_1, \dots, F_m be linear functionals on X , each of unit norm. As in [2], let consider the real number c defined by*

$$c = \sup_{x \neq 0} \left[\frac{\sum_{k=1}^m |F_k(x)|^2}{\|x\|^2} \right];$$

it then follows that $1 \leq c \leq m$. Suppose the vectors x_1, \dots, x_n whenever $x_i \neq 0$, satisfy


$$(2.7) \quad 0 \leq r_k \|x_i\| \leq \operatorname{Re} F_k(x_i), \quad i = 1, \dots, n, \quad k = 1, \dots, m.$$

Then one has the following reverse of the generalised triangle inequality [2]

$$(2.8) \quad \left(\frac{\sum_{k=1}^m r_k^2}{c} \right)^{\frac{1}{2}} \sum_{i=1}^n \|x_i\| \leq \left\| \sum_{i=1}^n x_i \right\|,$$

where equality holds if and only if both

$$(2.9) \quad F_k \left(\sum_{i=1}^n x_i \right) = r_k \sum_{i=1}^n \|x_i\|, \quad k = 1, \dots, m$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 10 of 99

and

$$(2.10) \quad \sum_{k=1}^m \left[F_k \left(\sum_{i=1}^n x_i \right) \right]^2 = c \left\| \sum_{i=1}^n x_i \right\|^2.$$

If $X = H$, an inner product space, then, for $F_k(x) = \langle x, e_k \rangle$, where $\{e_k\}_{k=1, \overline{n}}$ is an orthonormal family in H , i.e., $\langle e_i, e_j \rangle = \delta_{ij}$, $i, j \in \{1, \dots, k\}$, δ_{ij} is Kronecker delta, the condition (2.7) may be replaced by

$$(2.11) \quad 0 \leq r_k \|x_i\| \leq \operatorname{Re} \langle x_i, e_k \rangle, \quad i = 1, \dots, n, \quad k = 1, \dots, m;$$

implying the following reverse of the generalised triangle inequality


$$(2.12) \quad \left(\sum_{k=1}^m r_k^2 \right)^{\frac{1}{2}} \sum_{i=1}^n \|x_i\| \leq \left\| \sum_{i=1}^n x_i \right\|,$$

where the equality holds if and only if

$$(2.13) \quad \sum_{i=1}^n x_i = \left(\sum_{i=1}^n \|x_i\| \right) \sum_{k=1}^m r_k e_k.$$

The aim of the following sections is to present recent reverses of the triangle inequality obtained by the author in [5] and [6]. New results are established for the general case of normed spaces. Their versions in inner product spaces are analyzed and applications for complex numbers are given as well.

For various classical inequalities related to the triangle inequality, see Chapter XVII of the book [20] and the references therein.



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 11 of 99

3. Inequalities of Diaz-Metcalf Type for m Functionals

3.1. The Case of Normed Spaces

The following result may be stated [5].

Theorem 3.1 (Dragomir, 2004). *Let $(X, \|\cdot\|)$ be a normed linear space over the real or complex number field \mathbb{K} and $F_k : X \rightarrow \mathbb{K}$, $k \in \{1, \dots, m\}$ continuous linear functionals on X . If $x_i \in X \setminus \{0\}$, $i \in \{1, \dots, n\}$ are such that there exists the constants $r_k \geq 0$, $k \in \{1, \dots, m\}$ with $\sum_{k=1}^m r_k > 0$ and*

$$(3.1) \quad \operatorname{Re} F_k(x_i) \geq r_k \|x_i\|$$

for each $i \in \{1, \dots, n\}$ and $k \in \{1, \dots, m\}$, then


$$(3.2) \quad \sum_{i=1}^n \|x_i\| \leq \frac{\|\sum_{k=1}^m F_k\|}{\sum_{k=1}^m r_k} \left\| \sum_{i=1}^n x_i \right\|.$$

The case of equality holds in (3.2) if both

$$(3.3) \quad \left(\sum_{k=1}^m F_k \right) \left(\sum_{i=1}^n x_i \right) = \left(\sum_{k=1}^m r_k \right) \sum_{i=1}^n \|x_i\|$$

and

$$(3.4) \quad \left(\sum_{k=1}^m F_k \right) \left(\sum_{i=1}^n x_i \right) = \left\| \sum_{k=1}^m F_k \right\| \left\| \sum_{i=1}^n x_i \right\|.$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 12 of 99

Proof. Utilising the hypothesis (3.1) and the properties of the modulus, we have

$$\begin{aligned}
 (3.5) \quad I &:= \left| \left(\sum_{k=1}^m F_k \right) \left(\sum_{i=1}^n x_i \right) \right| \geq \left| \operatorname{Re} \left[\left(\sum_{k=1}^m F_k \right) \left(\sum_{i=1}^n x_i \right) \right] \right| \\
 &\geq \sum_{k=1}^m \operatorname{Re} F_k \left(\sum_{i=1}^n x_i \right) = \sum_{k=1}^m \sum_{i=1}^n \operatorname{Re} F_k(x_i) \\
 &\geq \left(\sum_{k=1}^m r_k \right) \sum_{i=1}^n \|x_i\|.
 \end{aligned}$$

On the other hand, by the continuity property of F_k , $k \in \{1, \dots, m\}$ we obviously have

$$(3.6) \quad I = \left| \left(\sum_{k=1}^m F_k \right) \left(\sum_{i=1}^n x_i \right) \right| \leq \left\| \sum_{k=1}^m F_k \right\| \left\| \sum_{i=1}^n x_i \right\|.$$

Making use of (3.5) and (3.6), we deduce the desired inequality (3.2).

Now, if (3.3) and (3.4) are valid, then, obviously, the case of equality holds true in the inequality (3.2).

Conversely, if the case of equality holds in (3.2), then it must hold in all the inequalities used to prove (3.2). Therefore we have

$$(3.7) \quad \operatorname{Re} F_k(x_i) = r_k \|x_i\|$$

for each $i \in \{1, \dots, n\}$, $k \in \{1, \dots, m\}$;

$$(3.8) \quad \sum_{k=1}^m \operatorname{Im} F_k \left(\sum_{i=1}^n x_i \right) = 0$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 13 of 99

and

$$(3.9) \quad \sum_{k=1}^m \operatorname{Re} F_k \left(\sum_{i=1}^n x_i \right) = \left\| \sum_{k=1}^m F_k \right\| \left\| \sum_{i=1}^n x_i \right\|.$$

Note that, from (3.7), by summation over i and k , we get

$$(3.10) \quad \operatorname{Re} \left[\left(\sum_{k=1}^m F_k \right) \left(\sum_{i=1}^n x_i \right) \right] = \left(\sum_{k=1}^m r_k \right) \sum_{i=1}^n \|x_i\|.$$

Since (3.8) and (3.10) imply (3.3), while (3.9) and (3.10) imply (3.4) hence the theorem is proved. \square

Remark 1. *If the norms $\|F_k\|$, $k \in \{1, \dots, m\}$ are easier to find, then, from (3.2), one may get the (coarser) inequality that might be more useful in practice:*


$$(3.11) \quad \sum_{i=1}^n \|x_i\| \leq \frac{\sum_{k=1}^m \|F_k\|}{\sum_{k=1}^m r_k} \left\| \sum_{i=1}^n x_i \right\|.$$

3.2. The Case of Inner Product Spaces

The case of inner product spaces, in which we may provide a simpler condition for equality, is of interest in applications [5].

Theorem 3.2 (Dragomir, 2004). *Let $(H; \langle \cdot, \cdot \rangle)$ be an inner product space over the real or complex number field \mathbb{K} , $e_k, x_i \in H \setminus \{0\}$, $k \in \{1, \dots, m\}$, $i \in \{1, \dots, n\}$. If $r_k \geq 0$, $k \in \{1, \dots, m\}$ with $\sum_{k=1}^m r_k > 0$ satisfy*

$$(3.12) \quad \operatorname{Re} \langle x_i, e_k \rangle \geq r_k \|x_i\|$$



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents




Go Back

Close

Quit

Page 14 of 99



Title Page

Contents



Go Back

Close

Quit

Page 15 of 99

for each $i \in \{1, \dots, n\}$ and $k \in \{1, \dots, m\}$, then

$$(3.13) \quad \sum_{i=1}^n \|x_i\| \leq \frac{\|\sum_{k=1}^m e_k\|}{\sum_{k=1}^m r_k} \left\| \sum_{i=1}^n x_i \right\|.$$

The case of equality holds in (3.13) if and only if

$$(3.14) \quad \sum_{i=1}^n x_i = \frac{\sum_{k=1}^m r_k}{\|\sum_{k=1}^m e_k\|^2} \left(\sum_{i=1}^n \|x_i\| \right) \sum_{k=1}^m e_k.$$

Proof. By the properties of inner product and by (3.12), we have

$$(3.15) \quad \left| \left\langle \sum_{i=1}^n x_i, \sum_{k=1}^m e_k \right\rangle \right| \geq \left| \sum_{k=1}^m \operatorname{Re} \left\langle \sum_{i=1}^n x_i, e_k \right\rangle \right| \\
\geq \sum_{k=1}^m \operatorname{Re} \left\langle \sum_{i=1}^n x_i, e_k \right\rangle \\
= \sum_{k=1}^m \sum_{i=1}^n \operatorname{Re} \langle x_i, e_k \rangle \geq \left(\sum_{k=1}^m r_k \right) \sum_{i=1}^n \|x_i\| > 0.$$

Observe also that, by (3.15), $\sum_{k=1}^m e_k \neq 0$.

On utilising Schwarz's inequality in the inner product space $(H; \langle \cdot, \cdot \rangle)$ for $\sum_{i=1}^n x_i$, $\sum_{k=1}^m e_k$, we have

$$(3.16) \quad \left\| \sum_{i=1}^n x_i \right\| \left\| \sum_{k=1}^m e_k \right\| \geq \left| \left\langle \sum_{i=1}^n x_i, \sum_{k=1}^m e_k \right\rangle \right|.$$

Making use of (3.15) and (3.16), we can conclude that (3.13) holds.

Now, if (3.14) holds true, then, by taking the norm, we have

$$\begin{aligned} \left\| \sum_{i=1}^n x_i \right\| &= \frac{(\sum_{k=1}^m r_k) \sum_{i=1}^n \|x_i\|}{\left\| \sum_{k=1}^m e_k \right\|^2} \left\| \sum_{k=1}^m e_k \right\| \\ &= \frac{(\sum_{k=1}^m r_k)}{\left\| \sum_{k=1}^m e_k \right\|} \sum_{i=1}^n \|x_i\|, \end{aligned}$$

i.e., the case of equality holds in (3.13).

Conversely, if the case of equality holds in (3.13), then it must hold in all the inequalities used to prove (3.13). Therefore, we have

$$(3.17) \quad \operatorname{Re} \langle x_i, e_k \rangle = r_k \|x_i\|$$

for each $i \in \{1, \dots, n\}$ and $k \in \{1, \dots, m\}$,


$$(3.18) \quad \left\| \sum_{i=1}^n x_i \right\| \left\| \sum_{k=1}^m e_k \right\| = \left\langle \sum_{i=1}^n x_i, \sum_{k=1}^m e_k \right\rangle$$

and

$$(3.19) \quad \operatorname{Im} \left\langle \sum_{i=1}^n x_i, \sum_{k=1}^m e_k \right\rangle = 0.$$

From (3.17), on summing over i and k , we get

$$(3.20) \quad \operatorname{Re} \left\langle \sum_{i=1}^n x_i, \sum_{k=1}^m e_k \right\rangle = \left(\sum_{k=1}^m r_k \right) \sum_{i=1}^n \|x_i\|.$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 16 of 99

By (3.19) and (3.20), we have

$$(3.21) \quad \left\langle \sum_{i=1}^n x_i, \sum_{k=1}^m e_k \right\rangle = \left(\sum_{k=1}^m r_k \right) \sum_{i=1}^n \|x_i\|.$$

On the other hand, by the use of the following identity in inner product spaces

$$(3.22) \quad \left\| u - \frac{\langle u, v \rangle}{\|v\|^2} v \right\|^2 = \frac{\|u\|^2 \|v\|^2 - |\langle u, v \rangle|^2}{\|v\|^2}, \quad v \neq 0,$$

the relation (3.18) holds if and only if

$$(3.23) \quad \sum_{i=1}^n x_i = \frac{\langle \sum_{i=1}^n x_i, \sum_{k=1}^m e_k \rangle}{\|\sum_{k=1}^m e_k\|^2} \sum_{k=1}^m e_k.$$

Finally, on utilising (3.21) and (3.23), we deduce that the condition (3.14) is necessary for the equality case in (3.13). \square


Before we give a corollary of the above theorem, we need to state the following lemma that has been basically obtained in [4]. For the sake of completeness, we provide a short proof here as well.

Lemma 3.3 (Dragomir, 2004). *Let $(H; \langle \cdot, \cdot \rangle)$ be an inner product space over the real or complex number field \mathbb{K} and $x, a \in H, r > 0$ such that:*

$$(3.24) \quad \|x - a\| \leq r < \|a\|.$$

Then we have the inequality

$$(3.25) \quad \|x\| \left(\|a\|^2 - r^2 \right)^{\frac{1}{2}} \leq \operatorname{Re} \langle x, a \rangle$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 17 of 99

or, equivalently

$$(3.26) \quad \|x\|^2 \|a\|^2 - [\operatorname{Re} \langle x, a \rangle]^2 \leq r^2 \|x\|^2.$$

The case of equality holds in (3.25) (or in (3.26)) if and only if

$$(3.27) \quad \|x - a\| = r \quad \text{and} \quad \|x\|^2 + r^2 = \|a\|^2.$$

Proof. From the first part of (3.24), we have

$$(3.28) \quad \|x\|^2 + \|a\|^2 - r^2 \leq 2 \operatorname{Re} \langle x, a \rangle.$$

By the second part of (3.24) we have $(\|a\|^2 - r^2)^{\frac{1}{2}} > 0$, therefore, by (3.28), we may state that


$$(3.29) \quad 0 < \frac{\|x\|^2}{(\|a\|^2 - r^2)^{\frac{1}{2}}} + (\|a\|^2 - r^2)^{\frac{1}{2}} \leq \frac{2 \operatorname{Re} \langle x, a \rangle}{(\|a\|^2 - r^2)^{\frac{1}{2}}}.$$

Utilising the elementary inequality

$$\frac{1}{\alpha}q + \alpha p \geq 2\sqrt{pq}, \quad \alpha > 0, p > 0, q \geq 0;$$

with equality if and only if $\alpha = \sqrt{\frac{q}{p}}$, we may state (for $\alpha = (\|a\|^2 - r^2)^{\frac{1}{2}}$, $p = 1$, $q = \|x\|^2$) that

$$(3.30) \quad 2\|x\| \leq \frac{\|x\|^2}{(\|a\|^2 - r^2)^{\frac{1}{2}}} + (\|a\|^2 - r^2)^{\frac{1}{2}}.$$



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents




Go Back

Close

Quit

Page 18 of 99



Title Page

Contents



Go Back

Close

Quit

Page 19 of 99

The inequality (3.25) follows now by (3.29) and (3.30).

From the above argument, it is clear that the equality holds in (3.25) if and only if it holds in (3.29) and (3.30). However, the equality holds in (3.29) if and only if $\|x - a\| = r$ and in (3.30) if and only if $(\|a\|^2 - r^2)^{\frac{1}{2}} = \|x\|$.

The proof is thus completed. □

We may now state the following corollary [5].

Corollary 3.4. *Let $(H; \langle \cdot, \cdot \rangle)$ be an inner product space over the real or complex number field \mathbb{K} , $e_k, x_i \in H \setminus \{0\}$, $k \in \{1, \dots, m\}$, $i \in \{1, \dots, n\}$. If $\rho_k \geq 0$, $k \in \{1, \dots, m\}$ with*

$$(3.31) \quad \|x_i - e_k\| \leq \rho_k < \|e_k\|$$

for each $i \in \{1, \dots, n\}$ and $k \in \{1, \dots, m\}$, then

$$(3.32) \quad \sum_{i=1}^n \|x_i\| \leq \frac{\|\sum_{k=1}^m e_k\|}{\sum_{k=1}^m (\|e_k\|^2 - \rho_k^2)^{\frac{1}{2}}} \left\| \sum_{i=1}^n x_i \right\|.$$

The case of equality holds in (3.32) if and only if

$$\sum_{i=1}^n x_i = \frac{\sum_{k=1}^m (\|e_k\|^2 - \rho_k^2)^{\frac{1}{2}}}{\|\sum_{k=1}^m e_k\|^2} \left(\sum_{i=1}^n \|x_i\| \right) \sum_{k=1}^m e_k.$$

Proof. Utilising Lemma 3.3, we have from (3.31) that

$$\|x_i\| (\|e_k\|^2 - \rho_k^2)^{\frac{1}{2}} \leq \operatorname{Re} \langle x_i, e_k \rangle$$

for each $k \in \{1, \dots, m\}$ and $i \in \{1, \dots, n\}$.

Applying Theorem 3.2 for

$$r_k := (\|e_k\|^2 - \rho_k^2)^{\frac{1}{2}}, \quad k \in \{1, \dots, m\},$$

we deduce the desired result. □

Remark 2. If $\{e_k\}_{k \in \{1, \dots, m\}}$ are orthogonal, then (3.32) becomes

$$(3.33) \quad \sum_{i=1}^n \|x_i\| \leq \frac{(\sum_{k=1}^m \|e_k\|^2)^{\frac{1}{2}}}{\sum_{k=1}^m (\|e_k\|^2 - \rho_k^2)^{\frac{1}{2}}} \left\| \sum_{i=1}^n x_i \right\|$$

with equality if and only if


$$\sum_{i=1}^n x_i = \frac{\sum_{k=1}^m (\|e_k\|^2 - \rho_k^2)^{\frac{1}{2}}}{\sum_{k=1}^m \|e_k\|^2} \left(\sum_{i=1}^n \|x_i\| \right) \sum_{k=1}^m e_k.$$

Moreover, if $\{e_k\}_{k \in \{1, \dots, m\}}$ is assumed to be orthonormal and

$$\|x_i - e_k\| \leq \rho_k \text{ for } k \in \{1, \dots, m\}, i \in \{1, \dots, n\}$$

where $\rho_k \in [0, 1)$ for $k \in \{1, \dots, m\}$, then

$$(3.34) \quad \sum_{i=1}^n \|x_i\| \leq \frac{\sqrt{m}}{\sum_{k=1}^m (1 - \rho_k^2)^{\frac{1}{2}}} \left\| \sum_{i=1}^n x_i \right\|$$



**Reverses of the Triangle
Inequality in Banach Spaces**

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 20 of 99

with equality if and only if

$$\sum_{i=1}^n x_i = \frac{\sum_{k=1}^m (1 - \rho_k^2)^{\frac{1}{2}}}{m} \left(\sum_{i=1}^n \|x_i\| \right) \sum_{k=1}^m e_k.$$

The following lemma may be stated as well [3].

Lemma 3.5 (Dragomir, 2004). *Let $(H; \langle \cdot, \cdot \rangle)$ be an inner product space over the real or complex number field \mathbb{K} , $x, y \in H$ and $M \geq m > 0$. If*

$$(3.35) \quad \operatorname{Re} \langle My - x, x - my \rangle \geq 0$$

or, equivalently,

$$(3.36) \quad \left\| x - \frac{m + M}{2} y \right\| \leq \frac{1}{2} (M - m) \|y\|,$$

then


$$(3.37) \quad \|x\| \|y\| \leq \frac{1}{2} \cdot \frac{M + m}{\sqrt{mM}} \operatorname{Re} \langle x, y \rangle.$$

The equality holds in (3.37) if and only if the case of equality holds in (3.35) and

$$(3.38) \quad \|x\| = \sqrt{mM} \|y\|.$$

Proof. Obviously,

$$\operatorname{Re} \langle My - x, x - my \rangle = (M + m) \operatorname{Re} \langle x, y \rangle - \|x\|^2 - mM \|y\|^2.$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 21 of 99

Then (3.35) is clearly equivalent to

$$(3.39) \quad \frac{\|x\|^2}{\sqrt{mM}} + \sqrt{mM} \|y\|^2 \leq \frac{M+m}{\sqrt{mM}} \operatorname{Re} \langle x, y \rangle.$$

Since, obviously,

$$(3.40) \quad 2 \|x\| \|y\| \leq \frac{\|x\|^2}{\sqrt{mM}} + \sqrt{mM} \|y\|^2,$$

with equality iff $\|x\| = \sqrt{mM} \|y\|$, hence (3.39) and (3.40) imply (3.37).

The case of equality is obvious and we omit the details. \square

Finally, we may state the following corollary of Theorem 3.2, see [5].

Corollary 3.6. *Let $(H; \langle \cdot, \cdot \rangle)$ be an inner product space over the real or complex number field \mathbb{K} , $e_k, x_i \in H \setminus \{0\}$, $k \in \{1, \dots, m\}$, $i \in \{1, \dots, n\}$. If $M_k > \mu_k > 0$, $k \in \{1, \dots, m\}$ are such that either*


$$(3.41) \quad \operatorname{Re} \langle M_k e_k - x_i, x_i - \mu_k e_k \rangle \geq 0$$

or, equivalently,

$$\left\| x_i - \frac{M_k + \mu_k}{2} e_k \right\| \leq \frac{1}{2} (M_k - \mu_k) \|e_k\|$$

for each $k \in \{1, \dots, m\}$ and $i \in \{1, \dots, n\}$, then

$$(3.42) \quad \sum_{i=1}^n \|x_i\| \leq \frac{\left\| \sum_{k=1}^m e_k \right\|}{\sum_{k=1}^m \frac{2 \cdot \sqrt{\mu_k M_k}}{\mu_k + M_k}} \left\| \sum_{i=1}^n x_i \right\|.$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 22 of 99

The case of equality holds in (3.42) if and only if

$$\sum_{i=1}^n x_i = \frac{\sum_{k=1}^m \frac{2 \cdot \sqrt{\mu_k M_k}}{\mu_k + M_k} \|e_k\|}{\left\| \sum_{k=1}^m e_k \right\|^2} \sum_{i=1}^n \|x_i\| \sum_{k=1}^m e_k.$$

Proof. Utilising Lemma 3.5, by (3.41) we deduce


$$\frac{2 \cdot \sqrt{\mu_k M_k}}{\mu_k + M_k} \|x_i\| \|e_k\| \leq \operatorname{Re} \langle x_i, e_k \rangle$$

for each $k \in \{1, \dots, m\}$ and $i \in \{1, \dots, n\}$.

Applying Theorem 3.2 for

$$r_k := \frac{2 \cdot \sqrt{\mu_k M_k}}{\mu_k + M_k} \|e_k\|, \quad k \in \{1, \dots, m\},$$

we deduce the desired result. □



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 23 of 99

4. Diaz-Metcalf Inequality for Semi-Inner Products

In 1961, G. Lumer [17] introduced the following concept.


Definition 4.1. Let X be a linear space over the real or complex number field \mathbb{K} . The mapping $[\cdot, \cdot] : X \times X \rightarrow \mathbb{K}$ is called a semi-inner product on X , if the following properties are satisfied (see also [3, p. 17]):

- (i) $[x + y, z] = [x, z] + [y, z]$ for all $x, y, z \in X$;
- (ii) $[\lambda x, y] = \lambda [x, y]$ for all $x, y \in X$ and $\lambda \in \mathbb{K}$;
- (iii) $[x, x] \geq 0$ for all $x \in X$ and $[x, x] = 0$ implies $x = 0$;
- (iv) $|[x, y]|^2 \leq [x, x][y, y]$ for all $x, y \in X$;
- (v) $[x, \lambda y] = \bar{\lambda} [x, y]$ for all $x, y \in X$ and $\lambda \in \mathbb{K}$.

It is well known that the mapping $X \ni x \mapsto [x, x]^{\frac{1}{2}} \in \mathbb{R}$ is a norm on X and for any $y \in X$, the functional $X \ni x \mapsto [x, y] \in \mathbb{K}$ is a continuous linear functional on X endowed with the norm $\|\cdot\|$ generated by $[\cdot, \cdot]$. Moreover, one has $\|\varphi_y\| = \|y\|$ (see for instance [3, p. 17]).

Let $(X, \|\cdot\|)$ be a real or complex normed space. If $J : X \rightarrow {}_2X^*$ is the normalised duality mapping defined on X , i.e., we recall that (see for instance [3, p. 1])

$$J(x) = \{\varphi \in X^* | \varphi(x) = \|\varphi\| \|x\|, \|\varphi\| = \|x\|\}, \quad x \in X,$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 24 of 99

then we may state the following representation result (see for instance [3, p. 18]):

Each semi-inner product $[\cdot, \cdot] : X \times X \rightarrow K$ that generates the norm $\|\cdot\|$ of the normed linear space $(X, \|\cdot\|)$ over the real or complex number field \mathbb{K} , is of the form

$$[x, y] = \left\langle \tilde{J}(y), x \right\rangle \text{ for any } x, y \in X,$$

where \tilde{J} is a selection of the normalised duality mapping and $\langle \varphi, x \rangle := \varphi(x)$ for $\varphi \in X^*$ and $x \in X$.

Utilising the concept of semi-inner products, we can state the following particular case of the Diaz-Metcalf inequality.

Corollary 4.1. Let $(X, \|\cdot\|)$ be a normed linear space, $[\cdot, \cdot] : X \times X \rightarrow \mathbb{K}$ a semi-inner product generating the norm $\|\cdot\|$ and $e \in X$, $\|e\| = 1$. If $x_i \in X$, $i \in \{1, \dots, n\}$ and $r \geq 0$ such that


$$(4.1) \quad r \|x_i\| \leq \operatorname{Re} [x_i, e] \text{ for each } i \in \{1, \dots, n\},$$

then we have the inequality

$$(4.2) \quad r \sum_{i=1}^n \|x_i\| \leq \left\| \sum_{i=1}^n x_i \right\|.$$

The case of equality holds in (4.2) if and only if both

$$(4.3) \quad \left[\sum_{i=1}^n x_i, e \right] = r \sum_{i=1}^n \|x_i\|$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 25 of 99

and

$$(4.4) \quad \left[\sum_{i=1}^n x_i, e \right] = \left\| \sum_{i=1}^n x_i \right\|.$$

The proof is obvious from the Diaz-Metcalf theorem [2, Theorem 3] applied for the continuous linear functional $F_e(x) = [x, e]$, $x \in X$.

Before we provide a simpler necessary and sufficient condition of equality in (4.2), we need to recall the concept of strictly convex normed spaces and a classical characterisation of these spaces.


Definition 4.2. A normed linear space $(X, \|\cdot\|)$ is said to be strictly convex if for every x, y from X with $x \neq y$ and $\|x\| = \|y\| = 1$, we have $\|\lambda x + (1 - \lambda)y\| < 1$ for all $\lambda \in (0, 1)$.

The following characterisation of strictly convex spaces is useful in what follows (see [1], [13], or [3, p. 21]).

Theorem 4.2. Let $(X, \|\cdot\|)$ be a normed linear space over \mathbb{K} and $[\cdot, \cdot]$ a semi-inner product generating its norm. The following statements are equivalent:

- (i) $(X, \|\cdot\|)$ is strictly convex;
- (ii) For every $x, y \in X$, $x, y \neq 0$ with $[x, y] = \|x\| \|y\|$, there exists a $\lambda > 0$ such that $x = \lambda y$.

The following result may be stated.



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 26 of 99

Corollary 4.3. Let $(X, \|\cdot\|)$ be a strictly convex normed linear space, $[\cdot, \cdot]$ a semi-inner product generating the norm and e, x_i ($i \in \{1, \dots, n\}$) as in Corollary 4.1. Then the case of equality holds in (4.2) if and only if

$$(4.5) \quad \sum_{i=1}^n x_i = r \left(\sum_{i=1}^n \|x_i\| \right) e.$$

Proof. If (4.5) holds true, then, obviously

$$\left\| \sum_{i=1}^n x_i \right\| = r \left(\sum_{i=1}^n \|x_i\| \right) \|e\| = r \sum_{i=1}^n \|x_i\|,$$

which is the equality case in (4.2).

Conversely, if the equality holds in (4.2), then by Corollary 4.1, we have that (4.3) and (4.4) hold true. Utilising Theorem 4.2, we conclude that there exists a $\mu > 0$ such that

$$(4.6) \quad \sum_{i=1}^n x_i = \mu e.$$


Inserting this in (4.3) we get

$$\mu \|e\|^2 = r \sum_{i=1}^n \|x_i\|$$

giving

$$(4.7) \quad \mu = r \sum_{i=1}^n \|x_i\|.$$

Finally, by (4.6) and (4.7) we deduce (4.5) and the corollary is proved. \square



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 27 of 99

5. Other Multiplicative Reverses for m Functionals

Assume that $F_k, k \in \{1, \dots, m\}$ are bounded linear functionals defined on the normed linear space X .

For $p \in [1, \infty)$, define

$$(c_p) \quad c_p := \sup_{x \neq 0} \left[\frac{\sum_{k=1}^m |F_k(x)|^p}{\|x\|^p} \right]^{\frac{1}{p}}$$

and for $p = \infty$,

$$(c_\infty) \quad c_\infty := \sup_{x \neq 0} \left[\max_{1 \leq k \leq m} \left\{ \frac{|F_k(x)|}{\|x\|} \right\} \right].$$


Then, by the fact that $|F_k(x)| \leq \|F_k\| \|x\|$ for any $x \in X$, where $\|F_k\|$ is the norm of the functional F_k , we have that

$$c_p \leq \left(\sum_{k=1}^m \|F_k\|^p \right)^{\frac{1}{p}}, \quad p \geq 1$$

and

$$c_\infty \leq \max_{1 \leq k \leq m} \|F_k\|.$$

We may now state and prove a new reverse inequality for the generalised triangle inequality in normed linear spaces.



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 28 of 99

Theorem 5.1. Let $x_i, r_k, F_k, k \in \{1, \dots, m\}, i \in \{1, \dots, n\}$ be as in the hypothesis of Theorem 3.1. Then we have the inequalities

$$(5.1) \quad (1 \leq) \frac{\sum_{i=1}^n \|x_i\|}{\left\| \sum_{i=1}^n x_i \right\|} \leq \frac{c_\infty}{\max_{1 \leq k \leq m} \{r_k\}} \left(\leq \frac{\max_{1 \leq k \leq m} \|F_k\|}{\max_{1 \leq k \leq m} \{r_k\}} \right).$$

The case of equality holds in (5.1) if and only if

$$(5.2) \quad \operatorname{Re} \left[F_k \left(\sum_{i=1}^n x_i \right) \right] = r_k \sum_{i=1}^n \|x_i\| \quad \text{for each } k \in \{1, \dots, m\}$$

and


$$(5.3) \quad \max_{1 \leq k \leq m} \operatorname{Re} \left[F_k \left(\sum_{i=1}^n x_i \right) \right] = c_\infty \left\| \sum_{i=1}^n x_i \right\|.$$

Proof. Since, by the definition of c_∞ , we have

$$c_\infty \|x\| \geq \max_{1 \leq k \leq m} |F_k(x)|, \quad \text{for any } x \in X,$$

then we can state, for $x = \sum_{i=1}^n x_i$, that

$$(5.4) \quad c_\infty \left\| \sum_{i=1}^n x_i \right\| \geq \max_{1 \leq k \leq m} \left| F_k \left(\sum_{i=1}^n x_i \right) \right| \geq \max_{1 \leq k \leq m} \left[\operatorname{Re} F_k \left(\sum_{i=1}^n x_i \right) \right] \\ \geq \max_{1 \leq k \leq m} \left[\operatorname{Re} \sum_{i=1}^n F_k(x_i) \right] = \max_{1 \leq k \leq m} \left[\sum_{i=1}^n \operatorname{Re} F_k(x_i) \right].$$



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 29 of 99

Utilising the hypothesis (3.1) we obviously have

$$\max_{1 \leq k \leq m} \left[\sum_{i=1}^n \operatorname{Re} F_k(x_i) \right] \geq \max_{1 \leq k \leq m} \{r_k\} \cdot \sum_{i=1}^n \|x_i\|.$$

Also, $\sum_{i=1}^n x_i \neq 0$, because, by the initial assumptions, not all r_k and x_i with $k \in \{1, \dots, m\}$ and $i \in \{1, \dots, n\}$ are allowed to be zero. Hence the desired inequality (5.1) is obtained.

Now, if (5.2) is valid, then, taking the maximum over $k \in \{1, \dots, m\}$ in this equality we get

$$\max_{1 \leq k \leq m} \operatorname{Re} \left[F_k \left(\sum_{i=1}^n x_i \right) \right] = \max_{1 \leq k \leq m} \{r_k\} \left\| \sum_{i=1}^n x_i \right\|,$$

which, together with (5.3) provides the equality case in (5.1).

Now, if the equality holds in (5.1), it must hold in all the inequalities used to prove (5.1), therefore, we have


$$(5.5) \quad \operatorname{Re} F_k(x_i) = r_k \|x_i\| \quad \text{for each } i \in \{1, \dots, n\} \quad \text{and } k \in \{1, \dots, m\}$$

and, from (5.4),

$$c_\infty \left\| \sum_{i=1}^n x_i \right\| = \max_{1 \leq k \leq m} \operatorname{Re} \left[F_k \left(\sum_{i=1}^n x_i \right) \right],$$

which is (5.3).

From (5.5), on summing over $i \in \{1, \dots, n\}$, we get (5.2), and the theorem is proved. \square



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 30 of 99

The following result in normed spaces also holds.

Theorem 5.2. Let $x_i, r_k, F_k, k \in \{1, \dots, m\}, i \in \{1, \dots, n\}$ be as in the hypothesis of Theorem 3.1. Then we have the inequality

$$(5.6) \quad (1 \leq) \frac{\sum_{i=1}^n \|x_i\|}{\|\sum_{i=1}^n x_i\|} \leq \frac{c_p}{(\sum_{k=1}^m r_k^p)^{\frac{1}{p}}} \left(\leq \frac{\sum_{k=1}^m \|F_k\|^p}{\sum_{k=1}^m r_k^p} \right)^{\frac{1}{p}},$$

where $p \geq 1$.

The case of equality holds in (5.6) if and only if


$$(5.7) \quad \operatorname{Re} \left[F_k \left(\sum_{i=1}^n x_i \right) \right] = r_k \sum_{i=1}^n \|x_i\| \quad \text{for each } k \in \{1, \dots, m\}$$

and

$$(5.8) \quad \sum_{k=1}^m \left[\operatorname{Re} F_k \left(\sum_{i=1}^n x_i \right) \right]^p = c_p^p \left\| \sum_{i=1}^n x_i \right\|^p.$$

Proof. By the definition of $c_p, p \geq 1$, we have

$$c_p^p \|x\|^p \geq \sum_{k=1}^m |F_k(x)|^p \quad \text{for any } x \in X,$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 31 of 99

implying that

$$(5.9) \quad c_p^p \left\| \sum_{i=1}^n x_i \right\|^p \geq \sum_{k=1}^m \left| F_k \left(\sum_{i=1}^n x_i \right) \right|^p \geq \sum_{k=1}^m \left| \operatorname{Re} F_k \left(\sum_{i=1}^n x_i \right) \right|^p \\ \geq \sum_{k=1}^m \left[\operatorname{Re} F_k \left(\sum_{i=1}^n x_i \right) \right]^p = \sum_{k=1}^m \left[\sum_{i=1}^n \operatorname{Re} F_k(x_i) \right]^p.$$

Utilising the hypothesis (3.1), we obviously have that

$$(5.10) \quad \sum_{k=1}^m \left[\sum_{i=1}^n \operatorname{Re} F_k(x_i) \right]^p \geq \sum_{k=1}^m \left[\sum_{i=1}^n r_k \|x_i\| \right]^p = \sum_{k=1}^m r_k^p \left(\sum_{i=1}^n \|x_i\| \right)^p.$$

Making use of (5.9) and (5.10), we deduce


$$c_p^p \left\| \sum_{i=1}^n x_i \right\|^p \geq \left(\sum_{k=1}^m r_k^p \right) \left(\sum_{i=1}^n \|x_i\| \right)^p,$$

which implies the desired inequality (5.6).

If (5.7) holds true, then, taking the power p and summing over $k \in \{1, \dots, m\}$, we deduce

$$\sum_{k=1}^m \left[\operatorname{Re} \left[F_k \left(\sum_{i=1}^n x_i \right) \right] \right]^p = \sum_{k=1}^m r_k^p \left(\sum_{i=1}^n \|x_i\| \right)^p,$$

which, together with (5.8) shows that the equality case holds true in (5.6).



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 32 of 99

Conversely, if the case of equality holds in (5.6), then it must hold in all inequalities needed to prove (5.6), therefore, we must have:


$$(5.11) \quad \operatorname{Re} F_k(x_i) = r_k \|x_i\| \quad \text{for each } i \in \{1, \dots, n\} \quad \text{and } k \in \{1, \dots, m\}$$

and, from (5.9),

$$c_p^p \left\| \sum_{i=1}^n x_i \right\|^p = \sum_{k=1}^m \left[\operatorname{Re} F_k \left(\sum_{i=1}^n x_i \right) \right]^p,$$

which is exactly (5.8).

From (5.11), on summing over i from 1 to n , we deduce (5.7), and the theorem is proved. \square



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 33 of 99

6. An Additive Reverse for the Triangle Inequality

6.1. The Case of One Functional

In the following we provide an alternative of the Diaz-Metcalf reverse of the generalised triangle inequality [6].

Theorem 6.1 (Dragomir, 2004). *Let $(X, \|\cdot\|)$ be a normed linear space over the real or complex number field \mathbb{K} and $F : X \rightarrow \mathbb{K}$ a linear functional with the property that $|F(x)| \leq \|x\|$ for any $x \in X$. If $x_i \in X$, $k_i \geq 0$, $i \in \{1, \dots, n\}$ are such that*

$$(6.1) \quad (0 \leq) \|x_i\| - \operatorname{Re} F(x_i) \leq k_i \text{ for each } i \in \{1, \dots, n\},$$

then we have the inequality


$$(6.2) \quad (0 \leq) \sum_{i=1}^n \|x_i\| - \left\| \sum_{i=1}^n x_i \right\| \leq \sum_{i=1}^n k_i.$$

The equality holds in (6.2) if and only if both

$$(6.3) \quad F\left(\sum_{i=1}^n x_i\right) = \left\| \sum_{i=1}^n x_i \right\| \text{ and } F\left(\sum_{i=1}^n x_i\right) = \sum_{i=1}^n \|x_i\| - \sum_{i=1}^n k_i.$$

Proof. If we sum in (6.1) over i from 1 to n , then we get

$$(6.4) \quad \sum_{i=1}^n \|x_i\| \leq \operatorname{Re} \left[F\left(\sum_{i=1}^n x_i\right) \right] + \sum_{i=1}^n k_i.$$



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 34 of 99

Taking into account that $|F(x)| \leq \|x\|$ for each $x \in X$, then we may state that

$$(6.5) \quad \operatorname{Re} \left[F \left(\sum_{i=1}^n x_i \right) \right] \leq \left| \operatorname{Re} F \left(\sum_{i=1}^n x_i \right) \right| \\ \leq \left| F \left(\sum_{i=1}^n x_i \right) \right| \leq \left\| \sum_{i=1}^n x_i \right\|.$$

Now, making use of (6.4) and (6.5), we deduce (6.2).

Obviously, if (6.3) is valid, then the case of equality in (6.2) holds true.

Conversely, if the equality holds in (6.2), then it must hold in all the inequalities used to prove (6.2), therefore we have

$$\sum_{i=1}^n \|x_i\| = \operatorname{Re} \left[F \left(\sum_{i=1}^n x_i \right) \right] + \sum_{i=1}^n k_i$$

and


$$\operatorname{Re} \left[F \left(\sum_{i=1}^n x_i \right) \right] = \left| F \left(\sum_{i=1}^n x_i \right) \right| = \left\| \sum_{i=1}^n x_i \right\|,$$

which imply (6.3). □

The following corollary may be stated [6].

Corollary 6.2. *Let $(X, \|\cdot\|)$ be a normed linear space, $[\cdot, \cdot] : X \times X \rightarrow \mathbb{K}$ a semi-inner product generating the norm $\|\cdot\|$ and $e \in X$, $\|e\| = 1$. If $x_i \in X$, $k_i \geq 0$, $i \in \{1, \dots, n\}$ are such that*

$$(6.6) \quad (0 \leq) \|x_i\| - \operatorname{Re}[x_i, e] \leq k_i \text{ for each } i \in \{1, \dots, n\},$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 35 of 99

then we have the inequality

$$(6.7) \quad (0 \leq) \sum_{i=1}^n \|x_i\| - \left\| \sum_{i=1}^n x_i \right\| \leq \sum_{i=1}^n k_i.$$

The equality holds in (6.7) if and only if both

$$(6.8) \quad \left[\sum_{i=1}^n x_i, e \right] = \left\| \sum_{i=1}^n x_i \right\| \quad \text{and} \quad \left[\sum_{i=1}^n x_i, e \right] = \sum_{i=1}^n \|x_i\| - \sum_{i=1}^n k_i.$$

Moreover, if $(X, \|\cdot\|)$ is strictly convex, then the case of equality holds in (6.7) if and only if


$$(6.9) \quad \sum_{i=1}^n \|x_i\| \geq \sum_{i=1}^n k_i$$

and

$$(6.10) \quad \sum_{i=1}^n x_i = \left(\sum_{i=1}^n \|x_i\| - \sum_{i=1}^n k_i \right) \cdot e.$$

Proof. The first part of the corollary is obvious by Theorem 6.1 applied for the continuous linear functional of unit norm F_e , $F_e(x) = [x, e]$, $x \in X$. The second part may be shown on utilising a similar argument to the one from the proof of Corollary 4.3. We omit the details. \square

Remark 3. If $X = H$, $(H; \langle \cdot, \cdot \rangle)$ is an inner product space, then from Corollary 6.2 we deduce the additive reverse inequality obtained in Theorem 7 of [12]. For further similar results in inner product spaces, see [4] and [12].



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 36 of 99

6.2. The Case of m Functionals

The following result generalising Theorem 6.1 may be stated [6].

Theorem 6.3 (Dragomir, 2004). *Let $(X, \|\cdot\|)$ be a normed linear space over the real or complex number field \mathbb{K} . If F_k , $k \in \{1, \dots, m\}$ are bounded linear functionals defined on X and $x_i \in X$, $M_{ik} \geq 0$ for $i \in \{1, \dots, n\}$, $k \in \{1, \dots, m\}$ are such that*

$$(6.11) \quad \|x_i\| - \operatorname{Re} F_k(x_i) \leq M_{ik}$$

for each $i \in \{1, \dots, n\}$, $k \in \{1, \dots, m\}$, then we have the inequality

$$(6.12) \quad \sum_{i=1}^n \|x_i\| \leq \left\| \frac{1}{m} \sum_{k=1}^m F_k \right\| \left\| \sum_{i=1}^n x_i \right\| + \frac{1}{m} \sum_{k=1}^m \sum_{i=1}^n M_{ik}.$$

The case of equality holds in (6.12) if both


$$(6.13) \quad \frac{1}{m} \sum_{k=1}^m F_k \left(\sum_{i=1}^n x_i \right) = \left\| \frac{1}{m} \sum_{k=1}^m F_k \right\| \left\| \sum_{i=1}^n x_i \right\|$$

and

$$(6.14) \quad \frac{1}{m} \sum_{k=1}^m F_k \left(\sum_{i=1}^n x_i \right) = \sum_{i=1}^n \|x_i\| - \frac{1}{m} \sum_{k=1}^m \sum_{j=1}^n M_{jk}.$$

Proof. If we sum (6.11) over i from 1 to n , then we deduce

$$\sum_{i=1}^n \|x_i\| - \operatorname{Re} F_k \left(\sum_{i=1}^n x_i \right) \leq \sum_{i=1}^n M_{ik}$$



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 37 of 99

for each $k \in \{1, \dots, m\}$.

Summing these inequalities over k from 1 to m , we deduce

$$(6.15) \quad \sum_{i=1}^n \|x_i\| \leq \frac{1}{m} \sum_{k=1}^m \operatorname{Re} F_k \left(\sum_{i=1}^n x_i \right) + \frac{1}{m} \sum_{k=1}^m \sum_{i=1}^n M_{ik}.$$

Utilising the continuity property of the functionals F_k and the properties of the modulus, we have


$$(6.16) \quad \begin{aligned} \sum_{k=1}^m \operatorname{Re} F_k \left(\sum_{i=1}^n x_i \right) &\leq \left| \sum_{k=1}^m \operatorname{Re} F_k \left(\sum_{i=1}^n x_i \right) \right| \\ &\leq \left| \sum_{k=1}^m F_k \left(\sum_{i=1}^n x_i \right) \right| \\ &\leq \left\| \sum_{k=1}^m F_k \right\| \left\| \sum_{i=1}^n x_i \right\|. \end{aligned}$$

Now, by (6.15) and (6.16), we deduce (6.12).

Obviously, if (6.13) and (6.14) hold true, then the case of equality is valid in (6.12).

Conversely, if the case of equality holds in (6.12), then it must hold in all the inequalities used to prove (6.12). Therefore we have

$$\sum_{i=1}^n \|x_i\| = \frac{1}{m} \sum_{k=1}^m \operatorname{Re} F_k \left(\sum_{i=1}^n x_i \right) + \frac{1}{m} \sum_{k=1}^m \sum_{i=1}^n M_{ik},$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 38 of 99

$$\sum_{k=1}^m \operatorname{Re} F_k \left(\sum_{i=1}^n x_i \right) = \left\| \sum_{k=1}^m F_k \right\| \left\| \sum_{i=1}^n x_i \right\|$$

and

$$\sum_{k=1}^m \operatorname{Im} F_k \left(\sum_{i=1}^n x_i \right) = 0.$$

These imply that (6.13) and (6.14) hold true, and the theorem is completely proved. \square

Remark 4. If $F_k, k \in \{1, \dots, m\}$ are of unit norm, then, from (6.12), we deduce the inequality

$$(6.17) \quad \sum_{i=1}^n \|x_i\| \leq \left\| \sum_{i=1}^n x_i \right\| + \frac{1}{m} \sum_{k=1}^m \sum_{i=1}^n M_{ik},$$


which is obviously coarser than (6.12), but perhaps more useful for applications.

6.3. The Case of Inner Product Spaces

The case of inner product spaces, in which we may provide a simpler condition of equality, is of interest in applications [6].

Theorem 6.4 (Dragomir, 2004). Let $(X, \|\cdot\|)$ be an inner product space over the real or complex number field \mathbb{K} , $e_k, x_i \in H \setminus \{0\}$, $k \in \{1, \dots, m\}$, $i \in \{1, \dots, n\}$. If $M_{ik} \geq 0$ for $i \in \{1, \dots, n\}$, $\{1, \dots, n\}$ such that

$$(6.18) \quad \|x_i\| - \operatorname{Re} \langle x_i, e_k \rangle \leq M_{ik}$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 39 of 99

for each $i \in \{1, \dots, n\}$, $k \in \{1, \dots, m\}$, then we have the inequality

$$(6.19) \quad \sum_{i=1}^n \|x_i\| \leq \left\| \frac{1}{m} \sum_{k=1}^m e_k \right\| \left\| \sum_{i=1}^n x_i \right\| + \frac{1}{m} \sum_{k=1}^m \sum_{i=1}^n M_{ik}.$$

The case of equality holds in (6.19) if and only if

$$(6.20) \quad \sum_{i=1}^n \|x_i\| \geq \frac{1}{m} \sum_{k=1}^m \sum_{i=1}^n M_{ik}$$

and


$$(6.21) \quad \sum_{i=1}^n x_i = \frac{m \left(\sum_{i=1}^n \|x_i\| - \frac{1}{m} \sum_{k=1}^m \sum_{i=1}^n M_{ik} \right)}{\left\| \sum_{k=1}^m e_k \right\|^2} \sum_{k=1}^m e_k.$$

Proof. As in the proof of Theorem 6.3, we have

$$(6.22) \quad \sum_{i=1}^n \|x_i\| \leq \operatorname{Re} \left\langle \frac{1}{m} \sum_{k=1}^m e_k, \sum_{i=1}^n x_i \right\rangle + \frac{1}{m} \sum_{k=1}^m \sum_{i=1}^n M_{ik},$$

and $\sum_{k=1}^m e_k \neq 0$.

On utilising the Schwarz inequality in the inner product space $(H; \langle \cdot, \cdot \rangle)$ for



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 40 of 99

$\sum_{i=1}^n x_i, \sum_{k=1}^m e_k$, we have

$$(6.23) \quad \left\| \sum_{i=1}^n x_i \right\| \left\| \sum_{k=1}^m e_k \right\| \geq \left| \left\langle \sum_{i=1}^n x_i, \sum_{k=1}^m e_k \right\rangle \right|$$

$$\geq \left| \operatorname{Re} \left\langle \sum_{i=1}^n x_i, \sum_{k=1}^m e_k \right\rangle \right|$$

$$\geq \operatorname{Re} \left\langle \sum_{i=1}^n x_i, \sum_{k=1}^m e_k \right\rangle.$$

By (6.22) and (6.23) we deduce (6.19).

Taking the norm in (6.21) and using (6.20), we have

$$\left\| \sum_{i=1}^n x_i \right\| = \frac{m \left(\sum_{i=1}^n \|x_i\| - \frac{1}{m} \sum_{k=1}^m \sum_{i=1}^n M_{ik} \right)}{\left\| \sum_{k=1}^m e_k \right\|},$$


showing that the equality holds in (6.19).

Conversely, if the case of equality holds in (6.19), then it must hold in all the inequalities used to prove (6.19). Therefore we have

$$(6.24) \quad \|x_i\| = \operatorname{Re} \langle x_i, e_k \rangle + M_{ik}$$

for each $i \in \{1, \dots, n\}$, $k \in \{1, \dots, m\}$,

$$(6.25) \quad \left\| \sum_{i=1}^n x_i \right\| \left\| \sum_{k=1}^m e_k \right\| = \left| \left\langle \sum_{i=1}^n x_i, \sum_{k=1}^m e_k \right\rangle \right|$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 41 of 99

and

$$(6.26) \quad \operatorname{Im} \left\langle \sum_{i=1}^n x_i, \sum_{k=1}^m e_k \right\rangle = 0.$$

From (6.24), on summing over i and k , we get

$$(6.27) \quad \operatorname{Re} \left\langle \sum_{i=1}^n x_i, \sum_{k=1}^m e_k \right\rangle = m \sum_{i=1}^n \|x_i\| - \sum_{k=1}^m \sum_{i=1}^n M_{ik}.$$

On the other hand, by the use of the identity (3.22), the relation (6.25) holds if and only if

$$\sum_{i=1}^n x_i = \frac{\langle \sum_{i=1}^n x_i, \sum_{k=1}^m e_k \rangle}{\|\sum_{k=1}^m e_k\|^2} \sum_{k=1}^m e_k,$$

giving, from (6.26) and (6.27), that

$$\sum_{i=1}^n x_i = \frac{m \sum_{i=1}^n \|x_i\| - \sum_{k=1}^m \sum_{i=1}^n M_{ik}}{\|\sum_{k=1}^m e_k\|^2} \sum_{k=1}^m e_k.$$

If the inequality holds in (6.19), then obviously (6.20) is valid, and the theorem is proved. \square

Remark 5. If in the above theorem the vectors $\{e_k\}_{k=1, \overline{m}}$ are assumed to be orthogonal, then (6.19) becomes:

$$(6.28) \quad \sum_{i=1}^n \|x_i\| \leq \frac{1}{m} \left(\sum_{k=1}^m \|e_k\|^2 \right)^{\frac{1}{2}} \left\| \sum_{i=1}^n x_i \right\| + \frac{1}{m} \sum_{k=1}^m \sum_{i=1}^n M_{ik}.$$



Title Page

Contents



Go Back

Close

Quit

Page 42 of 99

Moreover, if $\{e_k\}_{k=1,m}$ is an orthonormal family, then (6.28) becomes

$$(6.29) \quad \sum_{i=1}^n \|x_i\| \leq \frac{\sqrt{m}}{m} \left\| \sum_{i=1}^n x_i \right\| + \frac{1}{m} \sum_{k=1}^m \sum_{i=1}^n M_{ik},$$

which has been obtained in [12].

Before we provide some natural consequences of Theorem 6.4, we need some preliminary results concerning another reverse of Schwarz's inequality in inner product spaces (see for instance [4, p. 27]).

Lemma 6.5 (Dragomir, 2004). *Let $(X, \|\cdot\|)$ be an inner product space over the real or complex number field \mathbb{K} and $x, a \in H, r > 0$. If $\|x - a\| \leq r$, then we have the inequality*

$$(6.30) \quad \|x\| \|a\| - \operatorname{Re} \langle x, a \rangle \leq \frac{1}{2} r^2.$$

The case of equality holds in (6.30) if and only if

$$(6.31) \quad \|x - a\| = r \text{ and } \|x\| = \|a\|.$$

Proof. The condition $\|x - a\| \leq r$ is clearly equivalent to


$$(6.32) \quad \|x\|^2 + \|a\|^2 \leq 2 \operatorname{Re} \langle x, a \rangle + r^2.$$

Since

$$(6.33) \quad 2 \|x\| \|a\| \leq \|x\|^2 + \|a\|^2,$$

with equality if and only if $\|x\| = \|a\|$, hence by (6.32) and (6.33) we deduce (6.30).

The case of equality is obvious. □



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 43 of 99

Utilising the above lemma we may state the following corollary of Theorem 6.4 [6].

Corollary 6.6. *Let $(H; \langle \cdot, \cdot \rangle)$, e_k , x_i be as in Theorem 6.4. If $r_{ik} > 0$, $i \in \{1, \dots, n\}$, $k \in \{1, \dots, m\}$ such that*

$$(6.34) \quad \|x_i - e_k\| \leq r_{ik} \text{ for each } i \in \{1, \dots, n\} \text{ and } k \in \{1, \dots, m\},$$

then we have the inequality

$$(6.35) \quad \sum_{i=1}^n \|x_i\| \leq \left\| \frac{1}{m} \sum_{k=1}^m e_k \right\| \left\| \sum_{i=1}^n x_i \right\| + \frac{1}{2m} \sum_{k=1}^m \sum_{i=1}^n r_{ik}^2.$$

The equality holds in (6.35) if and only if

$$\sum_{i=1}^n \|x_i\| \geq \frac{1}{2m} \sum_{k=1}^m \sum_{i=1}^n r_{ik}^2$$


and

$$\sum_{i=1}^n x_i = \frac{m \left(\sum_{i=1}^n \|x_i\| - \frac{1}{2m} \sum_{k=1}^m \sum_{i=1}^n r_{ik}^2 \right)}{\left\| \sum_{k=1}^m e_k \right\|^2} \sum_{k=1}^m e_k.$$

The following lemma may provide another sufficient condition for (6.18) to hold (see also [4, p. 28]).

Lemma 6.7 (Dragomir, 2004). *Let $(H; \langle \cdot, \cdot \rangle)$ be an inner product space over the real or complex number field \mathbb{K} and $x, y \in H$, $M \geq m > 0$. If either*

$$(6.36) \quad \operatorname{Re} \langle My - x, x - my \rangle \geq 0$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 44 of 99

or, equivalently,

$$(6.37) \quad \left\| x - \frac{m+M}{2}y \right\| \leq \frac{1}{2} (M-m) \|y\|,$$

holds, then

$$(6.38) \quad \|x\| \|y\| - \operatorname{Re} \langle x, y \rangle \leq \frac{1}{4} \cdot \frac{(M-m)^2}{m+M} \|y\|^2.$$

The case of equality holds in (6.38) if and only if the equality case is realised in (6.36) and

$$\|x\| = \frac{M+m}{2} \|y\|.$$

The proof is obvious by Lemma 6.5 for $a = \frac{M+m}{2}y$ and $r = \frac{1}{2} (M-m) \|y\|$.


Finally, the following corollary of Theorem 6.4 may be stated [6].

Corollary 6.8. Assume that $(H, \langle \cdot, \cdot \rangle)$, e_k , x_i are as in Theorem 6.4. If $M_{ik} \geq m_{ik} > 0$ satisfy the condition

$$\operatorname{Re} \langle M_k e_k - x_i, x_i - \mu_k e_k \rangle \geq 0$$

for each $i \in \{1, \dots, n\}$ and $k \in \{1, \dots, m\}$, then

$$\sum_{i=1}^n \|x_i\| \leq \left\| \frac{1}{m} \sum_{k=1}^m e_k \right\| \left\| \sum_{i=1}^n x_i \right\| + \frac{1}{4m} \sum_{k=1}^m \sum_{i=1}^n \frac{(M_{ik} - m_{ik})^2}{M_{ik} + m_{ik}} \|e_k\|^2.$$



Title Page

Contents



Go Back

Close

Quit

Page 45 of 99

7. Other Additive Reverses for m Functionals

A different approach in obtaining other additive reverses for the generalised triangle inequality is incorporated in the following new result:

Theorem 7.1. *Let $(X, \|\cdot\|)$ be a normed linear space over the real or complex number field \mathbb{K} . Assume $F_k, k \in \{1, \dots, m\}$, are bounded linear functionals on the normed linear space X and $x_i \in X, i \in \{1, \dots, n\}, M_{ik} \geq 0, i \in \{1, \dots, n\}, k \in \{1, \dots, m\}$ are such that*

$$(7.1) \quad \|x_i\| - \operatorname{Re} F_k(x_i) \leq M_{ik}$$

for each $i \in \{1, \dots, n\}$ and $k \in \{1, \dots, m\}$.

(i) If c_∞ is defined by (c_∞) , then we have the inequality


$$(7.2) \quad \sum_{i=1}^n \|x_i\| \leq c_\infty \left\| \sum_{i=1}^n x_i \right\| + \frac{1}{m} \sum_{k=1}^m \sum_{i=1}^n M_{ik}.$$

(ii) If c_p is defined by (c_p) for $p \geq 1$, then we have the inequality:

$$(7.3) \quad \sum_{i=1}^n \|x_i\| \leq \frac{1}{m^{\frac{1}{p}}} c_p \left\| \sum_{i=1}^n x_i \right\| + \frac{1}{m} \sum_{k=1}^m \sum_{i=1}^n M_{ik}.$$

Proof. (i) Since

$$\max_{1 \leq k \leq m} \|F_k(x)\| \leq c_\infty \|x\| \quad \text{for any } x \in X,$$



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 46 of 99

then we have

$$\sum_{k=1}^m \left| F_k \left(\sum_{i=1}^n x_i \right) \right| \leq m \max_{1 \leq k \leq m} \left| F_k \left(\sum_{i=1}^n x_i \right) \right| \leq m c_\infty \left\| \sum_{i=1}^n x_i \right\|.$$

Using (6.16), we may state that

$$\frac{1}{m} \sum_{k=1}^m \operatorname{Re} F_k \left(\sum_{i=1}^n x_i \right) \leq c_\infty \left\| \sum_{i=1}^n x_i \right\|,$$

which, together with (6.15) imply the desired inequality (7.2).

(ii) Using the fact that, obviously


$$\left(\sum_{k=1}^m |F_k(x)|^p \right)^{\frac{1}{p}} \leq c_p \|x\| \quad \text{for any } x \in X,$$

then, by Hölder's inequality for $p > 1$, $\frac{1}{p} + \frac{1}{q} = 1$, we have

$$\begin{aligned} \sum_{k=1}^m \left| F_k \left(\sum_{i=1}^n x_i \right) \right| &\leq m^{\frac{1}{q}} \left(\sum_{k=1}^m \left| F_k \left(\sum_{i=1}^n x_i \right) \right|^p \right)^{\frac{1}{p}} \\ &\leq c_p m^{\frac{1}{q}} \left\| \sum_{i=1}^n x_i \right\|, \end{aligned}$$

which, combined with (6.15) and (6.16) will give the desired inequality (7.3).

The case $p = 1$ goes likewise and we omit the details. \square



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 47 of 99

Remark 6. Since, obviously $c_\infty \leq \max_{1 \leq k \leq m} \|F_k\|$, then from (7.2) we have

$$(7.4) \quad \sum_{i=1}^n \|x_i\| \leq \max_{1 \leq k \leq m} \{\|F_k\|\} \cdot \left\| \sum_{i=1}^n x_i \right\| + \frac{1}{m} \sum_{k=1}^m \sum_{i=1}^n M_{ik}.$$

Finally, since $c_p \leq (\sum_{k=1}^m \|F_k\|^p)^{\frac{1}{p}}$, $p \geq 1$, hence by (7.3) we have

$$(7.5) \quad \sum_{i=1}^n \|x_i\| \leq \left(\frac{\sum_{k=1}^m \|F_k\|^p}{m} \right)^{\frac{1}{p}} \left\| \sum_{i=1}^n x_i \right\| + \frac{1}{m} \sum_{k=1}^m \sum_{i=1}^n M_{ik}.$$

The following corollary for semi-inner products may be stated as well.


Corollary 7.2. Let $(X, \|\cdot\|)$ be a real or complex normed space and $[\cdot, \cdot] : X \times X \rightarrow \mathbb{K}$ a semi-inner product generating the norm $\|\cdot\|$. Assume $e_k, x_i \in H$ and $M_{ik} \geq 0$, $i \in \{1, \dots, n\}$, $k \in \{1, \dots, m\}$ are such that

$$(7.6) \quad \|x_i\| - \operatorname{Re} [x_i, e_k] \leq M_{ik},$$

for any $i \in \{1, \dots, n\}$, $k \in \{1, \dots, m\}$.

(i) If

$$d_\infty := \sup_{x \neq 0} \left\{ \frac{\max_{1 \leq k \leq n} |[x, e_k]|}{\|x\|} \right\} \left(\leq \max_{1 \leq k \leq n} \|e_k\| \right),$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents




Go Back

Close

Quit

Page 48 of 99



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 49 of 99

then

$$(7.7) \quad \sum_{i=1}^n \|x_i\| \leq d_\infty \left\| \sum_{i=1}^n x_i \right\| + \frac{1}{m} \sum_{k=1}^m \sum_{i=1}^n M_{ik} \\ \left(\leq \max_{1 \leq k \leq n} \|e_k\| \cdot \left\| \sum_{i=1}^n x_i \right\| + \frac{1}{m} \sum_{k=1}^m \sum_{i=1}^n M_{ik} \right);$$

(ii) If

$$d_p := \sup_{x \neq 0} \left\{ \frac{\sum_{k=1}^m |[x, e_k]|^p}{\|x\|^p} \right\}^{\frac{1}{p}} \left(\leq \left(\sum_{k=1}^m \|e_k\|^p \right)^{\frac{1}{p}} \right),$$

where $p \geq 1$, then

$$(7.8) \quad \sum_{i=1}^n \|x_i\| \leq \frac{1}{m^{\frac{1}{p}}} d_p \left\| \sum_{i=1}^n x_i \right\| + \frac{1}{m} \sum_{k=1}^m \sum_{i=1}^n M_{ik} \\ \left(\leq \left(\frac{\sum_{k=1}^m \|e_k\|^p}{m} \right)^{\frac{1}{p}} \left\| \sum_{i=1}^n x_i \right\| + \frac{1}{m} \sum_{k=1}^m \sum_{i=1}^n M_{ik} \right).$$

8. Applications for Complex Numbers

Let \mathbb{C} be the field of complex numbers. If $z = \operatorname{Re} z + i \operatorname{Im} z$, then by $|\cdot|_p : \mathbb{C} \rightarrow [0, \infty)$, $p \in [1, \infty]$ we define the p -modulus of z as

$$|z|_p := \begin{cases} \max \{|\operatorname{Re} z|, |\operatorname{Im} z|\} & \text{if } p = \infty, \\ (|\operatorname{Re} z|^p + |\operatorname{Im} z|^p)^{\frac{1}{p}} & \text{if } p \in [1, \infty), \end{cases}$$

where $|a|$, $a \in \mathbb{R}$ is the usual modulus of the real number a .

For $p = 2$, we recapture the usual modulus of a complex number, i.e.,

$$|z|_2 = \sqrt{|\operatorname{Re} z|^2 + |\operatorname{Im} z|^2} = |z|, \quad z \in \mathbb{C}.$$


It is well known that $(\mathbb{C}, |\cdot|_p)$, $p \in [1, \infty]$ is a Banach space over the *real number field* \mathbb{R} .

Consider the Banach space $(\mathbb{C}, |\cdot|_1)$ and $F : \mathbb{C} \rightarrow \mathbb{C}$, $F(z) = az$ with $a \in \mathbb{C}$, $a \neq 0$. Obviously, F is linear on \mathbb{C} . For $z \neq 0$, we have

$$\frac{|F(z)|}{|z|_1} = \frac{|a||z|}{|z|_1} = \frac{|a| \sqrt{|\operatorname{Re} z|^2 + |\operatorname{Im} z|^2}}{|\operatorname{Re} z| + |\operatorname{Im} z|} \leq |a|.$$

Since, for $z_0 = 1$, we have $|F(z_0)| = |a|$ and $|z_0|_1 = 1$, hence

$$\|F\|_1 := \sup_{z \neq 0} \frac{|F(z)|}{|z|_1} = |a|,$$



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 50 of 99

showing that F is a bounded linear functional on $(\mathbb{C}, |\cdot|_1)$ and $\|F\|_1 = |a|$.

We can apply Theorem 3.1 to state the following reverse of the generalised triangle inequality for complex numbers [5].

Proposition 8.1. *Let $a_k, x_j \in \mathbb{C}$, $k \in \{1, \dots, m\}$ and $j \in \{1, \dots, n\}$. If there exist the constants $r_k \geq 0$, $k \in \{1, \dots, m\}$ with $\sum_{k=1}^m r_k > 0$ and*

$$(8.1) \quad r_k [|\operatorname{Re} x_j| + |\operatorname{Im} x_j|] \leq \operatorname{Re} a_k \cdot \operatorname{Re} x_j - \operatorname{Im} a_k \cdot \operatorname{Im} x_j$$

for each $j \in \{1, \dots, n\}$ and $k \in \{1, \dots, m\}$, then


$$(8.2) \quad \sum_{j=1}^n [|\operatorname{Re} x_j| + |\operatorname{Im} x_j|] \leq \frac{|\sum_{k=1}^m a_k|}{\sum_{k=1}^m r_k} \left[\left| \sum_{j=1}^n \operatorname{Re} x_j \right| + \left| \sum_{j=1}^n \operatorname{Im} x_j \right| \right].$$

The case of equality holds in (8.2) if both

$$\begin{aligned} & \operatorname{Re} \left(\sum_{k=1}^m a_k \right) \operatorname{Re} \left(\sum_{j=1}^n x_j \right) - \operatorname{Im} \left(\sum_{k=1}^m a_k \right) \operatorname{Im} \left(\sum_{j=1}^n x_j \right) \\ &= \left(\sum_{k=1}^m r_k \right) \sum_{j=1}^n [|\operatorname{Re} x_j| + |\operatorname{Im} x_j|] \\ &= \left| \sum_{k=1}^m a_k \right| \left[\left| \sum_{j=1}^n \operatorname{Re} x_j \right| + \left| \sum_{j=1}^n \operatorname{Im} x_j \right| \right]. \end{aligned}$$

The proof follows by Theorem 3.1 applied for the Banach space $(\mathbb{C}, |\cdot|_1)$ and $F_k(z) = a_k z$, $k \in \{1, \dots, m\}$ on taking into account that:

$$\left\| \sum_{k=1}^m F_k \right\|_1 = \left| \sum_{k=1}^m a_k \right|.$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 51 of 99

Now, consider the Banach space $(\mathbb{C}, |\cdot|_\infty)$. If $F(z) = dz$, then for $z \neq 0$ we have

$$\frac{|F(z)|}{|z|_\infty} = \frac{|d||z|}{|z|_\infty} = \frac{|d| \sqrt{|\operatorname{Re} z|^2 + |\operatorname{Im} z|^2}}{\max\{|\operatorname{Re} z|, |\operatorname{Im} z|\}} \leq \sqrt{2}|d|.$$

Since, for $z_0 = 1 + i$, we have $|F(z_0)| = \sqrt{2}|d|$, $|z_0|_\infty = 1$, hence

$$\|F\|_\infty := \sup_{z \neq 0} \frac{|F(z)|}{|z|_\infty} = \sqrt{2}|d|,$$

showing that F is a bounded linear functional on $(\mathbb{C}, |\cdot|_\infty)$ and $\|F\|_\infty = \sqrt{2}|d|$.


If we apply Theorem 3.1, then we can state the following reverse of the generalised triangle inequality for complex numbers [5].

Proposition 8.2. *Let $a_k, x_j \in \mathbb{C}$, $k \in \{1, \dots, m\}$ and $j \in \{1, \dots, n\}$. If there exist the constants $r_k \geq 0$, $k \in \{1, \dots, m\}$ with $\sum_{k=1}^m r_k > 0$ and*

$$r_k \max\{|\operatorname{Re} x_j|, |\operatorname{Im} x_j|\} \leq \operatorname{Re} a_k \cdot \operatorname{Re} x_j - \operatorname{Im} a_k \cdot \operatorname{Im} x_j$$

for each $j \in \{1, \dots, n\}$ and $k \in \{1, \dots, m\}$, then

$$(8.3) \quad \sum_{j=1}^n \max\{|\operatorname{Re} x_j|, |\operatorname{Im} x_j|\} \leq \sqrt{2} \cdot \frac{|\sum_{k=1}^m a_k|}{\sum_{k=1}^m r_k} \max \left\{ \left| \sum_{j=1}^n \operatorname{Re} x_j \right|, \left| \sum_{j=1}^n \operatorname{Im} x_j \right| \right\}.$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 52 of 99

The case of equality holds in (8.3) if both

$$\begin{aligned} \operatorname{Re} \left(\sum_{k=1}^m a_k \right) \operatorname{Re} \left(\sum_{j=1}^n x_j \right) - \operatorname{Im} \left(\sum_{k=1}^m a_k \right) \operatorname{Im} \left(\sum_{j=1}^n x_j \right) \\ = \left(\sum_{k=1}^m r_k \right) \sum_{j=1}^n \max \{ |\operatorname{Re} x_j|, |\operatorname{Im} x_j| \} \\ = \sqrt{2} \left| \sum_{k=1}^m a_k \right| \max \left\{ \left| \sum_{j=1}^n \operatorname{Re} x_j \right|, \left| \sum_{j=1}^n \operatorname{Im} x_j \right| \right\}. \end{aligned}$$

Finally, consider the Banach space $(\mathbb{C}, |\cdot|_{2p})$ with $p \geq 1$.

Let $F : \mathbb{C} \rightarrow \mathbb{C}$, $F(z) = cz$. By Hölder's inequality, we have


$$\frac{|F(z)|}{|z|_{2p}} = \frac{|c| \sqrt{|\operatorname{Re} z|^2 + |\operatorname{Im} z|^2}}{(|\operatorname{Re} z|^{2p} + |\operatorname{Im} z|^{2p})^{\frac{1}{2p}}} \leq 2^{\frac{1}{2} - \frac{1}{2p}} |c|.$$

Since, for $z_0 = 1 + i$ we have $|F(z_0)| = 2^{\frac{1}{2}} |c|$, $|z_0|_{2p} = 2^{\frac{1}{2p}}$ ($p \geq 1$), hence

$$\|F\|_{2p} := \sup_{z \neq 0} \frac{|F(z)|}{|z|_{2p}} = 2^{\frac{1}{2} - \frac{1}{2p}} |c|,$$

showing that F is a bounded linear functional on $(\mathbb{C}, |\cdot|_{2p})$, $p \geq 1$ and $\|F\|_{2p} = 2^{\frac{1}{2} - \frac{1}{2p}} |c|$.

If we apply Theorem 3.1, then we can state the following proposition [5].



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 53 of 99

Proposition 8.3. Let $a_k, x_j \in \mathbb{C}$, $k \in \{1, \dots, m\}$ and $j \in \{1, \dots, n\}$. If there exist the constants $r_k \geq 0$, $k \in \{1, \dots, m\}$ with $\sum_{k=1}^m r_k > 0$ and


$$r_k [|\operatorname{Re} x_j|^{2p} + |\operatorname{Im} x_j|^{2p}]^{\frac{1}{2p}} \leq \operatorname{Re} a_k \cdot \operatorname{Re} x_j - \operatorname{Im} a_k \cdot \operatorname{Im} x_j$$

for each $j \in \{1, \dots, n\}$ and $k \in \{1, \dots, m\}$, then

$$(8.4) \quad \sum_{j=1}^n [|\operatorname{Re} x_j|^{2p} + |\operatorname{Im} x_j|^{2p}]^{\frac{1}{2p}} \leq 2^{\frac{1}{2} - \frac{1}{2p}} \frac{\left| \sum_{k=1}^m a_k \right|}{\sum_{k=1}^m r_k} \left[\left| \sum_{j=1}^n \operatorname{Re} x_j \right|^{2p} + \left| \sum_{j=1}^n \operatorname{Im} x_j \right|^{2p} \right]^{\frac{1}{2p}}.$$

The case of equality holds in (8.4) if both:

$$\begin{aligned} & \operatorname{Re} \left(\sum_{k=1}^m a_k \right) \operatorname{Re} \left(\sum_{j=1}^n x_j \right) - \operatorname{Im} \left(\sum_{k=1}^m a_k \right) \operatorname{Im} \left(\sum_{j=1}^n x_j \right) \\ &= \left(\sum_{k=1}^m r_k \right) \sum_{j=1}^n [|\operatorname{Re} x_j|^{2p} + |\operatorname{Im} x_j|^{2p}]^{\frac{1}{2p}} \\ &= 2^{\frac{1}{2} - \frac{1}{2p}} \left| \sum_{k=1}^m a_k \right| \left[\left| \sum_{j=1}^n \operatorname{Re} x_j \right|^{2p} + \left| \sum_{j=1}^n \operatorname{Im} x_j \right|^{2p} \right]^{\frac{1}{2p}}. \end{aligned}$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 54 of 99

Remark 7. If in the above proposition we choose $p = 1$, then we have the following reverse of the generalised triangle inequality for complex numbers

$$\sum_{j=1}^n |x_j| \leq \frac{|\sum_{k=1}^m a_k|}{\sum_{k=1}^m r_k} \left| \sum_{j=1}^n x_j \right|$$

provided $x_j, a_k, j \in \{1, \dots, n\}, k \in \{1, \dots, m\}$ satisfy the assumption

$$r_k |x_j| \leq \operatorname{Re} a_k \cdot \operatorname{Re} x_j - \operatorname{Im} a_k \cdot \operatorname{Im} x_j$$

for each $j \in \{1, \dots, n\}, k \in \{1, \dots, m\}$. Here $|\cdot|$ is the usual modulus of a complex number and $r_k > 0, k \in \{1, \dots, m\}$ are given.


We can apply Theorem 6.3 to state the following reverse of the generalised triangle inequality for complex numbers [6].

Proposition 8.4. Let $a_k, x_j \in \mathbb{C}, k \in \{1, \dots, m\}$ and $j \in \{1, \dots, n\}$. If there exist the constants $M_{jk} \geq 0, k \in \{1, \dots, m\}, j \in \{1, \dots, n\}$ such that

$$(8.5) \quad |\operatorname{Re} x_j| + |\operatorname{Im} x_j| \leq \operatorname{Re} a_k \cdot \operatorname{Re} x_j - \operatorname{Im} a_k \cdot \operatorname{Im} x_j + M_{jk}$$

for each $j \in \{1, \dots, n\}$ and $k \in \{1, \dots, m\}$, then

$$(8.6) \quad \sum_{j=1}^n [|\operatorname{Re} x_j| + |\operatorname{Im} x_j|] \leq \frac{1}{m} \left| \sum_{k=1}^m a_k \right| \left[\left| \sum_{j=1}^n \operatorname{Re} x_j \right| + \left| \sum_{j=1}^n \operatorname{Im} x_j \right| \right] + \frac{1}{m} \sum_{k=1}^m \sum_{j=1}^n M_{jk}.$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 55 of 99

The proof follows by Theorem 6.3 applied for the Banach space $(\mathbb{C}, |\cdot|_1)$ and $F_k(z) = a_k z$, $k \in \{1, \dots, m\}$ on taking into account that:

$$\left\| \sum_{k=1}^m F_k \right\|_1 = \left| \sum_{k=1}^m a_k \right|.$$

If we apply Theorem 6.3 for the Banach space $(\mathbb{C}, |\cdot|_\infty)$, then we can state the following reverse of the generalised triangle inequality for complex numbers [6].

Proposition 8.5. *Let $a_k, x_j \in \mathbb{C}$, $k \in \{1, \dots, m\}$ and $j \in \{1, \dots, n\}$. If there exist the constants $M_{jk} \geq 0$, $k \in \{1, \dots, m\}$, $j \in \{1, \dots, n\}$ such that*

$$\max \{|\operatorname{Re} x_j|, |\operatorname{Im} x_j|\} \leq \operatorname{Re} a_k \cdot \operatorname{Re} x_j - \operatorname{Im} a_k \cdot \operatorname{Im} x_j + M_{jk}$$

for each $j \in \{1, \dots, n\}$ and $k \in \{1, \dots, m\}$, then

$$(8.7) \quad \sum_{j=1}^n \max \{|\operatorname{Re} x_j|, |\operatorname{Im} x_j|\} \leq \frac{\sqrt{2}}{m} \left| \sum_{k=1}^m a_k \right| \max \left\{ \left| \sum_{j=1}^n \operatorname{Re} x_j \right|, \left| \sum_{j=1}^n \operatorname{Im} x_j \right| \right\} + \frac{1}{m} \sum_{k=1}^m \sum_{j=1}^n M_{jk}.$$

Finally, if we apply Theorem 6.3, for the Banach space $(\mathbb{C}, |\cdot|_{2p})$ with $p \geq 1$, then we can state the following proposition [6].



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 56 of 99

Proposition 8.6. Let a_k, x_j, M_{jk} be as in Proposition 8.5. If

$$\left[|\operatorname{Re} x_j|^{2p} + |\operatorname{Im} x_j|^{2p} \right]^{\frac{1}{2p}} \leq \operatorname{Re} a_k \cdot \operatorname{Re} x_j - \operatorname{Im} a_k \cdot \operatorname{Im} x_j + M_{jk}$$

for each $j \in \{1, \dots, n\}$ and $k \in \{1, \dots, m\}$, then

$$(8.8) \quad \sum_{j=1}^n \left[|\operatorname{Re} x_j|^{2p} + |\operatorname{Im} x_j|^{2p} \right]^{\frac{1}{2p}} \\ \leq \frac{2^{\frac{1}{2} - \frac{1}{2p}}}{m} \left| \sum_{k=1}^m a_k \right| \left[\left| \sum_{j=1}^n \operatorname{Re} x_j \right|^{2p} + \left| \sum_{j=1}^n \operatorname{Im} x_j \right|^{2p} \right]^{\frac{1}{2p}} + \frac{1}{m} \sum_{k=1}^m \sum_{j=1}^n M_{jk}.$$

where $p \geq 1$.


Remark 8. If in the above proposition we choose $p = 1$, then we have the following reverse of the generalised triangle inequality for complex numbers

$$\sum_{j=1}^n |x_j| \leq \left| \frac{1}{m} \sum_{k=1}^m a_k \right| \left| \sum_{j=1}^n x_j \right| + \frac{1}{m} \sum_{k=1}^m \sum_{j=1}^n M_{jk}$$

provided $x_j, a_k, j \in \{1, \dots, n\}, k \in \{1, \dots, m\}$ satisfy the assumption

$$|x_j| \leq \operatorname{Re} a_k \cdot \operatorname{Re} x_j - \operatorname{Im} a_k \cdot \operatorname{Im} x_j + M_{jk}$$

for each $j \in \{1, \dots, n\}, k \in \{1, \dots, m\}$. Here $|\cdot|$ is the usual modulus of a complex number and $M_{jk} > 0, j \in \{1, \dots, n\}, k \in \{1, \dots, m\}$ are given.



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 57 of 99

9. Karamata Type Inequalities in Hilbert Spaces

Let $f : [a, b] \rightarrow \mathbb{K}$, $\mathbb{K} = \mathbb{C}$ or \mathbb{R} be a Lebesgue integrable function. The following inequality, which is the continuous version of the *triangle inequality*

$$(9.1) \quad \left| \int_a^b f(x) dx \right| \leq \int_a^b |f(x)| dx,$$

plays a fundamental role in Mathematical Analysis and its applications.

It appears, see [20, p. 492], that the first reverse inequality for (9.1) was obtained by J. Karamata in his book from 1949, [14]. It can be stated as

$$(9.2) \quad \cos \theta \int_a^b |f(x)| dx \leq \left| \int_a^b f(x) dx \right|$$

provided


$$-\theta \leq \arg f(x) \leq \theta, \quad x \in [a, b]$$

for given $\theta \in (0, \frac{\pi}{2})$.

This result has recently been extended by the author for the case of Bochner integrable functions with values in a Hilbert space H (see also [10]):

Theorem 9.1 (Dragomir, 2004). *If $f \in L([a, b]; H)$ (this means that $f : [a, b] \rightarrow H$ is strongly measurable on $[a, b]$ and the Lebesgue integral $\int_a^b \|f(t)\| dt$ is finite), then*

$$(9.3) \quad \int_a^b \|f(t)\| dt \leq K \left\| \int_a^b f(t) dt \right\|,$$



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 58 of 99

provided that f satisfies the condition

$$(9.4) \quad \|f(t)\| \leq K \operatorname{Re} \langle f(t), e \rangle \quad \text{for a.e. } t \in [a, b],$$

where $e \in H$, $\|e\| = 1$ and $K \geq 1$ are given.

The case of equality holds in (9.4) if and only if

$$(9.5) \quad \int_a^b f(t) dt = \frac{1}{K} \left(\int_a^b \|f(t)\| dt \right) e.$$

As some natural consequences of the above results, we have noticed in [10] that, if $\rho \in [0, 1)$ and $f \in L([a, b]; H)$ are such that

$$(9.6) \quad \|f(t) - e\| \leq \rho \quad \text{for a.e. } t \in [a, b],$$

then


$$(9.7) \quad \sqrt{1 - \rho^2} \int_a^b \|f(t)\| dt \leq \left\| \int_a^b f(t) dt \right\|$$

with equality if and only if

$$\int_a^b f(t) dt = \sqrt{1 - \rho^2} \left(\int_a^b \|f(t)\| dt \right) \cdot e.$$

Also, for e as above and if $M \geq m > 0$, $f \in L([a, b]; H)$ such that either

$$(9.8) \quad \operatorname{Re} \langle Me - f(t), f(t) - me \rangle \geq 0$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 59 of 99

or, equivalently,

$$(9.9) \quad \left\| f(t) - \frac{M+m}{2}e \right\| \leq \frac{1}{2}(M-m)$$


for a.e. $t \in [a, b]$, then

$$(9.10) \quad \int_a^b \|f(t)\| dt \leq \frac{M+m}{2\sqrt{mM}} \left\| \int_a^b f(t) dt \right\|,$$

with equality if and only if

$$\int_a^b f(t) dt = \frac{2\sqrt{mM}}{M+m} \left(\int_a^b \|f(t)\| dt \right) \cdot e.$$

The main aim of the following sections is to extend the integral inequalities mentioned above for the case of Banach spaces. Applications for Hilbert spaces and for complex-valued functions are given as well.



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 60 of 99

10. Multiplicative Reverses of the Continuous Triangle Inequality

10.1. The Case of One Functional

Let $(X, \|\cdot\|)$ be a Banach space over the real or complex number field. Then one has the following reverse of the continuous triangle inequality [11].

Theorem 10.1 (Dragomir, 2004). *Let F be a continuous linear functional of unit norm on X . Suppose that the function $f : [a, b] \rightarrow X$ is Bochner integrable on $[a, b]$ and there exists a $r \geq 0$ such that*

$$(10.1) \quad r \|f(t)\| \leq \operatorname{Re} F[f(t)] \quad \text{for a.e. } t \in [a, b].$$

Then


$$(10.2) \quad r \int_a^b \|f(t)\| dt \leq \left\| \int_a^b f(t) dt \right\|,$$

where equality holds in (10.2) if and only if both

$$(10.3) \quad F \left(\int_a^b f(t) dt \right) = r \int_a^b \|f(t)\| dt$$

and

$$(10.4) \quad F \left(\int_a^b f(t) dt \right) = \left\| \int_a^b f(t) dt \right\|.$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 61 of 99

Proof. Since the norm of F is one, then

$$|F(x)| \leq \|x\| \quad \text{for any } x \in X.$$

Applying this inequality for the vector $\int_a^b f(t) dt$, we get

$$(10.5) \quad \left\| \int_a^b f(t) dt \right\| \geq \left| F \left(\int_a^b f(t) dt \right) \right| \\ \geq \left| \operatorname{Re} F \left(\int_a^b f(t) dt \right) \right| = \left| \int_a^b \operatorname{Re} F(f(t)) dt \right|.$$

Now, by integration of (10.1), we obtain

$$(10.6) \quad \int_a^b \operatorname{Re} F(f(t)) dt \geq r \int_a^b \|f(t)\| dt,$$


and by (10.5) and (10.6) we deduce the desired inequality (10.2).

Obviously, if (10.3) and (10.4) hold true, then the equality case holds in (10.2).

Conversely, if the case of equality holds in (10.2), then it must hold in all the inequalities used before in proving this inequality. Therefore, we must have

$$(10.7) \quad r \|f(t)\| = \operatorname{Re} F(f(t)) \quad \text{for a.e. } t \in [a, b],$$

$$(10.8) \quad \operatorname{Im} F \left(\int_a^b f(t) dt \right) = 0$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 62 of 99

and

$$(10.9) \quad \left\| \int_a^b f(t) dt \right\| = \operatorname{Re} F \left(\int_a^b f(t) dt \right).$$

Integrating (10.7) on $[a, b]$, we get

$$(10.10) \quad r \int_a^b \|f(t)\| dt = \operatorname{Re} F \left(\int_a^b f(t) dt \right).$$

On utilising (10.10) and (10.8), we deduce (10.3) while (10.9) and (10.10) would imply (10.4), and the theorem is proved. \square

Corollary 10.2. *Let $(X, \|\cdot\|)$ be a Banach space, $[\cdot, \cdot] : X \times X \rightarrow \mathbb{R}$ a semi-inner product generating the norm $\|\cdot\|$ and $e \in X, \|e\| = 1$. Suppose that the function $f : [a, b] \rightarrow X$ is Bochner integrable on $[a, b]$ and there exists a $r \geq 0$ such that*

$$(10.11) \quad r \|f(t)\| \leq \operatorname{Re} [f(t), e] \quad \text{for a.e. } t \in [a, b].$$

Then

$$(10.12) \quad r \int_a^b \|f(t)\| dt \leq \left\| \int_a^b f(t) dt \right\|$$

where equality holds in (10.12) if and only if both

$$(10.13) \quad \left[\int_a^b f(t) dt, e \right] = r \int_a^b \|f(t)\| dt$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 63 of 99

and

$$(10.14) \quad \left[\int_a^b f(t) dt, e \right] = \left\| \int_a^b f(t) dt \right\|.$$

The proof follows from Theorem 10.1 for the continuous linear functional $F(x) = [x, e]$, $x \in X$, and we omit the details.

The following corollary of Theorem 10.1 may be stated [8].

Corollary 10.3. *Let $(X, \|\cdot\|)$ be a strictly convex Banach space, $[\cdot, \cdot] : X \times X \rightarrow \mathbb{K}$ a semi-inner product generating the norm $\|\cdot\|$ and $e \in X$, $\|e\| = 1$. If $f : [a, b] \rightarrow X$ is Bochner integrable on $[a, b]$ and there exists a $r \geq 0$ such that (10.11) holds true, then (10.12) is valid. The case of equality holds in (10.12) if and only if*

$$(10.15) \quad \int_a^b f(t) dt = r \left(\int_a^b \|f(t)\| dt \right) e.$$


Proof. If (10.15) holds true, then, obviously

$$\left\| \int_a^b f(t) dt \right\| = r \left(\int_a^b \|f(t)\| dt \right) \|e\| = r \int_a^b \|f(t)\| dt,$$

which is the equality case in (10.12).

Conversely, if the equality holds in (10.12), then, by Corollary 10.2, we must have (10.13) and (10.14). Utilising Theorem 4.2, by (10.14) we can conclude that there exists a $\mu > 0$ such that

$$(10.16) \quad \int_a^b f(t) dt = \mu e.$$



Title Page

Contents



Go Back

Close

Quit

Page 64 of 99

Replacing this in (10.13), we get

$$\mu \|e\|^2 = r \int_a^b \|f(t)\| dt,$$

giving

$$(10.17) \quad \mu = r \int_a^b \|f(t)\| dt.$$

Utilising (10.16) and (10.17) we deduce (10.15) and the proof is completed. \square

10.2. The Case of m Functionals


The following result may be stated [8]:

Theorem 10.4 (Dragomir, 2004). *Let $(X, \|\cdot\|)$ be a Banach space over the real or complex number field \mathbb{K} and $F_k : X \rightarrow \mathbb{K}$, $k \in \{1, \dots, m\}$ continuous linear functionals on X . If $f : [a, b] \rightarrow X$ is a Bochner integrable function on $[a, b]$ and there exists $r_k \geq 0$, $k \in \{1, \dots, m\}$ with $\sum_{k=1}^m r_k > 0$ and*

$$(10.18) \quad r_k \|f(t)\| \leq \operatorname{Re} F_k[f(t)]$$

for each $k \in \{1, \dots, m\}$ and a.e. $t \in [a, b]$, then

$$(10.19) \quad \int_a^b \|f(t)\| dt \leq \frac{\|\sum_{k=1}^m F_k\|}{\sum_{k=1}^m r_k} \left\| \int_a^b f(t) dt \right\|.$$



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 65 of 99

The case of equality holds in (10.19) if both

$$(10.20) \quad \left(\sum_{k=1}^m F_k \right) \left(\int_a^b f(t) dt \right) = \left(\sum_{k=1}^m r_k \right) \int_a^b \|f(t)\| dt$$

and


$$(10.21) \quad \left(\sum_{k=1}^m F_k \right) \left(\int_a^b f(t) dt \right) = \left\| \sum_{k=1}^m F_k \right\| \left\| \int_a^b f(t) dt \right\|.$$

Proof. Utilising the hypothesis (10.18), we have

$$(10.22) \quad \begin{aligned} I &:= \left| \sum_{k=1}^m F_k \left(\int_a^b f(t) dt \right) \right| \geq \left| \operatorname{Re} \left[\sum_{k=1}^m F_k \left(\int_a^b f(t) dt \right) \right] \right| \\ &\geq \operatorname{Re} \left[\sum_{k=1}^m F_k \left(\int_a^b f(t) dt \right) \right] = \sum_{k=1}^m \left(\int_a^b \operatorname{Re} F_k f(t) dt \right) \\ &\geq \left(\sum_{k=1}^m r_k \right) \cdot \int_a^b \|f(t)\| dt. \end{aligned}$$

On the other hand, by the continuity property of F_k , $k \in \{1, \dots, m\}$, we obviously have

$$(10.23) \quad I = \left| \left(\sum_{k=1}^m F_k \right) \left(\int_a^b f(t) dt \right) \right| \leq \left\| \sum_{k=1}^m F_k \right\| \left\| \int_a^b f(t) dt \right\|.$$



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 66 of 99

Making use of (10.22) and (10.23), we deduce (10.19).

Now, obviously, if (10.20) and (10.21) are valid, then the case of equality holds true in (10.19).

Conversely, if the equality holds in the inequality (10.19), then it must hold in all the inequalities used to prove (10.19), therefore we have

$$(10.24) \quad r_k \|f(t)\| = \operatorname{Re} F_k[f(t)]$$

for each $k \in \{1, \dots, m\}$ and a.e. $t \in [a, b]$,

$$(10.25) \quad \operatorname{Im} \left(\sum_{k=1}^m F_k \right) \left(\int_a^b f(t) dt \right) = 0,$$


$$(10.26) \quad \operatorname{Re} \left(\sum_{k=1}^m F_k \right) \left(\int_a^b f(t) dt \right) = \left\| \sum_{k=1}^m F_k \right\| \left\| \int_a^b f(t) dt \right\|.$$

Note that, by (10.24), on integrating on $[a, b]$ and summing over $k \in \{1, \dots, m\}$, we get

$$(10.27) \quad \operatorname{Re} \left(\sum_{k=1}^m F_k \right) \left(\int_a^b f(t) dt \right) = \left(\sum_{k=1}^m r_k \right) \int_a^b \|f(t)\| dt.$$

Now, (10.25) and (10.27) imply (10.20) while (10.25) and (10.26) imply (10.21), therefore the theorem is proved. \square

The following new results may be stated as well:



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 67 of 99

Theorem 10.5. Let $(X, \|\cdot\|)$ be a Banach space over the real or complex number field \mathbb{K} and $F_k : X \rightarrow \mathbb{K}$, $k \in \{1, \dots, m\}$ continuous linear functionals on X . Also, assume that $f : [a, b] \rightarrow X$ is a Bochner integrable function on $[a, b]$ and there exists $r_k \geq 0$, $k \in \{1, \dots, m\}$ with $\sum_{k=1}^m r_k > 0$ and

$$r_k \|f(t)\| \leq \operatorname{Re} F_k [f(t)]$$

for each $k \in \{1, \dots, m\}$ and a.e. $t \in [a, b]$.

(i) If c_∞ is defined by (c_∞) , then we have the inequality

$$(10.28) \quad (1 \leq) \frac{\int_a^b \|f(t)\| dt}{\left\| \int_a^b f(t) dt \right\|} \leq \frac{c_\infty}{\max_{1 \leq k \leq m} \{r_k\}} \left(\leq \frac{\max_{1 \leq k \leq m} \|F_k\|}{\max_{1 \leq k \leq m} \{r_k\}} \right)$$

with equality if and only if


$$\operatorname{Re} (F_k) \left(\int_a^b f(t) dt \right) = r_k \int_a^b \|f(t)\| dt$$

for each $k \in \{1, \dots, m\}$ and

$$\max_{1 \leq k \leq m} \left[\operatorname{Re} (F_k) \left(\int_a^b f(t) dt \right) \right] = c_\infty \int_a^b \|f(t)\| dt.$$

(ii) If c_p , $p \geq 1$, is defined by (c_p) , then we have the inequality

$$(1 \leq) \frac{\int_a^b \|f(t)\| dt}{\left\| \int_a^b f(t) dt \right\|} \leq \frac{c_p}{\left(\sum_{k=1}^m r_k^p \right)^{\frac{1}{p}}} \left(\leq \frac{\sum_{k=1}^m \|F_k\|^p}{\sum_{k=1}^m r_k^p} \right)^{\frac{1}{p}}$$



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 68 of 99

with equality if and only if

$$\operatorname{Re} (F_k) \left(\int_a^b f(t) dt \right) = r_k \int_a^b \|f(t)\| dt$$

for each $k \in \{1, \dots, m\}$ and

$$\sum_{k=1}^m \left[\operatorname{Re} F_k \left(\int_a^b f(t) dt \right) \right]^p = c_p^p \left\| \int_a^b f(t) dt \right\|^p$$

where $p \geq 1$.

The proof is similar to the ones from Theorems 5.1, 5.2 and 10.4 and we omit the details.


The case of Hilbert spaces for Theorem 10.4, which provides a simpler condition for equality, is of interest for applications [8].

Theorem 10.6 (Dragomir, 2004). *Let $(X, \|\cdot\|)$ be a Hilbert space over the real or complex number field \mathbb{K} and $e_k \in H \setminus \{0\}$, $k \in \{1, \dots, m\}$. If $f : [a, b] \rightarrow H$ is a Bochner integrable function and $r_k \geq 0$, $k \in \{1, \dots, m\}$ and $\sum_{k=1}^m r_k > 0$ satisfy*

$$(10.29) \quad r_k \|f(t)\| \leq \operatorname{Re} \langle f(t), e_k \rangle$$

for each $k \in \{1, \dots, m\}$ and for a.e. $t \in [a, b]$, then

$$(10.30) \quad \int_a^b \|f(t)\| dt \leq \frac{\left\| \sum_{k=1}^m e_k \right\|}{\sum_{k=1}^m r_k} \left\| \int_a^b f(t) dt \right\|.$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 69 of 99

The case of equality holds in (10.30) for $f \neq 0$ a.e. on $[a, b]$ if and only if

$$(10.31) \quad \int_a^b f(t) dt = \frac{(\sum_{k=1}^m r_k) \int_a^b \|f(t)\| dt}{\|\sum_{k=1}^m e_k\|^2} \sum_{k=1}^m e_k.$$

Proof. Utilising the hypothesis (10.29) and the modulus properties, we have


$$(10.32) \quad \left| \left\langle \int_a^b f(t) dt, \sum_{k=1}^m e_k \right\rangle \right| \\ \geq \left| \sum_{k=1}^m \operatorname{Re} \left\langle \int_a^b f(t) dt, e_k \right\rangle \right| \geq \sum_{k=1}^m \operatorname{Re} \left\langle \int_a^b f(t) dt, e_k \right\rangle \\ = \sum_{k=1}^m \int_a^b \operatorname{Re} \langle f(t), e_k \rangle dt \geq \left(\sum_{k=1}^m r_k \right) \int_a^b \|f(t)\| dt.$$

By Schwarz's inequality in Hilbert spaces applied for $\int_a^b f(t) dt$ and $\sum_{k=1}^m e_k$, we have

$$(10.33) \quad \left\| \int_a^b f(t) dt \right\| \left\| \sum_{k=1}^m e_k \right\| \geq \left| \left\langle \int_a^b f(t) dt, \sum_{k=1}^m e_k \right\rangle \right|.$$

Making use of (10.32) and (10.33), we deduce (10.30).

Now, if $f \neq 0$ a.e. on $[a, b]$, then $\int_a^b \|f(t)\| dt \neq 0$ and by (10.32) $\sum_{k=1}^m e_k \neq$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 70 of 99

0. Obviously, if (10.31) is valid, then taking the norm we have

$$\begin{aligned} \left\| \int_a^b f(t) dt \right\| &= \frac{(\sum_{k=1}^m r_k) \int_a^b \|f(t)\| dt}{\left\| \sum_{k=1}^m e_k \right\|^2} \left\| \sum_{k=1}^m e_k \right\| \\ &= \frac{\sum_{k=1}^m r_k}{\left\| \sum_{k=1}^m e_k \right\|} \int_a^b \|f(t)\| dt, \end{aligned}$$

i.e., the case of equality holds true in (10.30).

Conversely, if the equality case holds true in (10.30), then it must hold in all the inequalities used to prove (10.30), therefore we have

$$(10.34) \quad \operatorname{Re} \langle f(t), e_k \rangle = r_k \|f(t)\|$$

for each $k \in \{1, \dots, m\}$ and a.e. $t \in [a, b]$,


$$(10.35) \quad \left\| \int_a^b f(t) dt \right\| \left\| \sum_{k=1}^m e_k \right\| = \left| \left\langle \int_a^b f(t) dt, \sum_{k=1}^m e_k \right\rangle \right|,$$

and

$$(10.36) \quad \operatorname{Im} \left\langle \int_a^b f(t) dt, \sum_{k=1}^m e_k \right\rangle = 0.$$

From (10.34) on integrating on $[a, b]$ and summing over k from 1 to m , we get

$$(10.37) \quad \operatorname{Re} \left\langle \int_a^b f(t) dt, \sum_{k=1}^m e_k \right\rangle = \left(\sum_{k=1}^m r_k \right) \int_a^b \|f(t)\| dt,$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 71 of 99

and then, by (10.36) and (10.37), we have

$$(10.38) \quad \left\langle \int_a^b f(t) dt, \sum_{k=1}^m e_k \right\rangle = \left(\sum_{k=1}^m r_k \right) \int_a^b \|f(t)\| dt.$$

On the other hand, by the use of the identity (3.22), the relation (10.35) holds true if and only if

$$(10.39) \quad \int_a^b f(t) dt = \frac{\left\langle \int_a^b f(t) dt, \sum_{k=1}^m e_k \right\rangle}{\left\| \sum_{k=1}^m e_k \right\|} \sum_{k=1}^m e_k.$$

Finally, by (10.38) and (10.39) we deduce that (10.31) is also necessary for the equality case in (10.30) and the theorem is proved. \square

Remark 9. If $\{e_k\}_{k \in \{1, \dots, m\}}$ are orthogonal, then (10.30) can be replaced by


$$(10.40) \quad \int_a^b \|f(t)\| dt \leq \frac{\left(\sum_{k=1}^m \|e_k\|^2 \right)^{\frac{1}{2}}}{\sum_{k=1}^m r_k} \left\| \int_a^b f(t) dt \right\|,$$

with equality if and only if

$$(10.41) \quad \int_a^b f(t) dt = \frac{\left(\sum_{k=1}^m r_k \right) \int_a^b \|f(t)\| dt}{\sum_{k=1}^m \|e_k\|^2} \sum_{k=1}^m e_k.$$

Moreover, if $\{e_k\}_{k \in \{1, \dots, m\}}$ are orthonormal, then (10.40) becomes

$$(10.42) \quad \int_a^b \|f(t)\| dt \leq \frac{\sqrt{m}}{\sum_{k=1}^m r_k} \left\| \int_a^b f(t) dt \right\|,$$



with equality if and only if

$$(10.43) \quad \int_a^b f(t) dt = \frac{1}{m} \left(\sum_{k=1}^m r_k \right) \left(\int_a^b \|f(t)\| dt \right) \sum_{k=1}^m e_k.$$

The following corollary of Theorem 10.6 may be stated as well [8].

Corollary 10.7. *Let $(H; \langle \cdot, \cdot \rangle)$ be a Hilbert space over the real or complex number field \mathbb{K} and $e_k \in H \setminus \{0\}$, $k \in \{1, \dots, m\}$. If $f : [a, b] \rightarrow H$ is a Bochner integrable function on $[a, b]$ and $\rho_k > 0$, $k \in \{1, \dots, m\}$ with*

$$(10.44) \quad \|f(t) - e_k\| \leq \rho_k < \|e_k\|$$

for each $k \in \{1, \dots, m\}$ and a.e. $t \in [a, b]$, then


$$(10.45) \quad \int_a^b \|f(t)\| dt \leq \frac{\|\sum_{k=1}^m e_k\|}{\sum_{k=1}^m (\|e_k\|^2 - \rho_k^2)^{\frac{1}{2}}} \left\| \int_a^b f(t) dt \right\|.$$

The case of equality holds in (10.45) if and only if

$$(10.46) \quad \int_a^b f(t) dt = \frac{\sum_{k=1}^m (\|e_k\|^2 - \rho_k^2)^{\frac{1}{2}}}{\|\sum_{k=1}^m e_k\|^2} \left(\int_a^b \|f(t)\| dt \right) \sum_{k=1}^m e_k.$$

Proof. Utilising Lemma 3.3, we have from (10.44) that

$$\|f(t)\| (\|e_k\|^2 - \rho_k^2)^{\frac{1}{2}} \leq \operatorname{Re} \langle f(t), e_k \rangle$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 73 of 99

for any $k \in \{1, \dots, m\}$ and a.e. $t \in [a, b]$.

Applying Theorem 10.6 for

$$r_k := (\|e_k\|^2 - \rho_k^2)^{\frac{1}{2}}, \quad k \in \{1, \dots, m\},$$

we deduce the desired result. □

Remark 10. If $\{e_k\}_{k \in \{1, \dots, m\}}$ are orthogonal, then (10.45) becomes

$$(10.47) \quad \int_a^b \|f(t)\| dt \leq \frac{(\sum_{k=1}^m \|e_k\|^2)^{\frac{1}{2}}}{\sum_{k=1}^m (\|e_k\|^2 - \rho_k^2)^{\frac{1}{2}}} \left\| \int_a^b f(t) dt \right\|,$$

with equality if and only if

$$(10.48) \quad \int_a^b f(t) dt = \frac{\sum_{k=1}^m (\|e_k\|^2 - \rho_k^2)^{\frac{1}{2}}}{\sum_{k=1}^m \|e_k\|^2} \left(\int_a^b \|f(t)\| dt \right) \sum_{k=1}^m e_k.$$

Moreover, if $\{e_k\}_{k \in \{1, \dots, m\}}$ is assumed to be orthonormal and


$$\|f(t) - e_k\| \leq \rho_k \quad \text{for a.e. } t \in [a, b],$$

where $\rho_k \in [0, 1)$, $k \in \{1, \dots, m\}$, then

$$(10.49) \quad \int_a^b \|f(t)\| dt \leq \frac{\sqrt{m}}{\sum_{k=1}^m (1 - \rho_k^2)^{\frac{1}{2}}} \left\| \int_a^b f(t) dt \right\|,$$

with equality iff

$$(10.50) \quad \int_a^b f(t) dt = \frac{\sum_{k=1}^m (1 - \rho_k^2)^{\frac{1}{2}}}{m} \left(\int_a^b \|f(t)\| dt \right) \sum_{k=1}^m e_k.$$



Finally, we may state the following corollary of Theorem 10.6 [11].

Corollary 10.8. *Let $(H; \langle \cdot, \cdot \rangle)$ be a Hilbert space over the real or complex number field \mathbb{K} and $e_k \in H \setminus \{0\}$, $k \in \{1, \dots, m\}$. If $f : [a, b] \rightarrow H$ is a Bochner integrable function on $[a, b]$ and $M_k \geq \mu_k > 0$, $k \in \{1, \dots, m\}$ are such that either*

$$(10.51) \quad \operatorname{Re} \langle M_k e_k - f(t), f(t) - \mu_k e_k \rangle \geq 0$$

or, equivalently,

$$(10.52) \quad \left\| f(t) - \frac{M_k + \mu_k}{2} e_k \right\| \leq \frac{1}{2} (M_k - \mu_k) \|e_k\|$$

for each $k \in \{1, \dots, m\}$ and a.e. $t \in [a, b]$, then


$$(10.53) \quad \int_a^b \|f(t)\| dt \leq \frac{\|\sum_{k=1}^m e_k\|}{\sum_{k=1}^m \frac{2 \cdot \sqrt{\mu_k M_k}}{\mu_k + M_k} \|e_k\|} \left\| \int_a^b f(t) dt \right\|.$$

The case of equality holds if and only if

$$\int_a^b f(t) dt = \frac{\sum_{k=1}^m \frac{2 \cdot \sqrt{\mu_k M_k}}{\mu_k + M_k} \|e_k\|}{\|\sum_{k=1}^m e_k\|^2} \left(\int_a^b \|f(t)\| dt \right) \cdot \sum_{k=1}^m e_k.$$

Proof. Utilising Lemma 3.5, by (10.51) we deduce

$$\|f(t)\| \frac{2 \cdot \sqrt{\mu_k M_k}}{\mu_k + M_k} \|e_k\| \leq \operatorname{Re} \langle f(t), e_k \rangle$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit


Page 75 of 99

for each $k \in \{1, \dots, m\}$ and a.e. $t \in [a, b]$.

Applying Theorem 10.6 for

$$r_k := \frac{2 \cdot \sqrt{\mu_k M_k}}{\mu_k + M_k} \|e_k\|, \quad k \in \{1, \dots, m\}$$

we deduce the desired result. □



**Reverses of the Triangle
Inequality in Banach Spaces**

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 76 of 99

11. Additive Reverses of the Continuous Triangle Inequality

11.1. The Case of One Functional

The aim of this section is to provide a different approach to the problem of reversing the continuous triangle inequality. Namely, we are interested in finding upper bounds for the positive difference

$$\int_a^b \|f(t)\| dt - \left\| \int_a^b f(t) dt \right\|$$

under various assumptions for the Bochner integrable function $f : [a, b] \rightarrow X$.


In the following we provide an additive reverse for the continuous triangle inequality that has been established in [8].

Theorem 11.1 (Dragomir, 2004). *Let $(X, \|\cdot\|)$ be a Banach space over the real or complex number field \mathbb{K} and $F : X \rightarrow \mathbb{K}$ be a continuous linear functional of unit norm on X . Suppose that the function $f : [a, b] \rightarrow X$ is Bochner integrable on $[a, b]$ and there exists a Lebesgue integrable function $k : [a, b] \rightarrow [0, \infty)$ such that*

$$(11.1) \quad \|f(t)\| - \operatorname{Re} F[f(t)] \leq k(t)$$

for a.e. $t \in [a, b]$. Then we have the inequality

$$(11.2) \quad (0 \leq) \int_a^b \|f(t)\| dt - \left\| \int_a^b f(t) dt \right\| \leq \int_a^b k(t) dt.$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 77 of 99

The equality holds in (11.2) if and only if both

$$(11.3) \quad F \left(\int_a^b f(t) dt \right) = \left\| \int_a^b f(t) dt \right\|$$

and

$$(11.4) \quad F \left(\int_a^b f(t) dt \right) = \int_a^b \|f(t)\| dt - \int_a^b k(t) dt.$$

Proof. Since the norm of F is unity, then

$$|F(x)| \leq \|x\| \quad \text{for any } x \in X.$$


Applying this inequality for the vector $\int_a^b f(t) dt$, we get

$$(11.5) \quad \left\| \int_a^b f(t) dt \right\| \geq \left| F \left(\int_a^b f(t) dt \right) \right| \geq \left| \operatorname{Re} F \left(\int_a^b f(t) dt \right) \right| \\ = \left| \int_a^b \operatorname{Re} F[f(t)] dt \right| \geq \int_a^b \operatorname{Re} F[f(t)] dt.$$

Integrating (11.1), we have

$$(11.6) \quad \int_a^b \|f(t)\| dt - \operatorname{Re} F \left(\int_a^b f(t) dt \right) \leq \int_a^b k(t) dt.$$

Now, making use of (11.5) and (11.6), we deduce (11.2).



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 78 of 99

Obviously, if the equality hold in (11.3) and (11.4), then it holds in (11.2) as well. Conversely, if the equality holds in (11.2), then it must hold in all the inequalities used to prove (11.2). Therefore, we have

$$\int_a^b \|f(t)\| dt = \operatorname{Re} \left[F \left(\int_a^b f(t) dt \right) \right] + \int_a^b k(t) dt.$$

and

$$\operatorname{Re} \left[F \left(\int_a^b f(t) dt \right) \right] = \left| F \left(\int_a^b f(t) dt \right) \right| = \left\| \int_a^b f(t) dt \right\|$$

which imply (11.3) and (11.4). □

Corollary 11.2. *Let $(X, \|\cdot\|)$ be a Banach space, $[\cdot, \cdot] : X \times X \rightarrow \mathbb{K}$ a semi-inner product which generates its norm. If $e \in X$ is such that $\|e\| = 1$, $f : [a, b] \rightarrow X$ is Bochner integrable on $[a, b]$ and there exists a Lebesgue integrable function $k : [a, b] \rightarrow [0, \infty)$ such that*

$$(11.7) \quad (0 \leq) \|f(t)\| - \operatorname{Re} [f(t), e] \leq k(t),$$

for a.e. $t \in [a, b]$, then

$$(11.8) \quad (0 \leq) \int_a^b \|f(t)\| dt - \left\| \int_a^b f(t) dt \right\| \leq \int_a^b k(t) dt,$$

where equality holds in (11.8) if and only if both

$$(11.9) \quad \left[\int_a^b f(t) dt, e \right] = \left\| \int_a^b f(t) dt \right\|$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 79 of 99

and

$$(11.10) \quad \left[\int_a^b f(t) dt, e \right] = \left\| \int_a^b f(t) dt \right\| - \int_a^b k(t) dt.$$

The proof is obvious by Theorem 11.1 applied for the continuous linear functional of unit norm $F_e : X \rightarrow \mathbb{K}$, $F_e(x) = [x, e]$.

The following corollary may be stated.

Corollary 11.3. *Let $(X, \|\cdot\|)$ be a strictly convex Banach space, and $[\cdot, \cdot]$, e , f , k as in Corollary 11.2. Then the case of equality holds in (11.8) if and only if*

$$(11.11) \quad \int_a^b \|f(t)\| dt \geq \int_a^b k(t) dt$$

and

$$(11.12) \quad \int_a^b f(t) dt = \left(\int_a^b \|f(t)\| dt - \int_a^b k(t) dt \right) e.$$


Proof. Suppose that (11.11) and (11.12) are valid. Taking the norm on (11.12) we have

$$\left\| \int_a^b f(t) dt \right\| = \left| \int_a^b \|f(t)\| dt - \int_a^b k(t) dt \right| \|e\| = \int_a^b \|f(t)\| dt - \int_a^b k(t) dt,$$

and the case of equality holds true in (11.8).

Now, if the equality case holds in (11.8), then obviously (11.11) is valid, and by Corollary 11.2,

$$\left[\int_a^b f(t) dt, e \right] = \left\| \int_a^b f(t) dt \right\| \|e\|.$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 80 of 99

Utilising Theorem 4.2, we get

$$(11.13) \quad \int_a^b f(t) dt = \lambda e \text{ with } \lambda > 0.$$

Replacing $\int_a^b f(t) dt$ with λe in the second equation of (11.9) we deduce

$$(11.14) \quad \lambda = \int_a^b \|f(t)\| dt - \int_a^b k(t) dt,$$

and by (11.13) and (11.14) we deduce (11.12). □


Remark 11. *If $X = H$, $(H; \langle \cdot, \cdot \rangle)$ is a Hilbert space, then from Corollary 11.3 we deduce the additive reverse inequality obtained in [7]. For further similar results in Hilbert spaces, see [7] and [9].*

11.2. The Case of m Functionals

The following result may be stated [8]:

Theorem 11.4 (Dragomir, 2004). *Let $(X, \|\cdot\|)$ be a Banach space over the real or complex number field \mathbb{K} and $F_k : X \rightarrow \mathbb{K}$, $k \in \{1, \dots, m\}$ continuous linear functionals on X . If $f : [a, b] \rightarrow X$ is a Bochner integrable function on $[a, b]$ and $M_k : [a, b] \rightarrow [0, \infty)$, $k \in \{1, \dots, m\}$ are Lebesgue integrable functions such that*

$$(11.15) \quad \|f(t)\| - \operatorname{Re} F_k[f(t)] \leq M_k(t)$$



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 81 of 99

for each $k \in \{1, \dots, m\}$ and a.e. $t \in [a, b]$, then

$$(11.16) \quad \int_a^b \|f(t)\| dt \leq \left\| \frac{1}{m} \sum_{k=1}^m F_k \right\| \left\| \int_a^b f(t) dt \right\| + \frac{1}{m} \sum_{k=1}^m \int_a^b M_k(t) dt.$$

The case of equality holds in (11.16) if and only if both

$$(11.17) \quad \frac{1}{m} \sum_{k=1}^m F_k \left(\int_a^b f(t) dt \right) = \left\| \frac{1}{m} \sum_{k=1}^m F_k \right\| \left\| \int_a^b f(t) dt \right\|$$


and

$$(11.18) \quad \frac{1}{m} \sum_{k=1}^m F_k \left(\int_a^b f(t) dt \right) = \int_a^b \|f(t)\| dt - \frac{1}{m} \sum_{k=1}^m \int_a^b M_k(t) dt.$$

Proof. If we integrate on $[a, b]$ and sum over k from 1 to m , we deduce

$$(11.19) \quad \int_a^b \|f(t)\| dt \leq \frac{1}{m} \sum_{k=1}^m \operatorname{Re} \left[F_k \left(\int_a^b f(t) dt \right) \right] + \frac{1}{m} \sum_{k=1}^m \int_a^b M_k(t) dt.$$

Utilising the continuity property of the functionals F_k and the properties of the



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 82 of 99

modulus, we have:

$$\begin{aligned}
 (11.20) \quad \sum_{k=1}^m \operatorname{Re} F_k \left(\int_a^b f(t) dt \right) &\leq \left| \sum_{k=1}^m \operatorname{Re} \left[F_k \left(\int_a^b f(t) dt \right) \right] \right| \\
 &\leq \left| \sum_{k=1}^m F_k \left(\int_a^b f(t) dt \right) \right| \\
 &\leq \left\| \sum_{k=1}^m F_k \right\| \left\| \int_a^b f(t) dt \right\|.
 \end{aligned}$$

Now, by (11.19) and (11.20) we deduce (11.16).

Obviously, if (11.17) and (11.18) hold true, then the case of equality is valid in (11.16).

Conversely, if the case of equality holds in (11.16), then it must hold in all the inequalities used to prove (11.16). Therefore, we have


$$\int_a^b \|f(t)\| dt = \frac{1}{m} \sum_{k=1}^m \operatorname{Re} \left[F_k \left(\int_a^b f(t) dt \right) \right] + \frac{1}{m} \sum_{k=1}^m \int_a^b M_k(t) dt,$$

$$\sum_{k=1}^m \operatorname{Re} \left[F_k \left(\int_a^b f(t) dt \right) \right] = \left\| \int_a^b f(t) dt \right\| \left\| \sum_{k=1}^m F_k \right\|$$

and

$$\sum_{k=1}^m \operatorname{Im} \left[F_k \left(\int_a^b f(t) dt \right) \right] = 0.$$

These imply that (11.17) and (11.18) hold true, and the theorem is completely proved. \square



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 83 of 99

Remark 12. If $F_k, k \in \{1, \dots, m\}$ are of unit norm, then, from (11.16) we deduce the inequality

$$(11.21) \quad \int_a^b \|f(t)\| dt \leq \left\| \int_a^b f(t) dt \right\| + \frac{1}{m} \sum_{k=1}^m \int_a^b M_k(t) dt,$$

which is obviously coarser than (11.16) but, perhaps more useful for applications.

The following new result may be stated as well:

Theorem 11.5. Let $(X, \|\cdot\|)$ be a Banach space over the real or complex number field \mathbb{K} and $F_k : X \rightarrow \mathbb{K}, k \in \{1, \dots, m\}$ continuous linear functionals on X . Assume also that $f : [a, b] \rightarrow X$ is a Bochner integrable function on $[a, b]$ and $M_k : [a, b] \rightarrow [0, \infty), k \in \{1, \dots, m\}$ are Lebesgue integrable functions such that

$$(11.22) \quad \|f(t)\| - \operatorname{Re} F_k[f(t)] \leq M_k(t)$$

for each $k \in \{1, \dots, m\}$ and a.e. $t \in [a, b]$.

(i) If c_∞ is defined by (c_∞) , then we have the inequality

$$(11.23) \quad \int_a^b \|f(t)\| dt \leq c_\infty \left\| \int_a^b f(t) dt \right\| + \frac{1}{m} \sum_{k=1}^m \int_a^b M_k(t) dt.$$

(ii) If $c_p, p \geq 1$, is defined by (c_p) , then we have the inequality

$$\int_a^b \|f(t)\| dt \leq \frac{c_p}{m^{1/p}} \left\| \int_a^b f(t) dt \right\| + \frac{1}{m} \sum_{k=1}^m \int_a^b M_k(t) dt.$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 84 of 99

The proof is similar to the ones from Theorem 7.1 and 11.4 and we omit the details.

The case of Hilbert spaces for Theorem 11.4, in which one may provide a simpler condition for equality, is of interest in applications [8].

Theorem 11.6 (Dragomir, 2004). *Let $(H, \langle \cdot, \cdot \rangle)$ be a Hilbert space over the real or complex number field \mathbb{K} and $e_k \in H, k \in \{1, \dots, m\}$. If $f : [a, b] \rightarrow H$ is a Bochner integrable function on $[a, b]$, $f(t) \neq 0$ for a.e. $t \in [a, b]$ and $M_k : [a, b] \rightarrow [0, \infty), k \in \{1, \dots, m\}$ is a Lebesgue integrable function such that*

$$(11.24) \quad \|f(t)\| - \operatorname{Re} \langle f(t), e_k \rangle \leq M_k(t)$$

for each $k \in \{1, \dots, m\}$ and for a.e. $t \in [a, b]$, then


$$(11.25) \quad \int_a^b \|f(t)\| dt \leq \left\| \frac{1}{m} \sum_{k=1}^m e_k \right\| \left\| \int_a^b f(t) dt \right\| + \frac{1}{m} \sum_{k=1}^m \int_a^b M_k(t) dt.$$

The case of equality holds in (11.25) if and only if

$$(11.26) \quad \int_a^b \|f(t)\| dt \geq \frac{1}{m} \sum_{k=1}^m \int_a^b M_k(t) dt$$

and

$$(11.27) \quad \int_a^b f(t) dt = \frac{m \left(\int_a^b \|f(t)\| dt - \frac{1}{m} \sum_{k=1}^m \int_a^b M_k(t) dt \right)}{\left\| \sum_{k=1}^m e_k \right\|^2} \sum_{k=1}^m e_k.$$



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 85 of 99

Proof. As in the proof of Theorem 11.4, we have

$$(11.28) \quad \int_a^b \|f(t)\| dt \leq \operatorname{Re} \left\langle \frac{1}{m} \sum_{k=1}^m e_k, \int_a^b f(t) dt \right\rangle + \frac{1}{m} \sum_{k=1}^m \int_a^b M_k(t) dt$$

and $\sum_{k=1}^m e_k \neq 0$.

On utilising Schwarz's inequality in Hilbert space $(H, \langle \cdot, \cdot \rangle)$ for $\int_a^b f(t) dt$ and $\sum_{k=1}^m e_k$, we have


$$(11.29) \quad \begin{aligned} \left\| \int_a^b f(t) dt \right\| \left\| \sum_{k=1}^m e_k \right\| &\geq \left| \left\langle \int_a^b f(t) dt, \sum_{k=1}^m e_k \right\rangle \right| \\ &\geq \left| \operatorname{Re} \left\langle \int_a^b f(t) dt, \sum_{k=1}^m e_k \right\rangle \right| \\ &\geq \operatorname{Re} \left\langle \int_a^b f(t) dt, \sum_{k=1}^m e_k \right\rangle. \end{aligned}$$

By (11.28) and (11.29), we deduce (11.25).

Taking the norm on (11.27) and using (11.26), we have

$$\left\| \int_a^b f(t) dt \right\| = \frac{m \left(\int_a^b \|f(t)\| dt - \frac{1}{m} \sum_{k=1}^m \int_a^b M_k(t) dt \right)}{\left\| \sum_{k=1}^m e_k \right\|},$$

showing that the equality holds in (11.25).



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 86 of 99

Conversely, if the equality case holds in (11.25), then it must hold in all the inequalities used to prove (11.25). Therefore we have

$$(11.30) \quad \|f(t)\| = \operatorname{Re} \langle f(t), e_k \rangle + M_k(t)$$

for each $k \in \{1, \dots, m\}$ and for a.e. $t \in [a, b]$,

$$(11.31) \quad \left\| \int_a^b f(t) dt \right\| \left\| \sum_{k=1}^m e_k \right\| = \left| \left\langle \int_a^b f(t) dt, \sum_{k=1}^m e_k \right\rangle \right|$$

and

$$(11.32) \quad \operatorname{Im} \left\langle \int_a^b f(t) dt, \sum_{k=1}^m e_k \right\rangle = 0.$$

From (11.30) on integrating on $[a, b]$ and summing over k , we get


$$(11.33) \quad \operatorname{Re} \left\langle \int_a^b f(t) dt, \sum_{k=1}^m e_k \right\rangle = m \int_a^b \|f(t)\| dt - \sum_{k=1}^m \int_a^b M_k(t) dt.$$

On the other hand, by the use of the identity (3.22), the relation (11.31) holds if and only if

$$\int_a^b f(t) dt = \frac{\left\langle \int_a^b f(t) dt, \sum_{k=1}^m e_k \right\rangle}{\left\| \sum_{k=1}^m e_k \right\|^2} \sum_{k=1}^m e_k,$$

giving, from (11.32) and (11.33), that (11.27) holds true.

If the equality holds in (11.25), then obviously (11.26) is valid and the theorem is proved. \square



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 87 of 99

Remark 13. If in the above theorem, the vectors $\{e_k\}_{k \in \{1, \dots, m\}}$ are assumed to be orthogonal, then (11.25) becomes

$$(11.34) \quad \int_a^b \|f(t)\| dt \leq \frac{1}{m} \left(\sum_{k=1}^m \|e_k\|^2 \right)^{\frac{1}{2}} \left\| \int_a^b f(t) dt \right\| + \frac{1}{m} \sum_{k=1}^m \int_a^b M_k(t) dt.$$

Moreover, if $\{e_k\}_{k \in \{1, \dots, m\}}$ is an orthonormal family, then (11.34) becomes

$$(11.35) \quad \int_a^b \|f(t)\| dt \leq \frac{1}{\sqrt{m}} \left\| \int_a^b f(t) dt \right\| + \frac{1}{m} \sum_{k=1}^m \int_a^b M_k(t) dt$$

which has been obtained in [4].


The following corollaries are of interest.

Corollary 11.7. Let $(H; \langle \cdot, \cdot \rangle)$, e_k , $k \in \{1, \dots, m\}$ and f be as in Theorem 11.6. If $r_k : [a, b] \rightarrow [0, \infty)$, $k \in \{1, \dots, m\}$ are such that $r_k \in L^2[a, b]$, $k \in \{1, \dots, m\}$ and

$$(11.36) \quad \|f(t) - e_k\| \leq r_k(t),$$

for each $k \in \{1, \dots, m\}$ and a.e. $t \in [a, b]$, then

$$(11.37) \quad \int_a^b \|f(t)\| dt \leq \left\| \frac{1}{m} \sum_{k=1}^m e_k \right\| \left\| \int_a^b f(t) dt \right\| + \frac{1}{2m} \sum_{k=1}^m \int_a^b r_k^2(t) dt.$$



Title Page

Contents



Go Back

Close

Quit

Page 88 of 99

The case of equality holds in (11.37) if and only if

$$\int_a^b \|f(t)\| dt \geq \frac{1}{2m} \sum_{k=1}^m \int_a^b r_k^2(t) dt$$

and

$$\int_a^b f(t) dt = \frac{m \left(\int_a^b \|f(t)\| dt - \frac{1}{2m} \sum_{k=1}^m \int_a^b r_k^2(t) dt \right)}{\left\| \sum_{k=1}^m e_k \right\|^2} \sum_{k=1}^m e_k.$$


Finally, the following corollary may be stated.

Corollary 11.8. Let $(H; \langle \cdot, \cdot \rangle)$, e_k , $k \in \{1, \dots, m\}$ and f be as in Theorem 11.6. If $M_k, \mu_k : [a, b] \rightarrow \mathbb{R}$ are such that $M_k \geq \mu_k > 0$ a.e. on $[a, b]$, $\frac{(M_k - \mu_k)^2}{M_k + \mu_k} \in L[a, b]$ and

$$\operatorname{Re} \langle M_k(t) e_k - f(t), f(t) - \mu_k(t) e_k \rangle \geq 0$$

for each $k \in \{1, \dots, m\}$ and for a.e. $t \in [a, b]$, then

$$\begin{aligned} \int_a^b \|f(t)\| dt &\leq \left\| \frac{1}{m} \sum_{k=1}^m e_k \right\| \left\| \int_a^b f(t) dt \right\| \\ &\quad + \frac{1}{4m} \sum_{k=1}^m \|e_k\|^2 \int_a^b \frac{[M_k(t) - \mu_k(t)]^2}{M_k(t) + \mu_k(t)} dt. \end{aligned}$$



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 89 of 99

12. Applications for Complex-Valued Functions

We now give some examples of inequalities for complex-valued functions that are Lebesgue integrable on using the general result obtained in Section 10.

Consider the Banach space $(\mathbb{C}, |\cdot|_1)$ over the real field \mathbb{R} and $F : \mathbb{C} \rightarrow \mathbb{C}$, $F(z) = ez$ with $e = \alpha + i\beta$ and $|e|^2 = \alpha^2 + \beta^2 = 1$, then F is linear on \mathbb{C} . For $z \neq 0$, we have

$$\frac{|F(z)|}{|z|_1} = \frac{|e||z|}{|z|_1} = \frac{\sqrt{|\operatorname{Re} z|^2 + |\operatorname{Im} z|^2}}{|\operatorname{Re} z| + |\operatorname{Im} z|} \leq 1.$$

Since, for $z_0 = 1$, we have $|F(z_0)| = 1$ and $|z_0|_1 = 1$, hence

$$\|F\|_1 := \sup_{z \neq 0} \frac{|F(z)|}{|z|_1} = 1,$$

showing that F is a bounded linear functional on $(\mathbb{C}, |\cdot|_1)$.


Therefore we can apply Theorem 10.1 to state the following result for complex-valued functions.

Proposition 12.1. *Let $\alpha, \beta \in \mathbb{R}$ with $\alpha^2 + \beta^2 = 1$, $f : [a, b] \rightarrow \mathbb{C}$ be a Lebesgue integrable function on $[a, b]$ and $r \geq 0$ such that*

$$(12.1) \quad r [|\operatorname{Re} f(t)| + |\operatorname{Im} f(t)|] \leq \alpha \operatorname{Re} f(t) - \beta \operatorname{Im} f(t)$$

for a.e. $t \in [a, b]$. Then

$$(12.2) \quad r \left[\int_a^b |\operatorname{Re} f(t)| dt + \int_a^b |\operatorname{Im} f(t)| dt \right]$$



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 90 of 99

$$\leq \left| \int_a^b \operatorname{Re} f(t) dt \right| + \left| \int_a^b \operatorname{Im} f(t) dt \right|.$$

The equality holds in (12.2) if and only if both

$$\alpha \int_a^b \operatorname{Re} f(t) dt - \beta \int_a^b \operatorname{Im} f(t) dt = r \left[\int_a^b |\operatorname{Re} f(t)| dt + \int_a^b |\operatorname{Im} f(t)| dt \right]$$

and

$$\alpha \int_a^b \operatorname{Re} f(t) dt - \beta \int_a^b \operatorname{Im} f(t) dt = \left| \int_a^b \operatorname{Re} f(t) dt \right| + \left| \int_a^b \operatorname{Im} f(t) dt \right|.$$

Now, consider the Banach space $(\mathbb{C}, |\cdot|_\infty)$. If $F(z) = dz$ with $d = \gamma + i\delta$ and $|d| = \frac{\sqrt{2}}{2}$, i.e., $\gamma^2 + \delta^2 = \frac{1}{2}$, then F is linear on \mathbb{C} . For $z \neq 0$ we have


$$\frac{|F(z)|}{|z|_\infty} = \frac{|d||z|}{|z|_\infty} = \frac{\sqrt{2}}{2} \cdot \frac{\sqrt{|\operatorname{Re} z|^2 + |\operatorname{Im} z|^2}}{\max\{|\operatorname{Re} z|, |\operatorname{Im} z|\}} \leq 1.$$

Since, for $z_0 = 1 + i$, we have $|F(z_0)| = 1$, $|z_0|_\infty = 1$, hence

$$\|F\|_\infty := \sup_{z \neq 0} \frac{|F(z)|}{|z|_\infty} = 1,$$

showing that F is a bounded linear functional of unit norm on $(\mathbb{C}, |\cdot|_\infty)$.

Therefore, we can apply Theorem 10.1, to state the following result for complex-valued functions.



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 91 of 99

Proposition 12.2. Let $\gamma, \delta \in \mathbb{R}$ with $\gamma^2 + \delta^2 = \frac{1}{2}$, $f : [a, b] \rightarrow \mathbb{C}$ be a Lebesgue integrable function on $[a, b]$ and $r \geq 0$ such that

$$r \max \{ |\operatorname{Re} f(t)|, |\operatorname{Im} f(t)| \} \leq \gamma \operatorname{Re} f(t) - \delta \operatorname{Im} f(t)$$

for a.e. $t \in [a, b]$. Then

$$(12.3) \quad r \int_a^b \max \{ |\operatorname{Re} f(t)|, |\operatorname{Im} f(t)| \} dt \leq \max \left\{ \left| \int_a^b \operatorname{Re} f(t) dt \right|, \left| \int_a^b \operatorname{Im} f(t) dt \right| \right\}.$$

The equality holds in (12.3) if and only if both


$$\gamma \int_a^b \operatorname{Re} f(t) dt - \delta \int_a^b \operatorname{Im} f(t) dt = r \int_a^b \max \{ |\operatorname{Re} f(t)|, |\operatorname{Im} f(t)| \} dt$$

and

$$\gamma \int_a^b \operatorname{Re} f(t) dt - \delta \int_a^b \operatorname{Im} f(t) dt = \max \left\{ \left| \int_a^b \operatorname{Re} f(t) dt \right|, \left| \int_a^b \operatorname{Im} f(t) dt \right| \right\}.$$

Now, consider the Banach space $(\mathbb{C}, |\cdot|_{2p})$ with $p \geq 1$. Let $F : \mathbb{C} \rightarrow \mathbb{C}$, $F(z) = cz$ with $|c| = 2^{\frac{1}{2p}-\frac{1}{2}}$ ($p \geq 1$). Obviously, F is linear and by Hölder's inequality

$$\frac{|F(z)|}{|z|_{2p}} = \frac{2^{\frac{1}{2p}-\frac{1}{2}} \sqrt{|\operatorname{Re} z|^2 + |\operatorname{Im} z|^2}}{(|\operatorname{Re} z|^{2p} + |\operatorname{Im} z|^{2p})^{\frac{1}{2p}}} \leq 1.$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 92 of 99

Since, for $z_0 = 1 + i$ we have $|F(z_0)| = 2^{\frac{1}{p}}$, $|z_0|_{2p} = 2^{\frac{1}{2p}}$ ($p \geq 1$), hence

$$\|F\|_{2p} := \sup_{z \neq 0} \frac{|F(z)|}{|z|_{2p}} = 1,$$

showing that F is a bounded linear functional of unit norm on $(\mathbb{C}, |\cdot|_{2p})$, ($p \geq 1$). Therefore on using Theorem 10.1, we may state the following result.

Proposition 12.3. Let $\varphi, \phi \in \mathbb{R}$ with $\varphi^2 + \phi^2 = 2^{\frac{1}{2p}-\frac{1}{2}}$ ($p \geq 1$), $f : [a, b] \rightarrow \mathbb{C}$ be a Lebesgue integrable function on $[a, b]$ and $r \geq 0$ such that


$$r \left[|\operatorname{Re} f(t)|^{2p} + |\operatorname{Im} f(t)|^{2p} \right]^{\frac{1}{2p}} \leq \varphi \operatorname{Re} f(t) - \phi \operatorname{Im} f(t)$$

for a.e. $t \in [a, b]$, then

$$(12.4) \quad r \int_a^b \left[|\operatorname{Re} f(t)|^{2p} + |\operatorname{Im} f(t)|^{2p} \right]^{\frac{1}{2p}} dt \\ \leq \left[\left| \int_a^b \operatorname{Re} f(t) dt \right|^{2p} + \left| \int_a^b \operatorname{Im} f(t) dt \right|^{2p} \right]^{\frac{1}{2p}}, \quad (p \geq 1)$$

where equality holds in (12.4) if and only if both

$$\varphi \int_a^b \operatorname{Re} f(t) dt - \phi \int_a^b \operatorname{Im} f(t) dt = r \int_a^b \left[|\operatorname{Re} f(t)|^{2p} + |\operatorname{Im} f(t)|^{2p} \right]^{\frac{1}{2p}} dt$$



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 93 of 99

and

$$\begin{aligned} \varphi \int_a^b \operatorname{Re} f(t) dt - \psi \int_a^b \operatorname{Im} f(t) dt \\ = \left[\left| \int_a^b \operatorname{Re} f(t) dt \right|^{2p} + \left| \int_a^b \operatorname{Im} f(t) dt \right|^{2p} \right]^{\frac{1}{2p}}. \end{aligned}$$

Remark 14. If $p = 1$ above, and

$$r |f(t)| \leq \varphi \operatorname{Re} f(t) - \psi \operatorname{Im} f(t) \quad \text{for a.e. } t \in [a, b],$$

provided $\varphi, \psi \in \mathbb{R}$ and $\varphi^2 + \psi^2 = 1, r \geq 0$, then we have a reverse of the classical continuous triangle inequality for modulus:

$$r \int_a^b |f(t)| dt \leq \left| \int_a^b f(t) dt \right|,$$


with equality iff

$$\varphi \int_a^b \operatorname{Re} f(t) dt - \psi \int_a^b \operatorname{Im} f(t) dt = r \int_a^b |f(t)| dt$$

and

$$\varphi \int_a^b \operatorname{Re} f(t) dt - \psi \int_a^b \operatorname{Im} f(t) dt = \left| \int_a^b f(t) dt \right|.$$

If we apply Theorem 11.1, then, in a similar manner we can prove the following result for complex-valued functions.



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 94 of 99

Proposition 12.4. Let $\alpha, \beta \in \mathbb{R}$ with $\alpha^2 + \beta^2 = 1$, $f, k : [a, b] \rightarrow \mathbb{C}$ Lebesgue integrable functions such that

$$|\operatorname{Re} f(t)| + |\operatorname{Im} f(t)| \leq \alpha \operatorname{Re} f(t) - \beta \operatorname{Im} f(t) + k(t)$$

for a.e. $t \in [a, b]$. Then

$$(0 \leq) \int_a^b |\operatorname{Re} f(t)| dt + \int_a^b |\operatorname{Im} f(t)| dt - \left[\left| \int_a^b \operatorname{Re} f(t) dt \right| + \left| \int_a^b \operatorname{Im} f(t) dt \right| \right] \leq \int_a^b k(t) dt.$$

Applying Theorem 11.1, for $(\mathbb{C}, |\cdot|_\infty)$ we may state:


Proposition 12.5. Let $\gamma, \delta \in \mathbb{R}$ with $\gamma^2 + \delta^2 = \frac{1}{2}$, $f, k : [a, b] \rightarrow \mathbb{C}$ Lebesgue integrable functions on $[a, b]$ such that

$$\max \{|\operatorname{Re} f(t)|, |\operatorname{Im} f(t)|\} \leq \gamma \operatorname{Re} f(t) - \delta \operatorname{Im} f(t) + k(t)$$

for a.e. $t \in [a, b]$. Then

$$(0 \leq) \int_a^b \max \{|\operatorname{Re} f(t)|, |\operatorname{Im} f(t)|\} dt - \max \left\{ \left| \int_a^b \operatorname{Re} f(t) dt \right|, \left| \int_a^b \operatorname{Im} f(t) dt \right| \right\} \leq \int_a^b k(t) dt.$$

Finally, utilising Theorem 11.1, for $(\mathbb{C}, |\cdot|_{2p})$ with $p \geq 1$, we may state that:



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 95 of 99

Proposition 12.6. Let $\varphi, \phi \in \mathbb{R}$ with $\varphi^2 + \phi^2 = 2^{\frac{1}{2p}-\frac{1}{2}}$ ($p \geq 1$), $f, k : [a, b] \rightarrow \mathbb{C}$ be Lebesgue integrable functions such that

$$[|\operatorname{Re} f(t)|^{2p} + |\operatorname{Im} f(t)|^{2p}]^{\frac{1}{2p}} \leq \varphi \operatorname{Re} f(t) - \phi \operatorname{Im} f(t) + k(t)$$

for a.e. $t \in [a, b]$. Then


$$(0 \leq) \int_a^b [|\operatorname{Re} f(t)|^{2p} + |\operatorname{Im} f(t)|^{2p}]^{\frac{1}{2p}} dt - \left[\left| \int_a^b \operatorname{Re} f(t) dt \right|^{2p} + \left| \int_a^b \operatorname{Im} f(t) dt \right|^{2p} \right]^{\frac{1}{2p}} \leq \int_a^b k(t) dt.$$

Remark 15. If $p = 1$ in the above proposition, then, from

$$|f(t)| \leq \varphi \operatorname{Re} f(t) - \psi \operatorname{Im} f(t) + k(t) \quad \text{for a.e. } t \in [a, b],$$

provided $\varphi, \psi \in \mathbb{R}$ and $\varphi^2 + \psi^2 = 1$, we have the additive reverse of the classical continuous triangle inequality

$$(0 \leq) \int_a^b |f(t)| dt - \left| \int_a^b f(t) dt \right| \leq \int_a^b k(t) dt.$$



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back


Close

Quit

Page 96 of 99

References

- [1] E. BERKSON, Some types of Banach spaces, Hermitian systems and Bade functionals, *Trans. Amer. Math. Soc.*, **106** (1965), 376–385.
- [2] J.B. DIAZ AND F.T. METCALF, A complementary triangle inequality in Hilbert and Banach spaces, *Proc. Amer. Math. Soc.*, **17**(1) (1966), 88–97.
- [3] S.S. DRAGOMIR, *Semi-Inner Products and Applications*, Nova Science Publishers Inc., New York, 2004, pp. 222.
- [4] S.S. DRAGOMIR, *Advances in Inequalities of the Schwarz, Grüss and Bessel Type in Inner Product Spaces*, Nova Science Publishers Inc., New York, 2005, pp. 249.
- [5] S.S. DRAGOMIR, A reverse of the generalised triangle inequality in normed spaces and applications, *RGMIA Res. Rep. Coll.*, **7**(2004), Supplement, Article 15. [ONLINE: [http://rgmia.vu.edu.au/v7\(E\).html](http://rgmia.vu.edu.au/v7(E).html)].
- [6] S.S. DRAGOMIR, Additive reverses of the generalised triangle inequality in normed spaces, *RGMIA Res. Rep. Coll.*, **7**(2004), Supplement, Article 17. [ONLINE: [http://rgmia.vu.edu.edu.au/v7\(E\).html](http://rgmia.vu.edu.au/v7(E).html)].
- [7] S.S. DRAGOMIR, Additive reverses of the continuous triangle inequality for Bochner integral of vector-valued functions in Hilbert spaces, *RGMIA Res. Rep. Coll.*, **7**(2004), Supplement, Article 12. [ONLINE: [http://rgmia.vu.edu.au/v7\(E\).html](http://rgmia.vu.edu.au/v7(E).html)].



Reverses of the Triangle Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents




Go Back

Close

Quit

Page 97 of 99



Reverses of the Triangle
Inequality in Banach Spaces

S.S. Dragomir

Title Page

Contents



Go Back


Close

Quit

Page 98 of 99

- [8] S.S. DRAGOMIR, Additive reverses of the continuous triangle inequality for Bochner integral of vector-valued functions in Banach spaces, *RGMIA Res. Rep. Coll.*, **7**(2004), Supplement, Article 16[ONLINE: [http://rgmia.vu.edu.au/v7\(E\).html](http://rgmia.vu.edu.au/v7(E).html)].
- [9] S.S. DRAGOMIR, Quadratic reverses of the continuous triangle inequality for Bochner integral of vector-valued functions in Hilbert spaces, *RGMIA Res. Rep. Coll.*, **7**(2004), Supplement, Article 10. [ONLINE: [http://rgmia.vu.edu.au/v7\(E\).html](http://rgmia.vu.edu.au/v7(E).html)].
- [10] S.S. DRAGOMIR, Reverses of the continuous triangle inequality for Bochner integral of vector-valued functions in Hilbert spaces, *RGMIA Res. Rep. Coll.*, **7**(2004), Supplement, Article 11. [ONLINE: [http://rgmia.vu.edu.au/v7\(E\).html](http://rgmia.vu.edu.au/v7(E).html)].
- [11] S.S. DRAGOMIR, Reverses of the continuous triangle inequality for Bochner integral of vector-valued functions in Banach spaces, *RGMIA Res. Rep. Coll.*, **7**(2004), Supplement, Article 14 [ONLINE: [http://rgmia.vu.edu.au/v7\(E\).html](http://rgmia.vu.edu.au/v7(E).html)].
- [12] S.S. DRAGOMIR, Reverses of the triangle inequality in inner product spaces, *RGMIA Res. Rep. Coll.*, **7**(2004), Supplement, Article 7. [ONLINE: [http://rgmia.vu.edu.au/v7\(E\).html](http://rgmia.vu.edu.au/v7(E).html)].
- [13] S. GUDDER AND D. STRAWTHER, Strictly convex normed linear spaces, *Proc. Amer. Math. Soc.*, **59**(2) (1976), 263–267.
- [14] J. KARAMATA, *Stieltjes Integral, Theory and Practice* (Serbo-Croatian) SANU, Posebna Izdonija, 154, Beograd, 1949.

- [15] S.M. KHALEELULA, On Diaz-Metcalf's complementary triangle inequality, *Kyungpook Math. J.*, **15** (1975), 9–11.
- [16] S. KUREPA, On an inequality, *Glasnik Math.*, **3**(23)(1968), 193–196.
- [17] G. LUMER, Semi-inner product spaces, *Trans. Amer. Math. Soc.*, **100** (1961), 29–43.
- [18] M. MARDEN, The Geometry of the Zeros of a Polynomial in a Complex Variable, *Amer. Math. Soc. Math. Surveys*, **3**, New York, 1949.
- [19] P.M. MILIČIĆ, On a complementary inequality of the triangle inequality (French), *Mat. Vesnik* **41**(2) (1989), 83–88.
- [20] D.S. MITRINOVIĆ, J.E. PEČARIĆ AND A.M. FINK, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht/Boston/London, 1993.
- [21] B. NATH, On a generalisation of semi-inner product spaces, *Math. J. Okayama Univ.*, **15**(10) (1971), 1–6.
- [22] M. PETROVICH, Module d'une somme, *L'Enseignement Mathématique*, **19** (1917), 53–56.
- [23] H.S. WILF, Some applications of the inequality of arithmetic and geometric means to polynomial equations, *Proceedings Amer. Math. Soc.*, **14** (1963), 263–265.



**Reverses of the Triangle
Inequality in Banach Spaces**

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 99 of 99